



Contents lists available at ScienceDirect

## Trauma Case Reports

journal homepage: [www.elsevier.com/locate/tcr](http://www.elsevier.com/locate/tcr)



### Case Report

# Blunt pharyngeal trauma detected by ultrasound sonography: A case report

Tadao Kubota \*, Mayu Shimaguchi, Hideki Katagiri, Ken Mizokami

Todaijima 3-4-32, Urayasu, Chiba 279-0001, Japan

#### ARTICLE INFO

##### Article history:

Accepted 28 May 2016

Available online 1 July 2016

##### Keywords:

Blunt neck trauma

Pharyngeal perforation

Ultrasound sonography

#### ABSTRACT

Pharyngeal perforation related to blunt neck trauma is a rare clinical entity. Here in, we report a case of pharyngeal perforation secondary to minor blunt neck trauma. A 46 year old female was brought to our emergency room with neck pain. She fell down and hit her neck directly to the edge of a bed. There was no crepitation in physical examination. Neck ultrasound showed a small amount of air in her deep neck space. Followed CT and nasopharyngeal scope supported the presence of pharyngeal perforation. So emergency exploration was done. We found a laceration of hypopharynx and it was repaired. She could discharged without any complication on day 7. The indirect finding of pharyngeal perforation is subcutaneous emphysema. However if the air is localized only in deep cervical space, physical assessment is difficult. Although CT scan is potent modality to find air in the deep organs, ultrasound may be alternatives. Emphasis is based on the suspicion that minor blunt neck trauma may cause pharyngeal perforation.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Background

Blunt pharyngeal perforation is rare. It is under 2% of all pharyngeal lesions [1]. Most of them occurred due to high energy injury such as motor vehicle accidents [2], so surgically important pharyngeal injury secondary to minor trauma is extremely rare. We report a case of pharyngeal perforation related to low energy injury. Delay to diagnosis of this condition may lead to severe retro-mediastinal infection which needs long time hospitalization or sometimes causes death. Early diagnosis and surgical repair are critical. In our case, we detected suspicious pharyngeal perforation using ultrasound sonography.

\* Corresponding author. Tel.: +81 47 351 3101; fax: +81 47 352 6237.

E-mail address: [tadaokubota@mac.com](mailto:tadaokubota@mac.com) (T. Kubota).

## Case report

A 46 year old female was referred from the emergency department with a neck pain and difficulty to swallow. According to a witness, she fell down in her room and hit her neck directly to the edge of a bed. She had mental retardation and epilepsy. Her mother said that she often fell down because of uncontrolled general convulsion. There were mixed old bruises and new ones in her face. She had already lost all her front teeth. Primary survey was done with no abnormality. In the secondary survey, her neck was swollen with a fresh bruise on the left side (Fig. 1). On physical examination there was localized tenderness on her neck, crepitation was unclear. She was unable to swallow saliva at all, so she constantly spat out bloody saliva. We thought she might have pharyngeal injury. At first, we had tried to see her pharyngeal mucosa using nasopharyngeal scope to get a direct finding. However it failed with poor examination. The left side of the pharyngeal mucosa was bleeding and edematous, so we could suspect the presence of a pharyngeal injury but couldn't get an evidence of the perforation. Next, we examined her neck using ultrasound sonography. It revealed some hyperechoic spots which meant extraluminal air in the deep neck space (Fig. 2). Air was distributed mainly around the pharynx and cervical esophagus rather than larynx and trachea. Finally neck CT scan was done, it showed deep cervical emphysema which was definitive evidence of the perforation in either the air way or alimentary tract (Fig. 3). There were no fractures of the cartilages. According to these findings, she should have symptomatic pharyngeal perforation, so we decided to do emergent neck exploration. Under general anesthesia, a standard neck oblique incision was made. Larynx and trachea were intact. We found a small amount of hematoma and a laceration, 1 cm in diameter on the left inferior pharyngeal muscle (Fig. 4). The laceration was closed by interrupted suture with polyfilament absorbable stitch. The retrocervical space was drained with a closed suction system. The day after surgery, she could swallow saliva. One week after surgery she discharged without any complications.

## Discussion

The most common cause of pharyngeal injury is iatrogenic, especially after difficult endotracheal intubation [3]. Surgically repaired pharyngeal perforation secondary to blunt trauma is rare [2,4]. She must be unconscious at that time because of general convulsion, so her body weight was concentrated on her neck. The force of blow might be relatively high. Even so, isolated blunt neck trauma with pharyngeal perforation caused by low energy is extremely rare. When we encounter an isolated minor blunt neck trauma patient, we rarely think that patient may have pharyngeal perforation. In that situation, CT scan is not ordered routinely. It is easy for us to suspect it if we find crepitation around the neck of a patient. But if a patient is asymptomatic, most physicians tend to think that further examination is unnecessary.

On the other hand, delayed diagnosed pharyngeal perforation is known as a source of severe retropharyngeal or mediastinal infection which sometimes needs aggressive drainage and long time intensive care [5]. Its mortality and morbidity rates remain high. In abdominal trauma, ultrasound sonography is



**Fig. 1.** Neck bruise. An arrow showed a bruise on left side of the neck.

Download English Version:

<https://daneshyari.com/en/article/4100509>

Download Persian Version:

<https://daneshyari.com/article/4100509>

[Daneshyari.com](https://daneshyari.com)