




Available online at  
**ScienceDirect**  
[www.sciencedirect.com](http://www.sciencedirect.com)

Elsevier Masson France  
**EM|consulte**  
[www.em-consulte.com/en](http://www.em-consulte.com/en)



Original article

# Criteria for admission of odontogenic infections at high risk of deep neck space infection



N. Alotaibi\*, L. Cloutier, E. Khaldoun, E. Bois, M. Chirat, D. Salvan

Service d'ORL et de chirurgie cervico-maxillo-faciale, centre hospitalier Sud Francilien, 116, boulevard Jean-Jaurès, 91100 Corbeil-Essonnes, France

## ARTICLE INFO

### Keywords:

Odontogenic infection  
 Dental abscess  
 Deep neck space infection  
 Criteria for admission

## ABSTRACT

**Objectives:** Many patients with odontogenic infections are hospitalised because of the risk of deep neck space infection. The objective of this study was to identify risk factors allowing more reliable selection of patients requiring hospitalisation for both specialists and emergency physicians.

**Material and methods:** This retrospective study was based on a cohort of 97 patients hospitalised for odontogenic infection in the Department of Otorhinolaryngology and Head and Neck Surgery of Centre hospitalier Sud Francilien, Île-de-France, from January 2008 to June 2012.

**Results:** The majority of patients presented with dental abscess (66 patients; 68%). Nineteen patients (20%) presented with deep neck space infection. The frequency of deep neck space infection was significantly higher in patients with mandibular odontogenic infection (16/55 patients (29%) than in those with maxillary odontogenic infection (3/42 (7%);  $P \leq 0.009$ ). The incidence of deep neck space infection was significantly higher in patients with dental abscess (17/66, (26%) than in those without dental abscess (2/31 (6%);  $P \leq 0.03$ ).

**Conclusion:** In addition to the well-known classical criteria (fever, neck swelling, dyspnoea, dysphagia, trismus, leukocytosis, elevated C reactive protein (CRP)), the criteria for admission for odontogenic infection should include mandibular odontogenic infection and/or the presence of dental abscess.

© 2015 Elsevier Masson SAS. All rights reserved.

## 1. Introduction

Deep neck space infections can spread along the fascial spaces of the head and neck, inducing life-threatening deep space infection associated with a high risk of complications (upper airway obstruction, mediastinitis, thoracic empyema, pericarditis, septic shock) [1–4]. Although the incidence of these infections has decreased considerably as a result of modern antibiotics and improvement of oral hygiene [5], these infections continue to be a source of severe morbidity and high mortality [6–8]. The portals of entry most commonly encountered are odontogenic infections, but tonsillar abscess and foreign bodies may also be responsible [9,10]. Odontogenic infections can spread and evolve very rapidly from a localized infection to deep neck space infection, corresponding to extensive necrotizing fasciitis requiring urgent medical and surgical management [11].

Odontogenic infections are a common medical and surgical emergency in our department (Centre hospitalier Sud Francilien Department of Otorhinolaryngology and Head and Neck Surgery).

Some patients are hospitalised due to an excess of caution, while others are treated on an outpatient basis when they should have been admitted to hospital. However, to our knowledge, no flow chart useful for the management of these odontogenic infections has been published in the literature. Criteria for admission can vary according to each practitioner's experience. The definition of objective criteria for admission of cases of odontogenic infection is important to improve patient management and to limit the risk of deep infections.


The objective of this retrospective study was to identify criteria of severity of odontogenic infections on the basis of identified risk factors for complications, such as the maxillary or mandibular site of infection [12] and the presence or absence of dental abscess.

## 2. Patients and methods

This retrospective study analysed patients hospitalised for odontogenic infection in the Department of Otorhinolaryngology and Head and Neck Surgery, Centre Hospitalier Sud Francilien, Île-de-France, from January 2008 to June 2012. These patients presented clinical signs such as dyspnoea, stridor, dysphagia, odynophagia, inflammatory neck swelling, fever, painful cervical lymphadenopathy, trismus, soft tissue crepitations (Fig. 1) and

\* Corresponding author.

E-mail address: [notaibi@alfaisal.edu](mailto:notaibi@alfaisal.edu) (N. Alotaibi).



**Fig. 1.** Contrast-enhanced head and neck CT, axial section through the thyroid cartilage in a patient with deep neck space infection. Air bubbles may be observed in the soft tissues.

were divided into 2 groups: group 1 comprising maxillary odontogenic infections and group 2 comprising mandibular odontogenic infections.

Clinical, laboratory, radiological data as well as outcome and complications were analysed. Statistical analysis was performed with SPSS statistical software (SPSS for Windows, Version 16.0. Chicago, SPSS Inc.). Results were considered to be statistically significant for  $P \leq 0.05$ .


### 2.1. Therapeutic management

After performing a laboratory work-up (complete blood count [CRP]) and dental panoramic radiography, treatment of hospitalised patients consisted of empirical antibiotic therapy with a combination of intravenous amoxicillin and clavulanic acid. Patients allergic to amoxicillin received clindamycin. Therapeutic management also comprised incision and drainage of the abscess under local anaesthesia whenever indicated and extraction of the infected teeth. Deep neck abscesses were drained surgically and broader antibiotic therapy was prescribed with the addition of ceftriaxone 2g daily and metronidazole 1500 mg daily (Fig. 2). In the absence of clinical improvement or in the presence of signs of extensive deep neck space infection, contrast-enhanced head and neck CT was performed. All hospitalised patients were evaluated by a dentist during the first 48 hours after admission for dental treatment. Patients were discharged according to the local and systemic clinical improvement.

### 3. Results

Two hundred eighty one patients attended the emergency department with dental symptoms and signs during this period and 97 patients (35%), comprising 35 women and 62 men, were hospitalised. The mean age of hospitalised patients was 41 years (range: 13 to 92) and the mean length of hospital stay was 4.6 days (Table 1).

All patients were treated by intravenous antibiotics. Dental extraction was performed in 73 patients (maxillary teeth in 30 cases, i.e. 31% of hospitalised patients, and mandibular teeth in 43



**Fig. 2.** Clinical appearance of deep neck space infection secondary to odontogenic infection after surgical drainage.

**Table 1**  
Study population and characteristics.

| Characteristics                       | Values          | Percentage |
|---------------------------------------|-----------------|------------|
| Number of patients                    | 97              |            |
| Age (years)                           | 41 ± 18 (13–92) |            |
| Gender                                |                 |            |
| Male                                  | 62              | 64         |
| Female                                | 35              | 36         |
| Anatomical site                       |                 |            |
| Mandibular                            | 55              | 57         |
| Maxillary                             | 42              | 43         |
| Dental abscess with surgical drainage |                 |            |
| Total number                          | 19/97           | 20         |
| Mandibular                            | 16/55           | 29         |
| Maxillary                             | 3/42            | 7          |
| Length of hospital stay (days)        |                 |            |
| Mean                                  | 5 ± 8           |            |
| Mandibular                            | 6 ± 11          |            |
| Maxillary                             | 3 ± 3           |            |
| P value                               | 0.02            |            |
| Treatment in hospital                 |                 |            |
| Antibiotics                           | 97              | 100        |
| Dental extraction                     | 73              | 75         |

Data expressed as number, mean ± standard deviation (range), percentage in parentheses (%), or number per subgroup (%).

cases, i.e. 44%) (Table 1). Eighty-five patients (88%) had a dental abscess, requiring local drainage in 66 cases and surgical drainage under general anaesthesia in 19 cases. Three patients (5%) presented extensive deep neck space infection, including 2 cases of mediastinitis treated by bilateral neck incision with drainage, repeated dressings and intensive care (Table 1). Of the 19 patients requiring surgical drainage of a dental abscess under general anaesthesia, the odontogenic infection arose in the mandible in 16 cases and in the maxilla in 3 cases.

#### 3.1. Criteria of severity according to the site of infection

Group 2 had a significantly higher risk of dental abscess requiring surgical drainage ( $P=0.009$ ) and a significantly longer hospital stay ( $P=0.01$ ). All patients admitted to the intensive care unit with deep neck space infection belonged to group 2.

Download English Version:

<https://daneshyari.com/en/article/4109984>

Download Persian Version:

<https://daneshyari.com/article/4109984>

[Daneshyari.com](https://daneshyari.com)