

HOSTED BY




ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/ijpam>



INSTRUCTIVE CASE

Childhood systemic sarcoid-like necrotizing granulomatous disease: Another piece of the puzzle



Ashwaq AlE'ed ^a, Hadeel AlManea ^b, Sulaiman M. Al-Mayouf ^{a,*}

^a Section of Rheumatology, Department of Pediatrics, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia

^b Pathology and Laboratory Medicine, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia

Received 25 July 2013; accepted 29 October 2013
Available online 7 October 2014

KEYWORDS

Childhood;
Granulomatous
disease;
Sarcoidosis

Abstract Necrotizing sarcoid-like granulomatous disease is a disorder of unknown etiology that is rarely described in childhood; it affects various organs and has a diverse clinical course. Here, we report the case of a girl with multisystem recalcitrant necrotizing sarcoid-like granulomatous disease. The diagnosis is based on histopathology and the exclusion of other disease entities known to cause granulomatous disease.

Copyright © 2014, King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

1. Introduction

Granuloma formation is a special type of inflammation that can occur in a wide variety of diseases, both infectious and non-infectious. However, the offending agent is often unknown to be sarcoidosis or granulomatous angiitis. Diagnosis relies on clinical history, laboratory findings demonstrating an inflammatory process and histological examination of the suspected lesions. It is necessary to

differentiate inflammatory granulomatous entities from tuberculosis and other infections or malignancies [1]. The rarity of non-infectious granulomatous disorders including necrotizing sarcoid-like granulomatosis and the scarcity of awareness may delay the appropriate diagnosis and treatment [2,3].

We describe a girl with recalcitrant necrotizing sarcoid-like granulomatous disease; the diagnosis is based on histopathology and the exclusion of other disease entities known to cause granulomatous disease.

2. Case report

A 13-year-old Pakistani girl was referred to our hospital with a three-year history of recurrent fever associated with generalized body weakness and diffuse musculoskeletal pain. She had a history of weight loss and recurrent

* Corresponding author. Alfaisal University, Rheumatology, Department of Pediatrics, MBC-58, King Faisal Specialist Hospital and Research Centre, PO Box 3354, Riyadh 11211, Saudi Arabia. Tel.: +966 11 442 7761; fax: +966 11 442 7784.

E-mail address: mayouf@kfshrc.edu.sa (S.M. Al-Mayouf).

Peer review under responsibility of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia.

abdominal pain without a change in bowel habits. She was observed and evaluated in different hospitals; she had persistently high white blood cell (WBC) and platelet counts as well as a high erythrocyte sedimentation rate (ESR). Other workup including bone marrow aspiration (BMA) and cultures were negative. However, Bartonella IgG levels were elevated. The initial clinical diagnosis was systemic onset juvenile idiopathic arthritis. Accordingly, she was treated with naproxen (250 mg twice daily), a high dose of prednisone (2 mg/kg/day), methotrexate (1 mg/kg/week) and tocilizumab (IL6 inhibitor). Because partial improvement was observed, tocilizumab was switched to anakinra (IL1 inhibitor) (100 mg daily). However, her symptoms recurred with bloody diarrhea. Further evaluation showed negative stool analysis and culture for any infectious disease. After undergoing upper and lower endoscopy, it was determined that there was no ulceration, and histopathology showed mild non-specific inflammation, which is not consistent with inflammatory bowel disease. An abdominal ultrasound and magnetic resonance imaging showed para-aortic lymphadenopathy, hepatic and splenic lesions. A lymph node biopsy revealed necrotizing granulomatous lymphadenitis with negative culture for mycobacterial and fungal infection. The family decided to use a mixed herbal medicine that was a mixture of NSAIDs. There was a family history of tuberculosis (TB).

Upon arrival to our hospital, she was febrile (38.9 °C) and appeared unwell with the same complaints mentioned above. She had scattered lymphadenopathy, decreased air entry bilaterally with crepitation, and hepatosplenomegaly. She was determined to have proximal muscle weakness and localized thoracic vertebral tenderness. Another physical examination was unremarkable.

The clinical impression at that time was infection (specifically TB), Crohn's disease, chronic granulomatous disease and autoimmune disease (namely sarcoidosis); other diagnoses such as malignancy i.e., lymphoma were considered as well.

The laboratory findings revealed a white blood cell count of $19.4 \times 10^9/l$, hemoglobin level of 90 g/l, platelet level of $679 \times 10^9/l$, elevated ESR level (108 mm/h) and elevated C-reactive protein level (213 mg/l). Her liver enzymes were elevated (ALT 136 U/l and AST 214 U/l), Gamma-glutamyltransferase level was GGT 104 IU/l and albumin

level was 23 g/l. However, her renal function and muscle enzymes were within normal range. Her complement level (C3, C4) was normal, and her autoantibodies (antinuclear antibodies and antineutrophil cytoplasmic antibodies) were negative.

A chest computed tomography (CT) scan showed pleural effusion with normal lung parenchymal tissue. However, there were enlarged bilateral supraclavicular and paratracheal lymph nodes. A tuberculin skin test (TST) was negative along with a normal QuantIFERON TB gold test. The angiotensin-converting enzymes were borderline at 54 U/l (normal range <52). She had normal serum immunoglobulin levels, T and B-lymphocytes markers as well as a normal oxidative burst assay.

She completed a capsule endoscopy, which revealed a normal intestine.

A thoracolumbar X-ray showed multiple wedge compression vertebral fractures and osteopenia, which were confirmed by bone densitometry (Z-score <-2).

A BMA and biopsy were negative for malignancy, granulomas or infections. An abdominal CT scan confirmed multiple enlarged mesenteric and retroperitoneal lymph nodes with multiple focal hypodense splenic lesions (Fig. 1). The histopathology of lymph node and splenic biopsy revealed necrotizing granulomatous inflammation (Fig. 2). An acid-fast bacilli (AFB) stain was negative in tissue culture. A nucleic acid amplification test using strand displacement amplification was negative for AFB. The final report of tissue culture including mycobacterial and fungal was negative. A serology (IgG and IgM) and polymerase chain reaction (PCR) test for Bartonella was negative.

We proposed a diagnosis of systemic sarcoid-like necrotizing granulomatous disease especially with the presence of necrotizing granulomatous inflammation and inconclusive extensive workup, which failed to suggest an alternative diagnoses. Accordingly, we started intravenous methyl prednisolone (30 mg/kg) weekly for four doses, oral prednisone (0.5 mg/kg/day), azathioprine (2 mg/kg/day), calcium and vitamin D supplements and cyclic pamidronate (every three months) treatment. Recently, infliximab and cyclosporine (4 mg/kg) were initiated, and she received four doses of infliximab (5 mg/kg). Fortunately, she experienced significant clinical improvement, and her inflammatory markers improved as well.




Figure 1 Abdominal CT scan that showed multiple enlarged mesenteric and retroperitoneal lymph nodes (1A) and a bulky spleen (measured to be 12.7 cm) with multiple focal hypodense lesions; the largest was 2.2 cm in its transverse dimension (1B).

Download English Version:

<https://daneshyari.com/en/article/4153740>

Download Persian Version:

<https://daneshyari.com/article/4153740>

[Daneshyari.com](https://daneshyari.com)