



A novel minimal residual disease model of neuroblastoma in mice

Jeremy R. Jackson^a, Youngleem Kim^a, Robert C. Seeger^b, Eugene S. Kim^{a,*}

^a Division of Pediatric Surgery, Department of Surgery, Keck School of Medicine, University of Southern California, Los Angeles, CA

^b Department of Pediatrics, Keck School of Medicine, University of Southern California, Los Angeles, CA



ARTICLE INFO

Article history:

Received 21 February 2016

Accepted 26 February 2016

Key words:

Neuroblastoma

Minimal residual disease

Mouse

Model

Resection

ABSTRACT

Introduction: Patients with high-risk neuroblastoma rarely succumb to their primary tumor but rather from relapsed metastatic disease after surgery. We, therefore, sought to create an *in vivo* model of minimal residual disease (MRD), which clinically replicates tumor recurrence and metastasis after surgical resection.

Methods: Neuroblastoma cell lines CHLA-255, CHLA-136, and SH-SY5Y were used. After establishing orthotopic xenografts, mice were divided into control tumor group (sham operation at 14 days) and tumor resection group (resection at 14 days). Mice were monitored by bioluminescent imaging and sacrificed when institutional criteria for euthanasia were met.

Results: In the CHLA-255 and CHLA-136 cell lines, mice experienced significantly longer survival following tumor resection ($p < 0.007$). There was no survival benefit seen in the SH-SY5Y cell line ($p = 0.29$). Bioluminescent imaging demonstrated metastatic disease in 100% of all tumor resection mice and varying rates of metastasis in control mice (4 of 5 CHLA-255, 2 of 4 CHLA-136, and 7 of 7 SH-SY5Y).

Conclusion: In this study, we describe a novel neuroblastoma model of MRD in mice. This MRD model serves as an innovative means to test preclinical therapies as well as elucidate mechanisms of metastatic disease in experimental neuroblastoma.

© 2016 Elsevier Inc. All rights reserved.

Neuroblastoma is a heterogeneous, embryonal malignancy of early childhood that is derived from primitive sympathetic ganglion cells. While this cancer can arise anywhere along the sympathetic nerve chain, the majority of these tumors are adrenal in origin. It is the most common pediatric extracranial solid tumor and represents a disproportionately high 15% of all cancer related deaths in children [1]. Despite 5-year survival rates of greater than 90% for low- and intermediate-risk neuroblastoma groups, the survival rate in children with high-risk neuroblastoma remains 40%–50% [2,3]. High-risk neuroblastoma is generally defined by patient age older than 12 months, evidence of metastatic disease, tumor MYCN gene amplification, and unfavorable pathologic findings. While nearly 80% of high-risk patients will achieve remission through high-dose chemotherapy, surgery, radiation and stem cell transplantation, the majority of patients will eventually succumb to recurrent metastatic disease. Relapse from the presence of minimal residual disease continues to be the most significant barrier to improving the prognosis of those with high-risk tumors.

In research, there are currently three major preclinical mouse models of neuroblastoma: heterotopic, orthotopic and transgenic models. The heterotopic model typically involves the subcutaneous injection of neuroblastoma cells into the flanks of mice [4]. Although this model is easy to perform, the subcutaneous space is not a clinically relevant site for neuroblastoma, and metastases rarely develop. A frequently used orthotopic model entails the injection of tumor cells directly into the subrenal

capsule. Despite being difficult to perform successfully in very young mice and requiring increased technical skill, it better simulates the micro-environment in which these tumors normally grow, and the resulting xenografts produce metastases to clinically relevant sites such as the bone marrow, bone, and liver [5,6]. Transgenic mouse models have also been extensively utilized. The TH-MYCN model, in particular, has an overexpression of the oncogene MYCN targeted to neural crest cells, which leads to spontaneous tumorigenesis in the thoracic cavity and adrenal glands of these mice [6,7]. Variances in tumor latency and challenges in breeding frequency make this model more difficult to utilize. However, current mouse models, while useful in evaluating treatment outcomes in the setting of a primary tumor, have provided little insight into the study of tumor recurrence and metastatic disease.

To this end, we developed a novel neuroblastoma model of minimal residual disease to evaluate the processes involved in tumor relapse and metastatic spread as well as their mechanistic origins. Herein, we investigate in several cell lines of human neuroblastoma the incidence of metastasis, recurrence of tumor, and the overall survival rate after surgical resection of orthotopic xenografts when compared to control mice with primary tumors.

1. Material and methods

1.1. Cell culture and reagents

Three human luciferase-expressing neuroblastoma cell lines (CHLA-255, CHLA-136, SH-SY5Y) were cultured and utilized. The CHLA-255 cell line is MYCN amplified and derived from the brain metastasis of a

* Corresponding author at: Division of Pediatric Surgery, Children's Hospital Los Angeles, 4650 Sunset Blvd, MS #100, Los Angeles, CA, 90027, USA. Tel.: +1 323 361 8332; fax: +1 323 361 3534.

E-mail address: eugenekim@chla.usc.edu (E.S. Kim).

patient. The CHLA-136 cell line is a particularly chemoresistant cell line with MYCN amplification and was established from the peripheral blood of a patient after chemotherapy and bone marrow transplantation. The SH-SY5Y cell line is non-MYCN amplified and a subclone of the SK-N-SH cell line, derived from a bone marrow biopsy of a patient with thoracic neuroblastoma. CHLA-255 and CHLA-136 were provided by Dr. Robert Seeger (Children's Hospital Los Angeles) and the SH-SY5Y cell line was a gift of Dr. Darrell Yamashiro (Columbia University). Briefly, cell lines were maintained in Iscove's Modified Dulbecco's Medium [CHLA-255 and CHLA-136] or RPMI-1640 Medium [SH-SY5Y]. All media were supplemented with 10% heat-inactivated FBS. On the day of injection, the cells were trypsinized using Puck's EDTA and centrifuged. The cells were washed with PBS twice and were twice counted using a hemacytometer. The cells were then resuspended in 100 μ l PBS per 1×10^6 cells and stored on ice.

Neuroblastoma cell lines expressing firefly luciferase were generated using a lentiviral transduction technique using standard protocols and a commercially available vector (pGL2-Control vector, Promega, Madison, WI). In brief, 5×10^6 293T cells (ATCC #11,268, Manassas, VA) were seeded on 10 cm cell culture dishes and incubated overnight. CaPO₄ was used for transfection of the luciferase vector followed by sodium butyrate induction about 17 h later. After 48 h of incubation, viral supernatant was ultracentrifuged, and the subsequent viral pellet was dissolved in PBS and stored in a -80°C freezer. Neuroblastoma cells, 2×10^5 , were incubated overnight in a 12 well plate. Lentiviral-luciferase was used to transduce the neuroblastoma cells for one to two cycles overnight in an incubator. Luciferase positive clones were identified by bioluminescent imaging and subsequently cultured using standard, cell-line specific tissue culture media.

1.2. Animal model

Six to eight week old inbred NOD SCID gamma (NSG) mice were used for all studies. Our NSG mouse colony was originally established from mice purchased from Jackson Laboratory and was housed in pathogen-free conditions. All in vivo experiments were approved by the Institutional Animal Care and Use Committee of Children's Hospital Los Angeles (IACUC protocol #363-14). Each mouse was implanted with 1×10^6 NB cells (CHLA-255, $n = 10$; CHLA-136, $n = 8$; SH-SY5Y, $n = 14$) into the left subrenal capsule as previously described [5]. In brief, the left flank of anesthetized NSG mice was prepared in a sterile fashion. A transverse left flank incision was performed and the muscle sharply divided. The left kidney was identified and exteriorized out of the wound, and one million neuroblastoma cells in 0.1 ml of PBS were injected into the kidney. The left kidney was placed back into the retroperitoneal space, and the muscle closed with a single 4-0 Vicryl stitch and the skin closed with a skin clip.

After 14 days, a complete resection of the xenograft was performed in half the mice (tumor resection group), while a sham surgery was performed on the other half (control tumor group). All mice were induced in an anesthetic chamber using 2%–2.5% inhaled isoflurane. The mice were transferred onto a sterile field and their left flank was prepared with betadine. Maintenance anesthesia was delivered through a nose cone with 2% isoflurane. A scalpel was used to reopen the previous left flank incision, and the muscle layer was sharply opened. For the control groups, the muscle layer was then reapproximated using 4-0 Vicryl and the skin was closed using a surgical clip. For the tumor resection mice, the left kidney was identified and any adhesions were bluntly dissected. The left kidney was then externalized and a 4-0 Vicryl free tie was used to ligate its hilum. The kidney was then excised. The muscle layer and skin were then closed in layers. Mice were monitored daily and were sacrificed when they met institutional criteria for euthanasia which included weakness/paralysis, seizures, inability to eat or drink, moribund state and dyspnea.

Mice underwent bioluminescent imaging to monitor tumor growth and metastatic disease (Xenogen IVIS Spectrum System, Caliper Life

Sciences). The incidence of metastasis was determined by the presence of a bioluminescent signal at the site of the femurs, liver and/or brain.

1.3. Statistical analysis

Survival data were expressed as mean \pm SD and were analyzed using the Kaplan–Meier method with significance being determined by log-rank test. $P < 0.05$ was considered significant.

2. Results

2.1. Minimal residual disease is found in mice after primary tumor resection

In the CHLA-136 group, from the initial 12 mice, 1 mouse died during the orthotopic tumor implantation secondary to anesthesia complications, and 3 mice subsequently died from hemorrhage during tumor resection. Refinement of techniques was made, and all mice in the CHLA-255 and SH-SY5Y groups survived both the orthotopic tumor cell implantation and the subsequent tumor resection surgical procedures.

During early postresection imaging (postoperative day 2), the CHLA-255 tumor resection group showed evidence of MRD in 4 of 5 mice, and at the time of late imaging prior to sacrifice, showed MRD in 5 of 5 mice (Fig. 1). At necropsy, recurrent tumor growth was found in 1 of 5 of the CHLA-255 tumor resection mice. In the CHLA-136 mice, early and late postresection imaging showed evidence of minimal residual disease in 4 of 4 tumor resection mice (Fig. 1). At sacrifice, 4 of 4 tumor resection mice developed recurrent tumors. In the SH-SY5Y group, on early postresection imaging, 6 of 7 tumor resection mice demonstrated evidence of minimal residual disease, and at the time of late imaging prior to sacrifice, showed MRD in 7 of 7 mice (Fig. 1). At necropsy, despite widely aggressive metastatic disease, there were no recurrent tumors in the tumor resection mice.

2.2. Increased survival after tumor resection in two of three cell lines

In the CHLA-255 group, mice that underwent tumor resection had a significantly increased mean survival of 47.4 ± 1.3 days compared to 33.8 ± 4.1 days for the control tumor group ($p = 0.003$) (Fig. 2a). Similarly, in the CHLA-136 group, mice that underwent tumor resection also had a significantly increased mean survival of 44 ± 3.2 days compared to 36 ± 2.7 days for the control tumor group ($p = 0.007$) (Fig. 2b). Interestingly, the SH-SY5Y mice showed no difference in mean survival following tumor resection (tumor resection group 27.6 ± 2.8 days vs. 29 ± 1.3 days control tumor group ($p = 0.29$) (Fig. 2c)).

2.3. All mice that undergo tumor resection develop metastatic disease

In the CHLA-255 cohort ($n = 10$), all mice in the tumor resection group (5 of 5) had evidence of metastasis to the brain, femur and/or liver at time of sacrifice, compared to 4 of 5 control tumor mice (Fig. 3). Specifically, all mice in the tumor resection group were found to have metastases to the liver; in the control group, 2 mice had metastases to the liver, 2 mice had normal appearing livers, and 1 mouse liver was too necrotic to determine. In the CHLA-136 group ($n = 8$), 4 of 4 tumor resection mice had evidence of metastasis to liver and/or femurs on the day of sacrifice, whereas only 2 of 4 control mice had metastasis (Fig. 3). All mice in the CHLA-136 tumor resection group demonstrated metastases to the liver, and in the control tumor group, only 2 mice demonstrated overt liver metastases. In the SH-SY5Y cohort ($n = 14$), at the time of sacrifice, all 7 of 7 tumor resection mice and 7 of 7 control tumor mice were found to have metastases to the liver on imaging as well as grossly (Fig. 3).

Download English Version:

<https://daneshyari.com/en/article/4154921>

Download Persian Version:

<https://daneshyari.com/article/4154921>

[Daneshyari.com](https://daneshyari.com)