



# Congenital esophagostenosis due to tracheobronchial remnant in infant: 3 cases report<sup>☆</sup>



Aloïse Sagna<sup>\*</sup>, Ibrahima Fall, Aïssata Ly, Mbaye Fall

## ARTICLE INFO

### Article history:

Received 5 January 2014

Received in revised form

2 March 2014

Accepted 5 March 2014

### Key words:

Stenosis

Esophageal

Tracheobronchial remnant

## ABSTRACT

Esophageal congenital stenosis due to tracheobronchial remnant is a rare deformation. Its scarcity incites us to report three observations of abnormal tracheal cartilage in the wall of the lower esophagus in infants (ages: 6; 12; 24 months). Authors present records of 3 children who underwent a radical procedure of surgery for esophageal congenital stenosis in Albert Royer's Pediatric Hospital from 2005 to 2009. Anatomopathological exam was performed to identify the cause of the stenosis and develop treatment recommendations. The study included two girls and one boy who presented vomiting, undernourishment and bad general state of health. The upper gastrointestinal radiological investigations established stenosis of distal esophagus with dilatation above. Endoscopy performed in one case showed not passable regular stenosis, located at 25 cm from dental arches. There was no abnormality of mucosal appearance above and dilation wasn't attempted for lack of pediatric dilator. Diagnosis of mega-esophagus was formulated and all the infants underwent esophageal distal resection with terminal end-to-end anastomosis, HIS corner repairing and anterior partial stomach fundoplication to prevent reflux. The postoperative follow up was good and the upper gastrointestinal radiological investigations were normal after one month. The anatomopathological exam revealed tracheobronchial remnant in all cases. Authors discuss about standpoint to adopt in front of esophageal congenital stenosis due to tracheobronchial remnant. We believe that radical surgery is essential for the removal of stenosis.

© 2014 The Authors. Published by Elsevier Inc. All rights reserved.

Esophageal congenital stenosis due to tracheobronchial remnant is defined as the abnormal presence of a ring of tracheal or bronchial cartilage in the wall of the esophagus. It is an uncommon cause of congenital esophagostenosis among infants, which is usually revealed by vomiting when introducing solid food in early childhood [1]. Differential diagnosis is to be done with achalasia of the cardiac, fibro-muscular stricture and secondary stenosis especially gastro-esophagus reflux. The presence of cartilaginous ring has been proved to fail dilation and surgery appears as the unique treatment likely to remove completely and permanently the obstruction [2]. We report on three cases of infants in a four year period.

## 1. Methods

We reviewed the medical records of 3 consecutive infants who were admitted to our hospital with the diagnosis of megaesophagus.

Gastrointestinal and esophagus Barium radiological investigation was performed in all cases. According to standard practice endoscopy examination is routinely required, but it was performed in one case as we had no easily available gastro-intestinal service support. The decision for surgical resection of the stenosis was made per-operatively and histological exam done on operative specimen. Satisfaction was based on recovery, vomit relief and weight gaining.

## 2. Results

### 2.1. Case 1

A 5-month 25-day-old female infant presents since she was 4-month-old, episodic early persistent vomiting. Then dehydration, severe undernutrition and moderate dyspnea were diagnosed following her hospitalization in pediatric unit with a weight of 3900 g in June, 28th 2004. On the 10th of July, face to persistent vomiting despite a well driven hydroelectrolytic resuscitation, an upper gastrointestinal radiological investigation was realized and showed stenosis of the distal quarter of the esophagus. On the 9th of September we performed gastrostomy of food supply with the aim of improving the nutritional state of the infant. The radical surgery decided 6 months later, on the 12th of January 2005,

<sup>☆</sup> This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

<sup>\*</sup> Corresponding author. Centre Hospitalier National d'Enfants Albert Royer, Dakar-Fann BP 25755, Senegal. Tel.: +221 777017241 (mobile); fax: +221 338258005, +338682154.

E-mail address: [alosagna@hotmail.com](mailto:alosagna@hotmail.com) (A. Sagna).



**Fig. 1.** Barium study.



**Fig. 2.** Operative findings.

The upper gastrointestinal radiological investigation was made on November 17th, showing a regular shrinkage of the terminal esophagus surmounted by a “socking” dilatation of the upstream segment. The radical surgery was made on March 17th, 2009 with simple postoperative consequences and a weight recovery of 11.800 kg after a recession of 19 months. The histological results of the operative specimen revealed tracheobronchial remnant ([Fig. 5](#)).

### 3. Discussion

The prevalence of esophagus congenital stenosis revolves around 1/25,000–1/50,000 births according to Japanese studies [[3](#)]. This very rare affection is associated nearly in one third of the cases


consisted of a resection of the stenosed esophagus with an end-to-end anastomosis followed by an anterior partial gastric fundoplication. Postoperative consequences were simple and the histology results revealed a tracheobronchial remnant.

#### 2.2. Case 2

A 9-month-old male infant without pathological history is admitted in pediatrics on September 17th, 2008 for early vomiting that runs since the age of 6 months in a context of moderate alteration of general state with an entrance weight of 8 kg. A correction of hydro-electrolytic disorders is made then an upper gastrointestinal radiological investigation is practiced on September 19th, 2008 showing a tight shrinkage of the terminal esophagus with a “socking” dilatation of the upstream segment ([Fig. 1](#)). Previously an endoscopy made on September 18th objectified a not passable, fibrous regular stenosis by the pediatric fiberscope located at 25 cm from dental arches; there was no abnormality of mucosal appearance above the stricture and dilation wasn't attempted for lack of pediatric dilator. The intervention is led by an upper umbilical median incision on December 24th, 2008 allowing resection of the terminal esophagus and end-to-end anastomosis with an anterior partial stomach fundoplication ([Figs. 2–4](#)). Postoperative consequences were simple and the hospital discharge was authorized after 8 days. The histological examination of the operative specimen found tracheobronchial remnant islet. After 24 months of follow-up the child is healthy and weights 12 kg.

#### 2.3. Case 3

A 24-month-old presents, since the age of food diversification, early persistent vomiting in spite of usual antiemetic agents. She had been hospitalized on October 30th, 2008 in pediatrics in a picture of protein-calorie malnutrition with a weight of 8.400 kg.



**Fig. 3.** End-to-end anastomosis after resection.

Download English Version:

<https://daneshyari.com/en/article/4161497>

Download Persian Version:

<https://daneshyari.com/article/4161497>

[Daneshyari.com](https://daneshyari.com)