



## Cardiac complications after laparoscopic large hiatal hernia repair. Is it related with staple fixation of the mesh? -Report of three cases



Maria del Carmen Fernandez <sup>a</sup>, María Diaz <sup>a</sup>, Fernando López <sup>a, b, \*</sup>, Roberto Martí-Obiol <sup>a</sup>, J. Ortega <sup>a, b</sup>

<sup>a</sup> Department of Surgery Clinic University Hospital, Valencia, Spain

<sup>b</sup> Department of Surgery, University of Valencia, Spain

### HIGHLIGHTS

- Three cases with different clinical expression of cardiac tamponade after laparoscopic large hiatal hernia repair.
- We review cardiac complications related to laparoscopic hiatal hernia repair.
- It is necessary to consider the risk of injury to the surrounding tissues during the anchorage of the mesh to the diaphragm.
- Cardiac complications must be considered in the postoperative period of mesh hiatoplasty.

### ARTICLE INFO

#### Article history:

Received 21 May 2015

Received in revised form

14 September 2015

Accepted 15 September 2015

#### Keywords:

Giant hiatal hernia  
Mesh hiatoplasty  
Cardiac complications  
Cardiac tamponade  
Mesh fixation

### ABSTRACT

**Introduction:** Laparoscopic Nissen operation with mesh reinforcement remains being the most popular operation for large hiatal hernia repair. Complications related to mesh placement have been widely described. Cardiac complications are rare, but have a fatal outcome if they are misdiagnosed.

**Presentation of cases:** We sought to outline our institutional experience of three patients who developed cardiac complications following a laparoscopic Nissen operation for large hiatal hernia repair.

**Discussion:** Laparoscopic hiatoplasty and Nissen fundoplication are safe and effective procedures for the hiatal hernia repair, but they are not exempt from complications. Fixation technique and material used must be taken into account. We have conducted a review of the literature on complications related to these procedures.

**Conclusion:** In the differential diagnosis of hemodynamic instability after laparoscopic hiatal hernia repair, cardiac tamponade and other cardiac complications should be considered.

© 2015 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Limited. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Laparoscopic mesh hiatoplasty for large hiatal hernias and antireflux surgery with prosthetic hiatal closure have been shown to be safe and effective procedures [1]. However, complications related to hiatal hernia repair with a mesh implant have been described [2]. Cardiac complications are rare and hardly reported in the literature.

We report our institutional experience in this surgery showing three different cases of cardiac complications after laparoscopic

hiatal hernia repair using a mesh implant and Nissen fundoplication.

## 2. Case report 1

A 46-year-old man was studied because of gastroesophageal reflux disease refractory to medical treatment. Endoscopic and contrast X-Ray studies were consistent with large hiatal hernia with most of the stomach bulging into the intrathoracic region.

He was eligible for surgery so a laparoscopic Nissen operation with mesh reinforcement was performed using Securestrap® (Ethicon, Norderstedt, Germany) for the mesh fixation. In the immediate postoperative period, the patient was doing well and remained hemodynamically stable. 48 h after surgery, patient developed hypotension, sweating and chest pain radiating to the

\* Corresponding author. Department of Surgery Clinic University Hospital, Valencia Spain.

E-mail addresses: [mcfm89@gmail.com](mailto:mcfm89@gmail.com) (M.C. Fernandez), [mariaadiaztozbarra@gmail.com](mailto:mariaadiaztozbarra@gmail.com) (M. Diaz), [ferlomo@gmail.com](mailto:ferlomo@gmail.com) (F. López), [romartob@gmail.com](mailto:romartob@gmail.com) (R. Martí-Obiol), [joaquin.ortega@uv.es](mailto:joaquin.ortega@uv.es) (J. Ortega).

left shoulder blade. EKG showed a diffuse ST-segment elevation. The echocardiogram was significant for mild pericardial effusion. The patient was then taken to Intensive Care Unit (ICU) for hemodynamic management.

Blood tests showed a troponin elevation supporting the diagnosis of myopericarditis secondary to thoraco-abdominal surgery. The patient condition improved after medical treatment with *Lysine-acetylsalicylic acid* so he could be discharged from ICU.

Despite the improvement, the patient persisted with chest pain and dyspnea. Cardiac MRI showed a large pericardial effusion (Fig. 1). Pericardiocentesis was performed with removing 650 cc of bloody fluid. The pericardial drain could be removed two days later.

The patient course was uncomplicated with resolution of the pericardial effusion on successive echocardiograms. He was discharged on the 15th postoperative day without medical treatments.

At present, patient has no recurrence of the gastric volvulus nor the pericardial effusion.

### 3. Case report 2

A 62-year-old female patient with history of arterial hypertension and osteoporosis and past surgical history of double adnexectomy and appendectomy, presented to our outpatient clinic with anemia and gastroesophageal reflux symptoms. After the diagnosis of a large hiatal hernia she was proposed for surgery. A Laparoscopic mesh hiatoplasty was performed using Securestrap® associating a Nissen fundoplication.

At the fifth postoperative day the patient developed acute respiratory distress syndrome with hypotension and oliguria that required an ICU management with intubation, mechanical ventilation and high doses of Noradrenaline. The thoracic CT-scan showed a bilateral pleural effusion and a large pericardial effusion that caused the collapse of the right atrium and ventricle (Fig. 2). She was taken back to the operating room for urgent surgical pericardial drainage. Exploration of the hiatus revealed an hematoma volume of 2.500–3.000 cc without active bleeding.

Patient condition got better improving her ventilatory function and without needs of amine support. The echocardiogram revealed a mild residual pericardial effusion and patient could be discharged from ICU.

After that, postoperative course was uneventful. She was discharged on the 23rd postoperative day with no needs of medical treatment. Nowadays, she does not present recurrence of hiatal hernia or pericardial complications.



Fig. 2. CT-scan showing a bilateral pleural effusion and a large pericardial effusion causing the collapse of the right atrium and ventricle.

### 4. Case report 3

A 70-year-old woman with medical history of hypothyroidism referred suffering from chest pain and heartburn for months. She was studied with CT-scan and upper GI X-Ray series, which demonstrated a giant hiatal hernia with an intrathoracic stomach (Fig. 3). The patient underwent to laparoscopic surgery and a Nissen fundoplication with mesh reinforcement using Securestrap® was performed. At the second postoperative day the patient presented hypotension, tachycardia and irregular pulse. EKG showed an atrial fibrillation with rapid ventricular response. Treatment with anticoagulation and amiodarone was initiated. Thoracic X-Ray showed a bilateral pleural effusion that did not require drainage. However, the patient persisted with precordial pain and dyspnea at rest, reason why an urgent thoracic CT-scan was performed. The scan revealed an extrapericardial hematoma in the hernia sac with a mild left atrium compromise (Figs. 4 and 5).

Conservative treatment was decided and the patient condition improved until she became asymptomatic. She was discharged on the 15th postoperative day with medical treatment determined by the cardiologist.




Fig. 1. Cardiac MRI showing a large pericardial effusion.

Download English Version:

<https://daneshyari.com/en/article/4195329>

Download Persian Version:

<https://daneshyari.com/article/4195329>

[Daneshyari.com](https://daneshyari.com)