

## SYSTEMATIC REVIEW

**Traditional Chinese Medicine for the treatment of influenza: a systematic review and Meta-analysis of randomized controlled trials**

Chunya Wang, Hao Wang, Xinying Liu, Dongping Xu, Yiping Tang, Ping Luo

**Chunya Wang, Yiping Tang, Ping Luo**, Emergency & Critical Care Center, Beijing Anzhen Hospital, Capital Medical University, Beijing 100029, China**Hao Wang**, Medical Information Center, Beijing 302 Hospital, Beijing 100039, China**Xinying Liu**, Traditional Chinese Medicine and Western Medicine Center, Beijing 302 Hospital, Beijing 100039, China**Dongping Xu**, Liver Failure Therapy and Research Center, Beijing 302 Hospital, Beijing 100039, China**Supported by** the Beijing 302 Hospital Inner Research Foundation (No. YNKT2012021) and the PLA 12th Five-Year Grand Project for Key Techniques and Devices in Management of Infectious Diseases (No. BWS11J048)**Correspondence to: Dr. Xinying Liu**, Traditional Chinese Medicine and Western Medicine Center, Beijing 302 Hospital, Beijing 100039, China. liuxinying302@163.com; **Prof. Dong-ping Xu**, Liver Failure Therapy and Research Center, Beijing 302 Hospital, Beijing 100039, China. xudongping@sina.com**Telephone:** +86-10-66933255; +86-10-66933020  
**Accepted:** September 29, 2013**Abstract****OBJECTIVE:** To justify the clinical use of Traditional Chinese Medicine (TCM) in the treatment of influenza.**METHODS:** MEDLINE, EMBASE, Chinese Biomedical Literature Database, China National Knowledge Infrastructure Database, China Science and Technology Journal Database, Wanfang Database and the Cochrane Database of Systematic Reviews were searched from the date of inception until January 1, 2013, for the literature on treatment of influenza with TCM.**RESULTS:** A total of 7 randomized controlled trials

were identified and reviewed. Of these trials, 2 compared a (modified) prescription of TCM with oseltamivir and 5 compared a patent traditional Chinese drug with oseltamivir. Based on the Meta-analysis, compared to oseltamivir, the (modified) prescription had similar effect in defervescence [ $WMD=5.66$ , 95%  $CI (-32.02, 43.35)$ ,  $P=0.77$ ] and viral shedding [ $WMD=-6.21$ , 95%  $CI (-84.19, 71.76)$ ,  $P=0.88$ ], and the patent traditional Chinese drug also had similar effect in viral shedding [ $WMD=-0.24$ , 95%  $CI (-4.79, 4.31)$ ,  $P=0.92$ ] but more effective in defervescence [ $WMD=-4.65$ , 95%  $CI (-8.91, -0.38)$ ,  $P=0.03$ ].

**CONCLUSION:** TCM has potential positive effects in the treatment of influenza.

© 2014 JTCM. All rights reserved.

**Key words:** Influenza, human; Medicine, Chinese Traditional; Oseltamivir; Systematic review; Randomized controlled trials**INTRODUCTION**

Up to now, the threat of pandemic influenza remains an internationally major public health concern.<sup>1</sup> As an important pharmaceutical intervention against the influenza virus, oseltamivir has been widely used against influenza virus infection in the world.<sup>2</sup> It has been demonstrated that oseltamivir can prevent the release of progeny viral particles from infected host cells and shorten the duration of influenza and decrease the development of serious complications.<sup>3,4</sup> However, a major shortcoming of oseltamivir is the emergence of viral resistance.<sup>5</sup> Therefore, it is of crucial importance to find an effective and safety alternative to treat influenza in-

fection, especially in rural areas of China where the supply of oseltamivir is often insufficient.

In China, Traditional Chinese Medicine (TCM) with a history of thousands of years is proved to be effective for the treatment of infectious diseases.<sup>6</sup> Some studies have demonstrated that TCM can be used as an alternative treatment of influenza virus infection.<sup>7-9</sup> In this study, the clinical effect of TCM in the treatment of influenza A virus infection is evaluated through comparison with oseltamivir.

## METHODS

### Search strategy

MEDLINE, EMBASE, Chinese Biomedical Literature Database (CBM), China National Knowledge Infrastructure Database (CNKI), China Science and Technology Journal Database (VIP), Wanfang Database and the Cochrane Database of Systematic Reviews were searched from the date of inception until January 1, 2013. Of these databases, CBM, CNKI, Chinese Scientific Journal Database and Wanfang Database provided literatures in Chinese. In this study, "Traditional Chinese Medicine", "oseltamivir" and "influenza" were designed for the search. Reference lists from retrieved documents were also searched.

### Criteria of inclusion

Subjects older than 3 years and hospitalized with a clinical or laboratory diagnosis of influenza infection were included in this study. Other inclusion criteria were: (a) study design: randomized controlled trial; (b) intervention: treatment group with a relatively fixed prescription of TCM or a patent traditional Chinese drug and control group with oseltamivir; (c) human studies.

### Quality control

Data were independently extracted from each study using pre-defined forms by two investigators, and disagreement was resolved by discussion among investiga-

tors and reference to the original article. When several publications pertaining to a single study were identified, the most complete publication was used. The concrete assessments were seen in the article<sup>10</sup> (Jadad score).

### Efficacy measures

Efficacy was measured with duration of fever and viral shedding. The safety of treatment was also assessed.

### Data analysis

Data analysis was carried out using Review Manager Software 4.2<sup>11</sup> (Cochrane Collaboration, Oxford, United Kingdom). The effect measures estimated were weighted mean difference (WMD), reported with 95% confidence intervals (CI). Fixed-effect or random-effect method in the analysis depended on the absence or presence of significant heterogeneity which was evaluated by the *Chi-square* and *I-square* (*I*<sup>2</sup>) tests. In the absence of statistically significant heterogeneity, the fixed-effect method was used to combine the results. When heterogeneity was confirmed, the random-effect method was used. The overall effect was tested using *Z* scores, with significance set at  $P < 0.05$ .

## RESULTS

### Selection and characteristics of study

Searches resulted in 2132 unique articles. Finally, 7 randomized controlled trials (RCTs) were left for analysis.<sup>12-18</sup> Figure 1 shows the flow chart of study selection process. 2 trials compared the efficacies between (modified) Yinqiao powder and oseltamivir and the total number of patients involved was 93.5 trials compared the efficacies between Lianhuaqingwen capsule and oseltamivir and the total number of patients involved was 607. Table 1 summarizes the main characteristics of the included studies.

### Clinical outcomes

Comparison of the effect of Yinqiao powder with oseltamivir: 2 trials in which the effect of Yinqiao powder




Figure 1 Flowchart of study selection process

Download English Version:

<https://daneshyari.com/en/article/4200996>

Download Persian Version:

<https://daneshyari.com/article/4200996>

[Daneshyari.com](https://daneshyari.com)