



Associations between parenting partners' objectively-assessed physical activity and Body Mass Index: A cross-sectional study

Jesmond Zahra ^{*}, Russell Jago, Simon J. Sebire

Centre for Exercise, Nutrition & Health Sciences, School for Policy Studies, University of Bristol, 8 Priory Road, Bristol, BS8 1TZ, UK

ARTICLE INFO

Available online 11 June 2015

Keywords:

Accelerometry
Physical activity
Obesity
Body weight
Health behaviour

ABSTRACT

Objective. Family members have the capacity to influence each other's health behaviours. This study examined whether there were associations in the objectively assessed physical activity and Body Mass Index (BMI) of mothers and fathers.

Methods. Recruitment took place in Bristol (UK) during 2012/13. Participants were 272 pairs of parents (dyads) that wore an accelerometer for at least 500 min on 3 or more days. Parents provided demographic information and self-reported height and weight. Multi-variable linear and logistic regression models examined the relationships between parents' moderate-to-vigorous physical activity (MVPA) and BMI.

Results. MVPA minutes ($r = 0.26, p < 0.001$) and Body Mass Index ($r = 0.20, p = 0.002$) of parents were correlated. Logistic regression analysis showed that mothers were almost twice (OR 1.87, $p < 0.05$) as likely to be overweight or obese when fathers were. Linear regression models showed that at the weekend every 9 min of paternal MVPA was associated with 3 min of maternal MVPA ($r = 0.34, p < 0.001$).

Conclusions. Both physical activity and BMI of parenting partners were associated. Since parents tend to share home environments and often perform activities together or as a family, then behavioural changes in one parent may have a ripple effect for other family members.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Regular physical activity (PA) has both preventive and rehabilitative effects for many chronic health conditions (Bauman, 2004; Donaldson, 2004; Kokkinos and Myers, 2010). Approximately 33% of men and 45% of women in the UK fail to meet the recommended 30 min of moderate-to-vigorous physical activity (MVPA) on 5 days per week (Health and Social Care Information Centre, 2012). Furthermore, it is estimated that 66% of men and 57% of women (aged ≥ 20) within the UK are overweight or obese (Ng et al., 2014). Low levels of PA have been associated with overweight and obesity in adults and children and recent data suggests that individuals who meet the PA guidelines are at reduced risk of being overweight or obese (Chau et al., 2012; Hills et al., 2011; Health and Social Care Information Centre, 2012). There is a need to develop strategies to increase PA and reduce obesity. Designing successful interventions requires an understanding of the determinants of PA behaviours; those interventions that target casual determinants of behaviour are likely to be more effective (Michie et al., 2008).

Families and family environments are proposed to be central components in both adult and child PA behaviours and choices. Diet and

physical activity are both influenced by core relationships within the home (Wilson, 2002) and therefore spouses are likely to be influential when it comes to modifying and maintaining healthy behaviours. Studies examining spousal correlations in PA levels have provided evidence of a relationship. A pedometer study conducted in Northern France identified an inter-spousal correlation during weekend days ($r = 0.14$); there was no evidence of an association during week days (Jacobi et al., 2011). These findings are consistent with the analysis from the Quebec Family study which highlighted an inter-spousal PA relationship (Simonen et al., 2002). This was the strongest for self-reported past year PA data ($r = 0.43$). Further analysis of previous three day PA data, recorded via diaries, demonstrated a correlation between spousal moderate-to-strenuous activity ($r = 0.22$). However, measuring PA using self-report can result in measurement error, bringing into question the validity of the findings and the accuracy of the reported correlations (Sternfeld et al., 2012; Prince et al., 2008). Previous studies have been limited by the use of pedometer or self-reported collected PA data; neither of which allows for the calculation of important output measures, such as minutes in MVPA. Further research into spousal PA association is required that utilises objective measures in order to provide a more reliable indication of spousal PA associations.

Research has demonstrated that health-related behaviours of spouses converge over time (Meyler et al., 2007). It is therefore possible

^{*} Corresponding author.

E-mail addresses: jez.zahra@bristol.ac.uk (J. Zahra), russ.jago@bristol.ac.uk (R. Jago), simon.sebire@bristol.ac.uk (S.J. Sebire).

that spousal body composition may be similar between parents. Both marriage and parenthood are associated with decreased PA and increased body weight in males and females (Hull et al., 2010; Averett et al., 2008) suggesting that individuals' behaviours may become similar once in a relationship. Previous research has demonstrated that changes from an unhealthy to a healthy behaviour in one spouse are associated with a positive change by the other partner (Falba and Sindelar, 2008). In addition, there are some evidences of associations in spousal Body Mass Index (BMI) (Jeffery and Rick, 2002; Abrevaya and Tang, 2011; Jacobi et al., 2011). Within the UK there are very few studies that have explored spousal BMI and even fewer with a specific focus on parents. The current study examined the spousal associations in parental PA and BMI among parents with children aged 5–6 years. Through utilising an objective measure of physical activity we hope to build on the findings of previous spousal PA studies. The main aim of the study is to investigate if the BMI and physical activity (MVPA) of parent dyads are associated.

Methods

Study design

Data are from a cross-sectional study (B-ProAct1v) carried out at the University of Bristol. The study aimed to examine factors that influenced young children's (5–6 years) and parents' PA, with a specific focus on the influence of parents on child PA behaviours (Jago et al., 2014b; Jago et al., 2014a). Data were collected from 57 primary schools in the greater Bristol area between January 2012 and May 2013, during this time 250 primary schools in Bristol and the surrounding areas were invited to participate. Ethical approval was granted by the School for Policy Studies research ethics committee at the University of Bristol and written informed consent was obtained from parents.

Participants

To take part in the project children and at least one parent needed to participate; second parents were encouraged to take part but it was not a requirement of the project. In total 1456 pupil and parent dyads agreed to take part. 1267 child–parent dyads wore and returned an accelerometer and were included in the final dataset. For the current study we were interested in parent dyads (e.g., mother–father) and therefore only cases where both parents participated are included in our analysis. Fig. 1 shows the study flow of participants.

Measures

Participating parents (first parents) were required to complete a questionnaire about family characteristics, personal demographics, overall child health and questions relating to screen-viewing behaviours. Second parents were asked to report demographic information. All parents that took part were asked to report their gender, height and weight to enable the calculation of BMI ($\text{BMI} = \text{kg}/\text{m}^2$). In addition, parents (first and second) and children wore an Actigraph GT3X accelerometer for five days; three week days and two weekend days. Uniaxial data were processed using Kinesoft (v3.3.75; Kinesoft, Saskatchewan, Canada) software which estimated minutes of MVPA for parents based on an age-appropriate cut point (≥ 2020 counts per minute) (Tudor-Locke et al., 2010). Sedentary behaviour and light activity were omitted from the analysis as they cannot be analysed in relation to UK PA government guidelines. Mean counts per minute (CPM) were also calculated. This measure provides an indication of the volume of physical activity in which participants engaged.

A valid day was specified as ≥ 500 min of wear time, this was to prevent further reductions in the dataset and to conform with the method used in previous studies working with the same data (Jago et al., 2014a). Any period of 60 min or more of consecutive zero counts, with an

allowance of 2 min of interruption, was classified as non-wear time and removed from subsequent analysis. Participants' home address and postcode were reported and then linked to lower layer super output data enabling the calculation of Index of Multiple Deprivation (IMD) scores based on the English Indices of Deprivation (<http://data.gov.uk/dataset/index-of-multiple-deprivation>).

Data management/reduction

In total 317 participating dyads met the inclusion criteria for at least three valid days of accelerometer data. Participants that did not provide personal demographic information were removed from the subsequent analysis. Following this criteria 272 participants were included in the final dataset, of which 269 provided at least one valid weekend day of wear time (wear time minutes, $M = 798$, $SD = 66$, Range = 579 to 949). Daily minutes of MVPA were calculated as a continuous variable. Participants were also categorised as either meeting or failing to meet the recommended 30 min/day (World Health Organization, 2010). Similarly BMI was computed as a continuous variable and also dichotomised to express weight status; normal weight or overweight/obese as defined by the UK National Health Service (<http://www.nhs.uk/Conditions/Obesity/Pages/Introduction.aspx>).

Statistics

Student t-tests were used to explore differences in demographic factors, BMI and MVPA between included and excluded participants. Descriptive statistics (i.e., means and standard deviation) were calculated and correlations were used to examine relationships between parents' MVPA and BMI. Linear regressions were conducted for each outcome variable and separately for weekday and weekend days to enable comparisons. Female data was used as the outcome and male data as the exposure. Logistic regression analysis was used to examine whether fathers meeting the recommend 30 min of daily MVPA was associated with mothers meeting the same guideline. Additional logistic regression analysis examined the odds of mothers being overweight/obese when fathers were above the BMI threshold for normal weight status. Each model was adjusted for female age, BMI, employment status, number of children, household cars and IMD scores as these have been previously associated with PA in adults (Trost et al., 2002; Cerin et al., 2009; Hull et al., 2010). Since participants were recruited via local primary schools, robust standard errors were used in all models to account for clustering of participants in schools. Analyses were conducted in Stata version 13.1 (Statacorp, College Station, TX). The R^2 value has been added to the regression model to provide an indication of the variance in outcome variables explained by exposure variables in this sample (Pearce, 2011).

Results

Descriptive statistics are shown in Table 1. On average males were approximately 1 year older than females and had slightly higher BMI. Males accumulated more MVPA minutes than females; fathers averaged 52 min of MVPA per day overall, and 56 min during weekdays. Both males and females performed approximately 10 min less MVPA on weekend days.

Parental BMI was positively correlated ($r = 0.20$ $p < 0.001$). Adjusted linear regression analysis showed that an increase of one unit in male BMI was associated with an increase of 0.22 units ($p < .001$) in mothers, which could be interpreted as an increase of one unit (kg/m^2) per five unit increase in father BMI. Adjusted logistic regression analysis showed a mother was 87% more likely to be overweight or obese if the father was overweight or obese (Table 2).

There was strong evidence of an association between parental PA ($r = 0.26$, $p < 0.001$). Table 3 displays the results from linear regressions performed on MVPA data. Mothers' weekend activity increased by 0.34 of a minute per extra minute of father MVPA, or an extra 3 min for every

Download English Version:

<https://daneshyari.com/en/article/4202456>

Download Persian Version:

<https://daneshyari.com/article/4202456>

[Daneshyari.com](https://daneshyari.com)