



Case report

Airway Obstruction Caused by Penicilliosis: A Case Report and Review of the Literature[☆]Viboon Boonsarngsuk,^{a,*} Dararat Eksombatchai,^a Wasana Kanoksil,^b Visasiri Tantrakul^a^a Division of Pulmonary and Critical Care Medicine, Department of Medicine, Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand^b Department of Pathology, Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand

ARTICLE INFO

Article history:

Received 6 March 2014

Accepted 25 April 2014

Available online 25 March 2015

Keywords:

Penicilliosis

Immunocompromised patient

Adult-onset immunodeficiency

ABSTRACT

Penicilliosis is an opportunistic infection in HIV-infected and other immunocompromised patients mostly in Southeast Asia, Southern China, Hong Kong, and Taiwan, with respiratory manifestations in about one-third of patients. We report the case of a 26-year-old non-HIV immunocompromised patient presenting with an airway obstruction caused by penicilliosis, together with a review of the literature of this rare condition.

© 2014 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

Penicilliosis endobronquial: presentación de un caso y revisión de la literatura

RESUMEN

La peniciliosis es una infección oportunista que se da en pacientes con infección por el VIH y otros pacientes inmunodeprimidos, sobre todo en el Sudeste Asiático, el sur de China, Hong Kong y Taiwán. Se producen manifestaciones respiratorias en alrededor de una tercera parte de los pacientes. Presentamos aquí el caso de un paciente de 26 años de edad inmunodeprimido, sin VIH, que comenzó con una peniciliosis endobronquial que obstruía las vías aéreas, junto con una revisión de la literatura de este trastorno muy poco frecuente.

© 2014 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Peniciliosis

Paciente inmunodeprimido

Inmunodeficiencia de inicio en la edad adulta

Introduction

Penicilliosis is an infection caused by *Penicillium marneffei*, a thermally dimorphic fungus. At room temperature (25 °C), it exhibits morphology characteristic of a mold, but grows in yeast-form when found in host tissues or in culture at 37 °C. *P. marneffei* is limited geographically to Southeast Asia, Southern China, Hong Kong and Taiwan. Although it has been suggested that bamboo rats are a reservoir for this fungus, the specific reservoir for transmission to humans remains unclear.

In man, *P. marneffei* is an opportunistic fungus that affects HIV-positive and other immunocompromised patients. Ingestion

or inhalation of fungus conidia could be the mode of transmission. Although the most common forms of presentation are non-specific and consist of low-grade fever, weight loss and anemia, the characteristic skin lesion is a central umbilicated papule. Since it is usually present, it is an important key to diagnosis.¹ Respiratory symptoms occur in around one third of patients, and diffuse reticulonodular, diffuse reticular, localized alveolar or localized reticular infiltrates, as well as cavitated lesions, can be identified on chest radiographs.² We describe the case of an immunocompromised patient who presented with obstruction of the right lower lobe bronchi caused by penicilliosis. We also present a literature review of this very rare condition.

Case Report

In April 2013, a 26-year-old man came in with fever and productive cough. He had been diagnosed with systemic lupus erythematosus (SLE) in 2002 on the basis of a malar rash, positive anti-nuclear antibody and anti-double-stranded DNA antibody

[☆] Please cite this article as: Boonsarngsuk V, Eksombatchai D, Kanoksil W, Tantrakul V. Penicilliosis endobronquial: presentación de un caso y revisión de la literatura. Arch Bronconeumol. 2015;51:e25–e28.

* Corresponding author.

E-mail address: bss-vb@hotmail.com (V. Boonsarngsuk).




Fig. 1. (A) Chest X-ray on initial presentation, in which a mottled ground glass opacity and consolidation at the level of the right lower lobe can be seen. (B) Chest computed axial tomography (CT) image in a mediastinal window setting, showing an area of consolidation at the level of the right lower lobe, with an endobronchial lesion obstructing the basal segmented bronchi (arrow). (C) Bronchoscopy revealed a whitish endobronchial mass (arrow) at the opening of the basal anterior, lateral and posterior segmented bronchi of the right lower lobe. The follow-up chest X-ray (D) and chest CT (E) showed improvement of this lesion. The basal segmented bronchi in the right lower lobe were restored and identified on the chest CT (E) and in the bronchoscopy image (F). RB7: medial basal segmented bronchus of the right lower lobe.

tests, and class IV lupic nephritis with rapidly progressive glomerulonephritis. His symptoms improved following corticosteroid and azathioprine treatment, which was gradually tapered off until it was discontinued in 2010. Two years later, the patient presented cervical lymphadenopathies. *Mycobacterium abscessus* was identified in lymph node and blood cultures. Treatment was established with clarithromycin, doxycycline and levofloxacin. Six months later, the blood cultures were negative for *Mycobacterium*, and the lymphadenopathies had regressed.

When the patient first came to the clinic, he had fever, with a body temperature of 38.2°C and normal oxygen saturation (SpO₂ 99%). Physical examination revealed fine crackles in the right lower lobe, but was otherwise unremarkable. There were no signs of skin lesions, lymphadenopathies or hepatomegaly.

Laboratory test results showed mild leukocytosis, normal routine serum biochemistry and negative HIV serology. Chest radiograph revealed a mottled ground glass opacity and consolidation at the level of the right lower lobe (Fig. 1A). Sputum Gram-staining and acid-fast bacilli testing were negative. Empirical treatment for bacterial pneumonia with imipenem–cilastatin was started, but the patient's clinical condition and chest radiograph did not improve. Chest computed tomography (CT) showed a consolidation in the right lower lobe and luminal stenosis of the basal segmented bronchi (Fig. 1B). There were also multiple small mediastinal lymph nodes.

Bronchoscopy performed one day later showed a whitish endobronchial mass at the opening of the basal anterior, lateral and posterior segmented bronchi of the right lower lobe (Fig. 1C). Histopathological examination of the endobronchial biopsy revealed acute and chronic inflammation with no granuloma formation. However, Grocott methenamine silver (GMS) staining showed a sausage-shaped yeast-type organism, long and

oval with clear central septa. Tissue and bronchial lavage cultures were positive for *P. marneffeii*.

The patient was treated with intravenous amphotericin B (1 mg/kg/day). The fever subsided after 5 days, and after 2 weeks treatment was switched from intravenous amphotericin B to oral itraconazole (400 mg/day). Two months later, the follow-up chest radiograph showed notable improvement in the lesions in the right lower lobe (Fig. 1D). The follow-up CT scan revealed only minimal residual ground glass opacity, interstitial thickening and slight peribronchial thickening in the previously affected areas (Fig. 1E). Repeat bronchoscopy showed complete resolution of the endobronchial lesion (Fig. 1F). After 6 months, oral itraconazole treatment was reduced to 200 mg/day for prophylaxis.

Given the patient's history of 2 opportunistic infections (*M. abscessus* and *P. marneffeii*) despite all immunosuppressant drugs having been discontinued for 2 years, underlying immunodeficiency was suspected. Anti-interferon-gamma (IFN- γ) antibodies were analyzed, confirming the diagnosis of adult-onset immune deficiency.³

Discussion

Endobronchial fungal infection is a very rare manifestation compared to other forms of presentation of fungal lung infections. The most common fungi causing endobronchial infection are *Aspergillus* spp., *Coccidioides immitis*, zygomycetes, *Candida* spp., *Cryptococcus neoformans* and *Histoplasma capsulatum*.⁴ Endoscopic findings differ in the various fungi, but definitive diagnosis requires microbiological identification (smears, culture or histopathology sections).

Pulmonary penicilliosis usually manifests as parenchymal or interstitial lesions. Endobronchial penicilliosis is much rarer. It can occur in immunocompromised hosts with or without HIV, and

Download English Version:

<https://daneshyari.com/en/article/4205352>

Download Persian Version:

<https://daneshyari.com/article/4205352>

[Daneshyari.com](https://daneshyari.com)