



Original Article

Evolution and Complications of Chest Trauma[☆]

Régulo José Ávila Martínez,^{a,*} Ana Hernández Voth,^b Carmen Marrón Fernández,^a
 Fátima Hermoso Alarza,^a Iván Martínez Serna,^a Andrea Mariscal de Alba,^a
 Mauricio Zuluaga Bedoya,^a María Dolores Trujillo,^a José Carlos Meneses Pardo,^a
 Vicente Díaz Hellin,^a Emilio Larru Cabrero,^a A. Pablo Gámez García^a

^a Servicio de Cirugía Torácica, Hospital Universitario 12 de Octubre, Madrid, Spain

^b Servicio de Neumología, Hospital Universitario 12 de Octubre, Madrid, Spain

ARTICLE INFO

Article history:

Received 26 September 2012

Accepted 18 December 2012

Available online 11 April 2013

Keywords:

Thoracic injuries

Comorbidities

Complications

ABSTRACT

Objective: To describe the clinical characteristics and risk factors of patients with chest trauma, and to evaluate their correlation with the development of complications.

Methods: Descriptive, prospective and analytical study of a patient cohort with chest trauma who underwent follow-up for a period of 30 days. Excluded from the study were those patients with moderate to severe traumatic brain injury, long-bone fractures, abdominal trauma and patients requiring mechanical ventilation.

Results: A total of 376 patients met the inclusion criteria, 220 of whom were males (58.5%). The most frequent causes of trauma were falls (218 cases; 57.9%) and motor vehicle accidents (57 cases; 15.1%). The most frequent type of trauma was rib contusion (248 cases; 65.9%) and rib fractures (61 cases; 16.2%). Complications were observed in 43 patients (11.4%), mainly hemothorax (13 cases), pneumothorax (9 cases), pneumonia (6 cases) and acute renal failure (4 cases). Four patients died due to pneumonia and hemothorax. Thirty-three patients were hospitalized (8.7%) and 10 (2.6%) required later re-admittance. The risk for complications increased significantly in patients with more than 2 rib fractures, in those over the age of 85 and in the presence of certain comorbidities, such as COPD and pathologies requiring anticoagulation therapy. The risk for re-admittance is higher in patients over the age of 60.

Conclusions: Patients with chest trauma who present certain comorbidities, are over the age of 85 and have more than 2 rib fractures may present more complications. These factors should be contemplated in the evaluation, management and follow-up of these subjects.

© 2012 SEPAR. Published by Elsevier España, S.L. All rights reserved.

Evolución y complicaciones del traumatismo torácico

RESUMEN

Objetivo: Describir las características clínicas y los factores de riesgo de los pacientes con traumatismo torácico, y evaluar su relación en el desarrollo de complicaciones.

Metodología: Estudio de tipo descriptivo, prospectivo y analítico de una cohorte de pacientes con traumatismo torácico a los que se les hizo seguimiento durante un periodo de 30 días. Se excluyeron pacientes con traumatismo craneoencefálico moderado a severo, fracturas de huesos largos, traumatismo abdominal, y pacientes que requirieron ventilación mecánica.

Resultados: Un total 376 pacientes cumplieron criterios de inclusión, y de ellos 220 eran varones (58,5%). Las causas más frecuentes de traumatismo fueron las caídas (218 casos; 57,9%) y los accidentes de tráfico (57 casos; 15,1%). El tipo de traumatismo más frecuente fue la contusión costal (248 casos; 65,9%) y la fractura de un arco costal (61 casos; 16,2%). Se observaron complicaciones en 43 pacientes (11,4%), principalmente por hemotórax (13 casos), neumotórax (9 casos), neumonía (6 casos) e insuficiencia renal aguda (4 casos). De estos pacientes, 4 fallecieron por neumonía y hemotórax. Treinta y tres pacientes (8,7%) fueron ingresados y 10 (2,6%) requirieron reingreso hospitalario. El riesgo de complicaciones aumenta

Palabras clave:

Traumatismo torácico

Comorbilidades

Complicaciones

[☆] Please cite this article as: Ávila Martínez RJ, et al. Evolución y complicaciones del traumatismo torácico. Arch Bronconeumol. 2013;49:177–80.

* Corresponding author.

E-mail address: reguloavila@gmail.com (R.J. Ávila Martínez).

significativamente en pacientes con más de 2 fracturas costales, en mayores de 85 años y en presencia de algunas comorbilidades como la EPOC y patologías que requieren anticoagulación. El riesgo de reingreso es mayor en pacientes con más de 60 años.

Conclusiones: Los pacientes con traumatismo torácico que presentan algunas comorbilidades, son mayores de 85 años y tienen más de 2 fracturas costales pueden presentar más complicaciones, y se deben considerar estos factores en su evaluación, manejo y seguimiento.

© 2012 SEPAR. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Introduction

Injuries are the leading public health problem in the European Union, with a mortality rate of 250 000 cases per year.¹ Chest trauma is responsible for more than 30% of these deaths, directly or indirectly. However, more than 85% of cases do not require surgery, but rather general measures available to emergency physicians.

Current demographic data show progressive aging of the European population. Spain is not immune to this problem, and in recent years has undergone an increasingly noticeable change in the population pyramid.² With this increase in the adult population, it is natural that these patients are involved in traumatic accidents.³

Chest trauma is a common problem in our setting; its main causes are motor vehicle, workplace and domestic accidents.⁴

Epidemiological data published on mild and moderate chest trauma are generally from retrospective studies.⁵ It is considered that patients with mild chest trauma, with no signs of complications, can be treated on an outpatient basis. However, it should be remembered that in elderly patients, low-energy traumas such as falls can cause serious problems.

There are many publications on moderate-severe chest trauma, or in the context of patients with multiple trauma.^{5,6}

A study was proposed to describe the clinical and demographic characteristics, comorbidities and risk factors of patients with chest trauma, and to follow them up closely to assess their evolution, since patients who initially survive may develop complications in the hours, days and weeks following the trauma, which may even lead to death.⁷

Methodology

Hospital Universitario 12 de Octubre is a tertiary referral hospital for the southern Madrid area and other districts in the region, and for other autonomous regions. It is one of the largest hospitals in Spain, with more than 1300 beds, 34 theaters for elective surgery and 4 for urgent procedures. It takes at least 44 000 hospital admissions annually from an estimated population of 412 930 persons, distributed in a population pyramid very similar to that of the rest of the country.

The study population was a total of 514 patients (of whom 138 were excluded) who consecutively attended the Emergency Department of our hospital for chest trauma between 1 February and 30 April 2011. Patients with moderate to severe traumatic brain injury (TBI), long-bone fractures, abdominal trauma and patients requiring mechanical ventilation were excluded.

This is a descriptive, prospective and analytical study of a patient cohort. All patients were evaluated in the Emergency department by a Thoracic Surgery resident within the first hour of arrival, following the Advanced Trauma Life Support protocol for the management of chest trauma.⁸

Based on the clinical and radiological evaluation, they were assigned to outpatient treatment or hospital admission. The criteria for admission were: 3 or more rib fractures and/or the presence of complications such as hemothorax and pneumothorax.

Patients were observed for 30 days following the trauma through the Madrid Community Healthcare System internal information network (HORUS), and followed up at the Thoracic Surgery

clinic between the first and second week after the trauma. The following were analyzed: age (as an independent, stratified variable), sex, cause, type (according to radiological involvement) and place where the trauma occurred, presence of comorbidities, development of complications and number of hospital admissions and readmissions, average length of stay of patients admitted and need for surgery.

The statistical program used was SPSS 17.0. Univariate analysis was performed to evaluate possible relationships between the development of complications and the other variables studied using the Chi-squared test (Table 1).

Results

Table 1 describes the variables studied and their relationship with the development of the complications observed in the patients in this series.

A total of 376 patients met the inclusion criteria, 220 males (58.2%) and 156 females (41.8%). The age group that most frequently visited the Emergency department was between 31 and 35 years (38 patients; 10.1%), followed by the group between 81 and 85 years (37 patients; 9.8%). The number of complications increased over the age of 60; their relationship was significant over the age of 85, independently of the type of trauma (Table 1).

The most common causes of chest trauma were falls (218 cases; 57.7%) mainly in patients over 71, and motor vehicle accidents (57 cases; 15.1%), predominantly in those aged between 31 and 55 years (Fig. 1).

The most common types of trauma were rib contusion (248 cases; 65.6%) and rib fractures (61 cases; 16.1%). With respect to the development of complications according to the number of rib fractures, the appearance of complications was statistically significant when there were more than 2 rib fractures (Table 1).

The most common accident site was the home (160 cases; 42.5%), road (145 cases; 38.5%) and nursing homes (68 cases; 18.1%).

The comorbidities observed in these patients were: hypertension (HT) (128 cases; 33.9%), dyslipidemia (64 cases; 16.9%), diabetes mellitus (DM) (32 cases; 8.5%), chronic obstructive pulmonary disease (COPD) (29 cases; 7.7%), pathologies associated with anticoagulation therapy (20 cases; 5.3%) and chronic renal




Fig. 1. Cause of trauma according to age group.

Download English Version:

<https://daneshyari.com/en/article/4205915>

Download Persian Version:

<https://daneshyari.com/article/4205915>

[Daneshyari.com](https://daneshyari.com)