



Case report

Bronchial leiomyoma, a case report and review of literature



José Cárdenas-García MD^{a,*}, Alfredo Lee-Chang MD^b, Virginia Chung MD^b,
Chang Shim MD^b, Stephen Factor MD^c, Amit Tibb MD^b

^a Division of Pulmonary, Critical Care and Sleep Medicine, Hofstra North-Shore LIJ School of Medicine, New Hyde Park, NY, USA

^b Pulmonary Division, Department of Medicine, Jacobi Medical Center, Albert Einstein College of Medicine, Bronx, NY, USA

^c Department of Pathology, Jacobi Medical Center, Albert Einstein College of Medicine, Bronx, NY, USA

A B S T R A C T

Keywords:
Leiomyoma
Endobronchial tumor

A 44 year old male former smoker from Ecuador presented with productive cough for 3 weeks, positive tuberculin skin test, 40 lbs weight loss and right lower lobe collapse. He denied wheezing or hemoptysis. He was treated with antibiotics and ruled out for tuberculosis with negative sputum smear. Bronchoscopy showed an endobronchial lesion at the distal end of bronchus intermedius as cause of the collapse. Endobronchial biopsy of the lesion revealed an endobronchial leiomyoma, a rare cause of endobronchial tumor. The patient underwent bilobectomy as definite therapy for the leiomyoma due to its large size and possible extra-luminal extension, which made it not amenable to bronchoscopic resection or bronchoplasty. Differential diagnoses of endobronchial lesions are discussed along with clinical, radiographic, pathologic characteristics and various treatment modalities for endobronchial leiomyomas.

© 2014 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Introduction

Leiomyomas of the respiratory tract can present anywhere along the tracheo-bronchial tree or within the lung parenchyma. If definitive diagnosis can be made with biopsy, then lung-preserving techniques such as endobronchial removal or bronchoplasty can be used.

Case report

A 44 year-old male former heavy smoker was referred to our hospital for an abnormal chest radiograph and cough. He reported productive cough, fevers and night sweats for three weeks. On review of systems he had an unintentional weight loss of 40 lbs over the previous four months. He never had a tuberculin skin test (TST) and denied tuberculosis contacts or wheezing. He was originally from Ecuador and had a history of right lower lobe pneumonia in the previous two years. A TST placed on admission was positive (20 mm). Decreased breath sounds on auscultation and dullness on percussion were appreciated at the right base. Blood tests revealed

only mild leukocytosis (13,300 cells/uL) without bandemia. His basic metabolic panel and liver function panel results were unremarkable.

The patient's posteroanterior and lateral chest radiographs are shown in Fig. 1A–B. Chest CT images (axial and coronal views) are shown in Fig. 2A–B. Pulmonary function tests were normal. Flexible bronchoscopy showed a smooth round white polypoid lesion with wide base at the distal end of bronchus intermedius almost completely occluding the lumen (Fig. 3). White thick mucoid substance could be seen exuding from the right middle lobe bronchus. Our patient underwent bilobectomy and a biopsy of the polypoid lesion is shown in Fig. 4A–B. The diagnosis of an endobronchial leiomyoma causing complete obstruction of bronchus intermedius was made.

Discussion

Clinical discussion

Endobronchial leiomyomas are extremely rare benign tumors; Forkel reported the first case in 1909 [1]. They account for 33–45% of all pulmonary leiomyomas [2,3], and are usually found in young middle age patients (39.1 years in average) without sex predilection [2,4].

Clinical presentation depends on the location of the tumor, its size and changes in the lung distal to the lesion. Bronchial lesions

* Corresponding author. Division of Pulmonary, Critical Care and Sleep Medicine, Hofstra North Shore LIJ School of Medicine, 410 Lakeville Road, New Hyde Park, NY 11042, USA.

E-mail address: jdecardenasg@gmail.com (J. Cárdenas-García).




Fig. 1. Chest radiograph shows right lower lobe (RLL) collapse, similar to that of 2 years ago A. Posteroanterior view. B. Lateral view.

produce symptoms due to partial or complete obstruction of the affected bronchus, which may include wheezing, orthopnea, hemoptysis, recurrent pneumonia and subsequent bronchiectasis [5]. Kwon described cough, dyspnea and fever as major symptoms in pulmonary leiomyomas in the case series he reported [6]. Our patient had all of those symptoms.

Endobronchial lesions comprise a heterogeneous group of pathologic entities (Table 1). Only 10% are benign tumors, papilloma being the most common form [7]. Our patient presented with recurrent pneumonia, significant weight loss and a positive TST, which lead to the presumptive diagnosis of active tuberculosis. However, sputum smear analysis was negative for acid-fast bacilli (AFB). Since endobronchial tuberculosis (EBTB) is sputum positive for AFB only in 16–53.3% [8], and our patient was originally from a high-endemic area, we decided to start empirical treatment for endobronchial tuberculosis. Interestingly, constitutional symptoms including fever, weight loss, anorexia and night sweat are not usually prominent in EBTB [8]. We discontinued antituberculosis therapy two months later; after both endobronchial biopsy and bronchoalveolar lavage cultures for mycobacteria returned negative.

The treatment for endobronchial leiomyoma depends on its location in the airway as well as the size and width of the lesion. Most endobronchial leiomyomas reported in the literature have been resected either by lobectomy or pneumonectomy [9]; however less invasive procedures have been reported as safe alternatives including simple bronchoscopic removal with or without laser [6], bronchoplasty, or bronchotomy [10] utilizing sleeve resection of the involved bronchus while sparing the distal lung

resection. Parenchymal resection is appropriate if there is a solitary parenchymal nodule or end-stage infection distal to the obstruction [11]. Bronchoscopic intervention is the modality of choice nowadays if the lesion is not wide based [6]. The prognosis of leiomyoma is excellent after complete resection with rare recurrence with only two cases reported [6,12]. Our patient underwent flexible bronchoscopy, during which limited endobronchial biopsies were obtained due to bleeding. He was referred then to thoracic surgery team for resection. A right lung bilobectomy (middle and lower lobes) was carried out via video assisted thoracoscopic surgery (VATS) due to its large size, wide base and suspected extraluminal extension, which would render it not amenable to bronchoscopic resection or bronchoplasty. The patient had an uneventful post-operative course and was followed up in our outpatient clinic. Post surgical pulmonary function tests were within normal limits.

Radiological discussion

The diagnosis of leiomyoma can not be made by imaging alone since there are no pathognomonic features. Atelectasis is the most frequent finding in chest radiographs for endobronchial leiomyomas [11], but other findings such as normal imaging, solitary round mass, pneumonic infiltration, unilateral emphysema or hyperlucency due to air trapping distal to the obstructed bronchus can be found [2]. Most benign endobronchial tumors produce non-specific masses in the wall of the airways, except for lipoma and cartilaginous tumors, which may show fat and calcium respectively. Hamartomas can also have both features [10]. Four cases of




Fig. 2. Chest CT scan reveals an endobronchial lesion (asterisk) in the bronchus intermedius with complete atelectasis of right lower lobe and a small pleural effusion. A, Axial window. B, Coronal lung window.

Download English Version:

<https://daneshyari.com/en/article/4212816>

Download Persian Version:

<https://daneshyari.com/article/4212816>

[Daneshyari.com](https://daneshyari.com)