

ORIGINAL ARTICLE

Spectrometric analysis and scanning electronic microscopy of two pleural plaques from mediaeval Portuguese period



T. Fernandes^{a,b,*}, R. Granja^a, P.L. Thillaud^c

^a Departamento de Biologia, Escola de Ciências e Tecnologia, Universidade de Évora, 7002-554, Portugal

^b Centro de Investigação em Antropologia e Saúde (CIAS), Universidade de Coimbra, Apartado 3046, 3001-401 Coimbra, Portugal

^c École des Hautes Études, Section des Sciences Historiques et Philologiques, La Sorbonne, Paris, France

Received 29 August 2013; accepted 10 January 2014

Available online 21 June 2014

KEYWORDS

Calcified soft tissue;
Infectious disease;
Mediaeval Portuguese
cemetery;
Pleural plaques

PALAVRAS-CHAVE

Tecidos moles
calcificados;
Doença infecciosa;
Cemitério mediaeval
português;
Placas pleurais

Abstract During an archaeological excavation at a mediaeval monastery (Flor da Rosa, Crato, Portugal), a skeleton of an adult woman was found with two calcifications in the thoracic cage. The location and the macroscopic analysis of the calcifications allowed them to be assigned as pleural plaques. Spectrometric analysis and scanning electronic microscopy enabled to establish that it originated with an infectious process. These results associated with the lesions found in the ribs and vertebrae strongly suggest tuberculosis as the cause of these pleural plaques.

© 2013 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. All rights reserved.

Análise espectrométrica e microscopia eletrónica de varrimento de 2 placas pleurais do período mediaeval Português

Resumo Durante uma escavação arqueológica de um mosteiro mediaeval (Flor da Rosa, Crato, Portugal) foi encontrado um esqueleto de uma mulher adulta, com 2 calcificações na caixa torácica. A localização e a análise macroscópica das calcificações permitiu que fossem consideradas como placas pleurais. A análise espectrométrica e a microscopia eletrónica de varrimento permitiram determinar que tiveram origem num processo infeccioso. Estes resultados, associados com as lesões encontradas nas costelas e vértebras, sugerem nitidamente a tuberculose como causa dessas placas pleurais.

© 2013 Sociedade Portuguesa de Pneumologia. Publicado por Elsevier España, S.L.U. Todos os direitos reservados.

* Corresponding author.

E-mail address: tmf@uevora.pt (T. Fernandes).

Introduction

Two pleural plaques (pathological calcifications of soft tissues) were found in the thoracic cage of the skeletal remains of a woman (FR 857), exhumed at the necropolis of the "Mosteiro da Flor da Rosa" (Crato, Portugal), during an archaeological excavation at this mediaeval monastery. The excavation took place between 1989 and 1992 and revealed about one hundred graves inside and around the church. The coins that accompanied some of the burials, which constituted almost the only grave goods, gave a dating for the cemetery between the XIV to the XIX centuries. The skeletons were found in a supine position, their lower limbs were extended and their upper limbs were folded according to Christian ritual practices.

Using infra-red spectroscopy and scanning electronic microscopy to identify the mineral content grades, the structure of the pleural plaque surface and the elemental composition, we suggest that the plaques had an infectious origin, as discussed below. In spite of the fact that the skeleton was in a poor state and incomplete, other interesting pathological lesions were observed: new bone formation on the pleural face of ribs and vertebral body osteolysis.

As far as we know pleural plaques are not a common finding on archaeological series: there are one from Israel,¹ two in France,² three in Switzerland,³⁻⁵ one in Slovakia,⁶ two in Hungary, one from Bácsalmás^{7,8} and the second from Pitvaros⁷. All of these pleural plaques, except the one from Bácsalmás, which is reported to date from the 17th century, were mediaeval. The oldest is dated from the 6th century (Saint-Denis, France) and the most recent from the 15th century (Bina, Slovakia). Although the exact dating of the present case is not confirmed, according to the archaeological report⁹ its stratigraphy strongly suggests the 15th century.

The aim of this study was to describe the pleural plaques and to identify the most probable aetiology.

Material and methods

Sex and age diagnoses were performed according to Buikstra & Ubelaker¹⁰ recommendations.

After studying both pleural plaques macroscopically, two samples were taken from the most irregular surface of one of them (fragment II – Fig. 1A and B): one sample of the edge and one sample of the inner side. In order to quantify mineral composition, these samples were analysed by infra-red spectroscopy (Fourier transform IR spectroscopy, according to Pavia et al.¹¹) by Dr. M. Daudon at the Biochemistry Laboratory of the "Hôpital Necker-Enfants Malades" (Paris). Each portion of the bone was ground, mixed with potassium bromide and pressed into pellets, which subsequently were analysed.

Results

Observation of this poorly preserved skeleton of a woman revealed, in addition to the pleural plaque, deposits of new bone in the pleural surfaces of the ribs, presenting a striated aspect, as well as some lytic lesions in several thoracic vertebral bodies.

Macroscopic observation of the plaques revealed the following features: solid, mineral, thin with spiculated margins, slightly bent (especially the larger one), irregular granular external convex side, smooth and striated internal concave side.

The spectroscopic analysis revealed a similar composition between the edge and the inner side of the plaque fragment (Fig. 2 A and B; Fragment II).

The following main mineral and organic components were found:

- 1 Carapatite, usually detected in the band of 1034 cm^{-1} , displayed by far the highest values in both graphs.
- 2 Crystalline apatite, commonly detected in the ranges between 564 cm^{-1} and 605 cm^{-1} , corresponded to the second highest values obtained.
- 3 The third highest values plotted in our graphs corresponded to a residual protein framework, identified by the bands of 1545 cm^{-1} (starch band 2) and 1650 cm^{-1} (starch band 1) usually connected to CoNH bounds, revealing the good conservation of the CH amino acids.




Figure 1 (A) Pleural plaques, external convex side. From left to right: Fragment I; Fragment II (true size). (B) Pleural plaques, internal concave side. From left to right: Fragment I; Fragment II (true size). Maximum length, maximum width and average thickness of fragments I and II are, respectively: 64.43/34.14/6.86 mm and 55.46/43.43/7.8 mm. Scanning electron microscopy (Hitachi TM 3000, SU-70, 30 keV) was performed to study the pleural plaques surface and its element composition. Because both fragments were found in the same skeleton and show similar macroscopic aspects, only one of them was analysed.

Download English Version:

<https://daneshyari.com/en/article/4214760>

Download Persian Version:

<https://daneshyari.com/article/4214760>

[Daneshyari.com](https://daneshyari.com)