




Egyptian Society of Radiology and Nuclear Medicine
The Egyptian Journal of Radiology and Nuclear Medicine

www.elsevier.com/locate/ejrnrm
www.sciencedirect.com



ORIGINAL ARTICLE

Prevalence of venous sinus stenosis in Pseudotumor cerebri (PTC) using digital subtraction angiography (DSA)



Mohamed Hamdy Ibrahim ^{a,*}, Sherif Hashim Abo Zeid ^b, Ahmed Abd Elbar ^b

^a Neurology Department, Faculty of Medicine, Ain Shams University, Cairo, Egypt

^b Neurosurgery Department, Ain Shams University, Cairo, Egypt

Received 4 August 2013; accepted 12 February 2014

Available online 6 March 2014

KEYWORDS

Pseudotumor cerebri;
Venous sinus stenosis;
Headache;
MRV;
Digital subtraction angiography (venous phase)

Abstract *Objectives:* To study the prevalence of intracranial venous stenosis in Pseudotumor cerebri patients.

Patients and methods: Thirty patients were diagnosed having PTC according to Dandy criteria. All underwent general and neurological assessment. Radiological assessment included CT scan brain $\pm$ MRI brain without contrast, MRV. All underwent digital subtraction angiography (DSA) (venous phase) to confirm the validity of filling gaps seen at the level of MRV.

Results: MRV brain showed that 24 patients (80%) showed filling gaps. Digital subtraction cerebral angiography (venous phase) showed 9 patients (30%) had stenosis in their dural sinuses. MRV showed to be a good screening tool since it had 100% sensitivity and negative predictive value. However, since it has a moderate specificity (62%) with a positive predictive value (PPV) of only 35%, then lesions detected should be confirmed with digital subtraction cerebral angiography (venous phase) particularly those involving the transverse and sigmoid sinus.

Conclusion: Studying the intracranial venous system in patients with PTC is an important step in understanding the pathophysiology of the disease. Detection of venous sinus stenosis opens the way to a novel therapeutic option for refractory patients like venous sinus stenting.

© 2014 Production and hosting by Elsevier B.V. on behalf of Egyptian Society of Radiology and Nuclear Medicine. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

* Corresponding author. Address: Ain shams University, Cairo, Egypt. Tel.: +20 971501903136.
E-mail address: mohamedhamdy_neuro2007@yahoo.com (M.H. Ibrahim).
Peer review under responsibility of Egyptian Society of Radiology and Nuclear Medicine.



Production and hosting by Elsevier

1. Introduction

Pseudotumor cerebri (PTC) is a neurological disorder presenting with symptoms of increased intracranial pressure (headache, visual disturbances, papilledema) without localizing neurological findings in an alert patient.

Various pathogenic mechanisms have been considered to explain the raised intracranial pressure in those patients. These included increased cerebral volume, bio-hormonal

mechanisms, disturbed normal CSF production/absorption balance, and increased cerebral venous pressure (1,2). None appears satisfactory on its own; however, it is well known that symptoms of PTC can be mimicked by cerebral venous sinus thrombosis and venous outflow obstruction (3). However, in practice not enough attention is paid to the role of venous disease in the etiology of PTC (4).

The role of such venous disease in PTC has been revisited as several groups using invasive monitoring, have documented high pressure in the venous sinuses in typical cases (5). It appears to be the result of focal stenotic lesions in the dural sinuses obstructing the venous outflow. This has led to the suggestion that undetected intracranial venous hypertension may after all be the substrate for PTC (6). King et al. (5) using venography, reported narrowing of the transverse sinuses with either smooth tapering of uncertain cause or intraluminal filling defects suggestive of a mural thrombus in some patients with PTC (7).

In our study we aimed at evaluating the prevalence of the venous sinus disease in the etiology of PTC using MRV and digital subtraction cerebral angiography (venous phase).

2. Patients and methods

2.1. Subjects

This study was conducted on 30 Egyptian patients with symptoms and signs of PTC. Patients were recruited from the neurology, neurosurgery departments and outpatient clinics of Ain-Shams University Hospitals. Ethics Committee approved the study and patients were informed about the details of the study and written consents were obtained.

2.1.1. Inclusion criteria were:

Modified Dandy criteria (7).

2.1.2. Exclusion criteria

1. Patients with true localizing findings on examination denoting focal brain dysfunction.
2. Patients with traumatic, neoplastic, infectious, structural or iatrogenic causes of intracranial hypertension.
3. Patients with clinical and neuroimaging evidence of acute primary dural sinus thrombosis or cortical vein thrombosis.

2.2. Methods

All patients included in this work were subjected to the following:

1. Complete general and neurological assessment.
2. Lumbar puncture (LP).
3. Full ophthalmologic assessment included:
 - A. Visual acuity measurement: using Snellen chart.
 - B. Direct and indirect ophthalmoscopic fundus examination: To assess and grade papilledema.
 - C. Automated perimetry.
4. Radiological investigations:

- a. CT scan brain $\pm$ MRI brain without contrast.
- b. Magnetic resonance venography (MRV) of the intracranial venous system by time of flight (TOF) or phase contrast techniques.
- c. Digital subtraction angiography (DSA) (venous phase).

5. Statistical methodology: Analysis of data was done by IBM computer using SPSS (statistical program for social science) (version 10).

2.3. Results

MRV brain showed that 24 patients (80%) showed filling gaps. Digital subtraction cerebral angiography (venous phase) showed 9 patients (30%) had stenosis in their dural sinuses. MRV showed to be a good screening tool since it had 100% sensitivity and negative predictive value. However, since it has a moderate specificity (62%) with a positive predictive value (PPV) of only 35%, then lesions detected should be confirmed with digital subtraction cerebral angiography (venous phase) particularly those involving the transverse and sigmoid sinus (see Fig. 1A).

3. Discussion

Pseudotumor cerebri (PTC), also known as benign intracranial hypertension (BIH), is a syndrome seen more frequently in young obese women characterized by symptoms and signs of increased intracranial pressure in the absence of an intracranial mass lesion, infection, or hydrocephalus in otherwise healthy and alert patient. The exact etiology of benign intracranial hypertension is still unknown. However, various pathological conditions may be associated with this syndrome. These include endocrine and metabolic disorders, intracranial venous sinus thrombosis, drugs and toxins, hematological and connective tissue disorders, high cerebrospinal fluid (CSF) protein content, "meningism" with systemic bacterial or viral infections, and empty sella syndrome (11).

The mechanism of increased intracranial pressure in these disorders is still unclear. Involvement of the venous flow was




Fig. 1 (A) MRV with a filling defect in the Rt transverse sinus suggestive of stenosis with aplasia of the left transverse and sigmoid sinus.

Download English Version:

<https://daneshyari.com/en/article/4224281>

Download Persian Version:

<https://daneshyari.com/article/4224281>

[Daneshyari.com](https://daneshyari.com)