



CASE REPORT

Subgaleal dermoid cyst of the anterior fontanelle in a child



Amit Agrawal^{a,*}, Vissa Santhi^b, V. Umamaheswara Reddy^c

^a Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh, India

^b Department of Pathology, Narayana Medical College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh, India

^c Department of Radiology, Narayana Medical College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh, India

Received 13 May 2015; accepted 25 August 2015

Available online 10 September 2015

KEYWORDS

Anterior fontanel;
Congenital inclusion cyst;
Dermoid cyst;
Subgaleal

Abstract The congenital inclusion dermoid cyst (CDIC) of the anterior fontanel is a benign, slow-growing lesion. Subgaleal location of dermoid cyst is extremely uncommon. We report a case of 14 year old female who had subgaleal dermoid cyst of the anterior fontanel where the cyst could be excised completely with good outcome. Computed tomography scan is the investigations of choice as it will show the greater details of the lesion and its relation to the underlying structures. Characteristic scalloping of the outer table of cranium on radiographs can help to differentiate intracranial extension from extracranial location of the lesion.

© 2015 The Authors. The Egyptian Society of Radiology and Nuclear Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The congenital inclusion dermoid cyst (CDIC) of the anterior fontanel is a benign, slow-growing, non-tender, soft swelling which is covered with intact skin and accounts for 0.1–0.5% of all cranial tumors (1–6). Total subgaleal location of dermoid cyst of the anterior fontanel is extremely uncommon (7,8). We report a case of subgaleal dermoid cyst of the anterior fontanel where the cyst could be excised completely with good outcome (9).

* Corresponding author at: Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Nellore 524003, Andhra Pradesh, India. Mobile: +91 8096410032.
E-mail addresses: dramitagrawal@gmail.com, dramit_in@yahoo.com (A. Agrawal).

Peer review under responsibility of Egyptian Society of Radiology and Nuclear Medicine.

<http://dx.doi.org/10.1016/j.ejrnm.2015.08.016>

0378-603X © 2015 The Authors. The Egyptian Society of Radiology and Nuclear Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2. Case report

A 14 year old female patient presented with the history of scalp swelling over anterior fontanel since birth. The swelling was gradually increasing in size. Now for last one month it was associated with dull aching type headache. There was no history of fever, vomiting, seizures or altered sensorium. Her general and systemic examination was unremarkable. Neurological examination including higher mental functions was normal. In cranial nerves, motor and sensory examination was normal. Local examination revealed a firm, non-tender, non-pulsatile, non-compressible swelling over the anterior fontanel. There was no bruit over the swelling, transillumination test was negative and the skin over the swelling was healthy and covered normal hair. Computed tomography (CT) demonstrated an extracranial homogenous hypodense midline swelling, overlying and covering the anterior fontanel (Figs. 1 and 2). The patient underwent total excision of the swelling.




Fig. 1 Axial plain CT images brain window (A) and bone window (B) showing soft tissue swelling at the confluence of the sutures; there was smooth scalloping of adjacent bone with just a focal breach of the inner table. Sagittal and Coronal bone window images (C, D) showing the relations of the lesion with underlying sinus.



Fig. 2 3D Volume rendered image (A) Soft tissue lesion of the vertex in better detail, (B) and (C) showing the remodeling and depression of the skull bones at the bregma caused by the lesion.

The swelling was found between the galea aponeurotica and pericranium (Fig. 3). There was no tissue connection either to the overlying skin or to the intracranial extension. Cyst wall lined by stratified squamous epithelium and lumen showed keratinous material and subepithelial fibrocollagenous tissue showed sebaceous glands, ducts and congested blood vessels (Fig. 4). The postoperative course was uneventful and the later outcome was good without recurrence.

3. Discussion

Dermoid cysts are developmental tumors that develop from germ cells displaced between the 3rd and 5th week of embryogenesis, when the ectoderm folds into the neural tube and lies along the midline at the neural groove closure (5,10–12). Histologically, these cysts have a fibrous capsule, lined by squamous epithelium and contain clear fluid with some adnexal appen-

Download English Version:

<https://daneshyari.com/en/article/4224491>

Download Persian Version:

<https://daneshyari.com/article/4224491>

[Daneshyari.com](https://daneshyari.com)