

## Key imaging characteristics for preoperative identification of cavernous sinus hemangioma

Roban Shah, MD, and Surya Nadimpalli, MD

Cavernous sinus hemangiomas are relatively rare but have characteristic imaging features. We present a case of a large extra-axial middle cranial fossa mass arising from the cavernous sinus in a 51-year-old female, and we describe the unique features and complications of this intracranial tumor. With the proper pre-operative diagnosis, an appropriate plan can be created for these highly vascular lesions, minimizing the risks involved with surgery.

### Introduction

Cavernous sinus hemangiomas (CSHs) are vascular malformations consisting of abnormal, dilated vessels without intervening neural tissue (1). They constitute 2-3% of all lesions in the cavernous sinus (2).

Pathologically, these tumors are highly vascular and contain “honeycombed vascular spaces” with stagnant blood flow (3). This vascularity and proximity to neurovascular structures complicates surgical procedures, with resulting morbidity and mortality of 36 to 38% (3). Extra-axial CH tumors rarely bleed spontaneously, in contrast to parenchymal cavernous angiomas, which have a spontaneous hemorrhage rate of 25% (3).

We present a case of a large, extra-axial, middle cranial fossa mass arising from the cavernous sinus in a 51-year-old female, and we describe the unique features and complications of this relatively rare intracranial tumor. Preoperative diagnosis is crucial to avoid massive bleeding during resection.

### Case report

#### *History*

A 51-year-old woman with a medical history of controlled hypertension became acutely nauseous and had an episode of emesis. A few hours later, she developed a headache and continued having emesis that persisted throughout the night. The following morning, the patient was found unresponsive and was taken to the emergency department.

#### *Examination*

The initial noncontrast head CT demonstrated a large, 7.5-cm right extra-axial mass in the right middle cranial fossa with right-to-left shift. There was expansion of the cavernous sinus with encasement of the right cavernous carotid, and scalloping of the bony margins of the middle cranial fossa with enlargement of the right foramen ovale (Fig. 1). Postcontrast head CT demonstrated homogeneous enhancement (Fig. 2).

Contrast-enhanced MRI demonstrated a homogeneously enhancing, lobulated, well-circumscribed mass that had homogeneously low signal intensity on T1 and high signal intensity on T2 and FLAIR (Figs. 3 through 6). The mass resulted in hydrocephalus due to obstruction of the left lateral ventricle, likely at the level of the foramen of monro. Due to avid enhancement, pre-operative embolization was performed.

#### *Angiography*

A cerebral angiogram demonstrated three hypertrophied branches from the petrous portion of the internal carotid artery, likely representing the caroticotympanic, cavernous,

---

**Citation:** Shah R, Nadimpalli S. Key imaging characteristics for preoperative identification of cavernous sinus hemangioma. *Radiology Case Reports*. (Online) 2015;10(1):1013.

**Copyright:** © 2015 The Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 2.5 License, which permits reproduction and distribution, provided the original work is properly cited. Commercial use and derivative works are not permitted.

Drs. Shah and Nadimpalli are in the Department of Radiology at the Advocate Illinois Masonic Medical Center, Chicago, IL. Contact Dr. Shah at [rshah31@gmail.com](mailto:rshah31@gmail.com).

**Competing Interests:** The authors have declared that no competing interests exist.

**DOI:** 10.2484/rcr.v10i1.1013

## Key imaging characteristics for preoperative identification of cavernous sinus hemangioma




Fig. 1. Noncontrast axial CT image demonstrates a large extra-axial mass in the right middle cranial fossa, expanding into the right cavernous sinus.

and hypophyseal arteries. Subselective catheterization of an internal maxillary branch of a right external carotid artery demonstrated an irregular blushing pattern throughout the intracranial mass (Fig. 7). Embolization of this branch was performed (Fig. 8).

### *Surgery findings*

Surgery demonstrated a vascular tumor, the majority of which was between the sphenoid wing and petrous bone. Excavation of the tumor was challenging due to its exten-




Fig. 2. Post-contrast axial CT image demonstrates homogeneous enhancement.

sive vascularity. To limit amount of blood loss, a second surgery was performed at a later date to remove the remainder of the tumor.

### *Pathological findings*

Pathology demonstrated a proliferation of blood vessels, which appeared venous in nature. Organizing hemorrhage was seen. Adjacent tissue comprised dense fibrous tissue and large nerve trunks with frequent ganglion cells, without evidence of cerebral parenchyma (Figs. 9 and 10).




Fig. 3. Sagittal T1 noncontrast image shows buckling of the cortex, supporting extra-axial location.




Fig. 4. Axial T2 MRI.

## Discussion

The etiology of CSHs is unknown. One hypothesis proposed that the tumor arises within the cavernous sinus, initially deriving its blood supply from the intracavernous ICA and recruiting additional supply from the middle meningeal and accessory middle meningeal arteries (4).

There is a 7:1 predominance in females (5), and it has been hypothesized that estrogen may play a role (2). This is

Download English Version:

<https://daneshyari.com/en/article/4248001>

Download Persian Version:

<https://daneshyari.com/article/4248001>

[Daneshyari.com](https://daneshyari.com)