

Original

Positive sentinel node risk in relation to oestrogen receptors in breast cancer in premenopausal and postmenopausal women

M.T. Bajén^{a,*}, A.M. Benítez^a, A. Domenech^a, Y. Ricart, J. Mora^a, P. Notta^a, D. Ramal^a, A. Urruticoechea^b, T. Soler^c and J. Martín-Comin^a

^a Nuclear Medicine Department, Hospital Universitario de Bellvitge -IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain

^b Medical Oncology Department, Instituto Catalán de Oncology (ICO)-IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain

^c Pathology Department, Hospital Universitario de Bellvitge -IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain

INFORMATION ON THE ARTICLE

History of the article:

Received: June 4, 2010

Accepted: July 20, 2010

Keywords:

Sentinel node

Breast cancer

Tumour oestrogen receptor status

Premenopausal and postmenopausal

women

Palabras clave:

Ganglio centinela

Cáncer de mama

Receptores estrogénicos tumorales

Mujeres premenopáusicas y

posmenopáusicas

ABSTRACT

Objective: The influence of the relationship between pre- and post-menopausal stage of patients with breast cancer (BC) and the risk of sentinel lymph node (SLN) metastasis as well as the influence of oestrogen receptor (ER) status within each one of these groups were analyzed.

Method: A BC database with 1,388 patients was analysed. Three age groups were studied: A, elderly postmenopausal, 200 patients, ≥ 70 years old; B, younger postmenopausal, 89 patients, 55-69 years old; C, premenopausal, 85 patients, < 55 years old. In each group 2 subgroups were analyzed: positive ER and negative ER tumours. Data analysed: SLN-positive patients, non-sentinel node (NSN)-positive patients, non-surgical detections (NSD) and non disease-free (NDF) patients after a follow-up of 52 months. Statistical analysis: chi-squared test, significance: $P \leq 0.05$.

Results: SLN metastasis was significantly ($P < 0.025$) more common in premenopausal than in postmenopausal patients, and within those, mainly in negative ER tumours. Positive-NSN increases slightly in premenopausal patients (exclusively in negative ER tumours). NDF patients are similar in the 3 groups and in all of them it is much more frequent in negative ER tumours.

Conclusions: SLN metastasis varies significantly according to hormonal state and not according to age, being more frequent in premenopausal patients and mainly in ER negative tumours.

© 2010 Elsevier España, S.L. and SEMNIM. All rights reserved.

Relación entre el riesgo de metástasis en el ganglio centinela y los receptores estrogénicos en pacientes pre y posmenopáusicas con cáncer de mama

RESUMEN

Objetivo: Investigar la relación entre el estado pre o posmenopáusicas de la paciente con carcinoma de mama (CM) y el riesgo de metástasis en el ganglio centinela (GC) y dentro de cada uno de esos grupos el riesgo según el estado de los receptores de estrógeno (RE).

Método: Análisis de la base de datos de GC con 1.388 pacientes. Se estudiaron tres grupos de edad: A) posmenopáusicas ancianas, 200 pacientes, ≥ 70 años; B) posmenopáusicas más jóvenes, 89 pacientes, 55-69 años, y C) premenopáusicas, 85 pacientes, < 55 años. En cada grupo se analizaron 2 subgrupos: tumores con RE positivos o negativos. Factores estudiados en cada grupo y subgrupo: pacientes con GC positivo, pacientes con ganglios no centinela (GNC) positivos, número de no detecciones quirúrgicas (NDQ) y pacientes no libres de enfermedad (NLE) tras 52 meses de seguimiento. Análisis estadístico: test de chi-cuadrado; significancia $p \leq 0,05$.

Resultados: En las premenopáusicas el número de GC positivos es significativamente ($p < 0,025$) mayor que en las posmenopáusicas y dentro de las premenopáusicas fundamentalmente en los tumores con RE negativos. El número de GNC positivos aumenta solo discretamente en las premenopáusicas y ocurre exclusivamente en tumores con RE negativos. El número de pacientes NLE es similar en los 3 grupos y en todos ellos es mucho más frecuente en pacientes con tumores con RE negativos.

Conclusiones: El número de pacientes con GC positivo varía significativamente con el estado hormonal y no con la edad de la paciente, siendo más frecuentes en las premenopáusicas y fundamentalmente en tumores con RE negativos.

© 2010 Elsevier España, S.L. y SEMNIM. Todos los derechos reservados.

* Corresponding author.

E-mail address: mtbajen@bellvitgehospital.cat (M.T. Bajén).

Introduction

The incidence of breast cancer increases with age, this being the most frequent cancer in women over 70 years of age.¹ Comorbidity also increases with age and may condition the duration and extension of the surgery. Therefore, if axillary surgery can be avoided without jeopardizing the control of the tumor, this would be a very important advantage in this group of patients. The need for lymph node staging in these patients is a controversial subject. The perception exists that regional staging in elderly patients has less influence in the subsequent treatment while there is a very extended belief that there is a lower risk of metastasis in regional lymph nodes in the older patients.²

There are various studies that confirm the clinical impression that the disease in elderly patients is more indolent and that they have a more favorable biological phenotype, as shown by the fact that there are lower rates of tumor cell proliferation, lower expression of human epidermal growth factor receptor 2 (HER 2), higher content of estrogen and/or progesterone receptors, lower rate of p53 accumulation and, what is more important, better evolution.³⁻⁵

It is well known that elderly patients have a greater likelihood of positive estrogen receptor tumors. Thus, hormone therapy in them plays a very important role, chemotherapy being relegated to a second place.⁶

It has already been demonstrated that the risk of metastasis in the sentinel node (SN) significantly decreases as age increases.² However, this risk has not been studied in relationship to the hormonal status of the patient (pre- and post-menopausal).

Since the utility of selective sentinel node biopsy in any population group depends on the pretest likelihood of detecting metastasis in the sentinel node, our study was conceived to investigate if the risk of metastasis in the sentinel node was related to, more than age, the pre- or post-menopausal status of the patient and to determine, within each one of these groups, what the risk would be according to whether the tumor has positive or negative estrogen receptors (ER).

Material and methods

Study design

This is a retrospective study of the prospective data base of sentinel nodes in breast cancer patients in the nuclear medicine department of the Hospital Universitario of Bellvitge. From June 2000 to December 2007, a total of 1388 patients suffering from breast cancer who had undergone a selective sentinel lymph node biopsy were included in the data base. Over the years, the cutoff age to consider a patient as elderly in the literature had been increasing, reaching up to 70 years, based on the prolongation that is being experienced in life expectancy.⁷ For this reason, our study has considered those women 70 years or older as an elderly patient. We have considered 55 years of age as the age threshold that separates the premenopausal patient from the post-menopausal.

The number of positive lymphadenectomies, that is, the existence of other metastatic lymph nodes in the lymphadenectomy specimen, has also been studied, taking into account that lymphadenectomy had been done because metastasis in the SN had been detected.

Furthermore, the follow-up of the patents was recorded in order to calculate how many were disease-free at the time of the study in each group.

Each one of these factors studied was also analyzed within each age group according to whether the patients had positive or negative estrogen receptor (ER) tumors.

Study population

Of the 1388 patients, 374 patients with invasive breast cancer whose tumor size was ≤ 3 cm were included in the present study.

These patients were followed-up at a mean of 52 months. Three groups were differentiated: group A, elderly post-menopausal, was made up of 200 patients, who were 70 years or older. They underwent a selective SN biopsy between June 2000 and December 2007 and were followed-up for a mean of 35 months. Group B is the group of the youngest post-menopausal subjects, made up of 89 patients, between 55 and 69 years, who underwent the SN technique between January 2001 and December 2004 and who were clinically followed-up for a mean of 66 months, and finally, group C, pre-menopausal, with 85 patients under 55 years of age, who were also operated on between January 2001 and December 2004 and had a mean follow-up of 54.5 months.

Pathological study

A SN was considered positive when metastasis was detected by hematoxylin and eosin stain or by immunohistochemistry. According to the TNM staging, if the size of the metastasis detected was >2 mm, it was considered a macrometastasis, if the size was between 0.2 and 2 mm, micrometastasis and if it was <0.2 mm, isolated tumor cells (ITC).

The lymph nodes of the lymphadenectomy specimen were only analyzed with hematoxylin and eosin stain.

Statistical analysis

Comparison between samples was made using the chi square test and the Yates correction was applied when the items calculated were between 3 and 5. Significance was determined with $p \leq 0.05$.

Results

The number of SN excised per patient was 1.46 in group A, 1.6 in group B and 1.7 in group C.

The number of patients in whom the SN was not detected surgically increased with age. In group A, elderly post-menopausal subjects, this occurred in 17 patients (17/200). In group B, the youngest post-menopausal, in 4 patients (4/89) and in group C, premenopausal, in one patient (1/85).

The number of patients with negative estrogen receptor tumors in group A was 18 (18/200), in group B, 14 patients (14/89) and in group C, 11 patients (11/85). No statistically significant differences were found between the different age groups.

Regarding patients with positive SN, this significantly increased ($p < 0.025$) in the premenopausal patient group compared to the postmenopausal patients. In Group A, we found 37 patients (37/200), in B, 16 patients (16/89) and in C, 27 patients (27/85). However, the number of these patients in whom there were other metastatic lymph nodes in the lymphadenectomy specimen only slightly increased with age: 6 patients in Group A (6/200), 5 in group B (5/89) and 6 in Group C (6/85), no significant differences ($p > 0.1$) being found between the different age groups.

The percentage of patients who were not disease-free at the time of the study was similar in the three age groups: 8 patients in the elderly patient group (8/200), 4 in the youngest postmenopausal group (4/89) and 5 in the premenopausal one (5/85) and no statistically significant differences ($p > 0.05$) were found.

All of these results are shown as percentages in table 1.

Table 2 shows the number of patients with positive SN and with other positive lymph nodes of the lymphadenectomy and the number of patients in which the SN was not detected surgically and the patients who were not disease-free at the time of the study in accordance with positive or negative ER of the tumors in each age group.

In the premenopausal group, 21 out of the 74 patients with positive ER had positive SN (28.4%) and 3 patients also had other positive lymph nodes of the lymphadenectomy (4%). Six of the 11 patients

Download English Version:

<https://daneshyari.com/en/article/4249481>

Download Persian Version:

<https://daneshyari.com/article/4249481>

[Daneshyari.com](https://daneshyari.com)