




Arab Journal of Urology (Official Journal of the Arab Association of Urology)

www.sciencedirect.com



PFUI-RELATED COMPLICATIONS REVIEW

The mechanism of continence after posterior urethroplasty



Herman S. Bagga^{*}, Kenneth W. Angermeier

Center for Genitourinary Reconstruction, Glickman Urological and Kidney Institute, Cleveland Clinic, Cleveland, OH, USA

Received 2 November 2014, Accepted 26 November 2014

Available online 7 February 2015

KEYWORDS

Posterior urethroplasty;
Urinary continence;
Genitourinary trauma;
Urethral trauma;
Pelvic fracture

ABBREVIATIONS

PFUI, pelvic fracture urethral injury;
PU, posterior urethroplasty

Abstract The standard of care after a pelvic fracture urethral injury is a repair via a one-stage anastomotic posterior urethroplasty using a step-wise perineal approach. The initial injury, immediate postoperative management, and surgical repair can all affect urinary continence in these patients. Proximal continence mechanisms, particularly the bladder neck, are particularly important in maintaining urinary continence in these patients. Patients with bladder neck dysfunction should be counselled about the greater risk of urinary incontinence.

© 2015 Production and hosting by Elsevier B.V. on behalf of Arab Association of Urology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

The anatomy of pelvic fracture urethral injury (PFUI)

In a typical man urinary continence depends on two urethral segments, conceptually separated at the level of the prostatic verumontanum. Proximally is the internal

sphincter, comprising the bladder neck and proximal prostatic urethra. Distally there are several features, including a small segment of remaining prostatic urethra, the membranous urethra, and the external sphincter, which includes the urethral sphincter mechanism and the extrinsic pelvic floor musculature [1–3].

The urethral pressure responsible for continence is normally greatest at the membranous urethra, where the skeletal urethral sphincter applies its maximum pressure [4]. During trauma associated with PFUI in men, this region of the urethra is at particular risk, putting the continence mechanisms in jeopardy. This location of injury is due to tethering of the posterior urethra with

^{*} Corresponding author at: Cleveland Clinic, 9500 Euclid Avenue, Q10, Cleveland, OH 44195, USA.

E-mail address: hermanbagga@gmail.com (H.S. Bagga).

Peer review under responsibility of Arab Association of Urology.



Production and hosting by Elsevier

the pubic arch by the puboprostatic ligaments and the perineal membrane. It is between these two relatively fixed points that the membranous urethra and the urethral sphincter mechanism lie. The region is therefore vulnerable to injury during pelvic fracture. Such trauma begins with a stretch injury and can progress to partial or complete urethral rupture, particularly when a ligamentous attachment is torn from the urethra. Adherence is thought to be tighter at the puboprostatic ligament, leaving the distal site of attachment more vulnerable to injury. Therefore, although the effects of the injury can be wider, the disruption itself is most commonly noted at the level of the bulbomembranous junction, just distal to the urethral sphincter mechanism [5–9].

This pattern of injury is in contrast to that occurring in females, where it is more common to see a partial anterior urethral tear rather than a complete urethral disruption. Injury to children can also vary, as they have less developed prostate glands which have decreased support, and thus tend more commonly to have proximal injuries through the bladder neck or prostatic urethra [7,8,10,11].

Effects of the initial management

In addition to the direct urethral damage caused by PFUI, subsequent management can result in additional urethral compromise which can further affect urinary continence. The standard of care for management begins with the immediate placement of a suprapubic catheter for urinary drainage. This is sometimes accompanied by immediate primary realignment over a catheter. If early realignment is used the patient is then managed expectantly. In cases of failure future interventions might include anastomotic urethroplasty. Historically, realignment was done as an open procedure. The invasiveness of this approach was found to cause a greater risk of urinary incontinence for the patient [12,13]. In the modern era, open realignment is rarely indicated, except if there are concomitant bladder, bladder neck, or rectal injuries. In these cases immediate open repair and realignment is recommended to prevent urinary incontinence or pelvic sepsis. Currently, immediate primary realignment is generally achieved by endoscopic means through a cystotomy, with no retropubic or perineal exploration. No additional risk to urinary incontinence has been noted for those who undergo this mode of immediate management [7].

The effects of posterior urethroplasty (PU)

Ultimately, the standard of care for managing urethral obliteration as a result of PFUI is a single-stage anastomotic urethroplasty. Generally, a perineal approach is adequate (and preferred), but a combined abdominoperineal procedure is sometimes required when a tension-free anastomosis cannot be made from below.

Before anastomosis, the scarred urethral and periurethral tissue must be resected from the obliterated portions of the bulbar and membranous urethra, with additional excision of adjacent injured tissue. Given that the dismembered region of the urethra is most often near the bulbomembranous junction, the repair is at times a bulbomembranous anastomosis. However, if the injured and scarred urethra continues proximally, it must also be excised before the anastomosis. Therefore, although there is the potential for preserving at least part of the urethral sphincter mechanism, the entire distal continence component can be put at risk during repair [1,5,7,8,14].

The mechanism of continence after surgery

After PU the proximal urethral component (comprising the bladder neck and supramontanal urethra), might serve as the primary urinary continence mechanism. For some patients, the smooth muscle of this region might represent the only remaining active part of the sphincter. However, this area is known to contribute a lower pressure to the urethral continence mechanism than its distal counterpart [1,8,14]. Koraitim et al. [14] investigated the urethral pressures after anastomotic PU in these patients, and confirmed this fact. They found a lower maximum urethral pressure and urethral closing pressure in repaired patients than in healthy controls, with decreased pressures of 39 and 48 cm H₂O vs. 65 and 75 cm H₂O, respectively.

In the same study the functional urethral length of repaired patients was found to be half of that in normal patients, at 2.4 vs. 4.8 cm. Under stress conditions intra-abdominal pressure changes were transmitted throughout this entire shortened functional urethra in repaired patients, compared to just the proximal urethral portion of healthy controls [14]. Anatomically this is not surprising, given that the proximal urethra is supported in a retropubic, intra-abdominal position. However, this finding gives more support to the contention that the proximal urethra might serve as the primary continence mechanism in patients after anastomotic urethroplasty for PFUI. In their study, Koraitim et al. [14] also investigated the contribution of periurethral skeletal muscles to continence after PU. From pressure studies while patients were asked to use a 'hold manoeuvre', they found an increased urethral pressure in all normal patients, but only in 65% of patients who had PU. This further reinforces the importance of the proximal urethral component for continence in these patients.

Bladder neck dysfunction might be secondary to sacral nerve or pelvic plexus injury, but a more common cause would be direct trauma to the bladder neck itself [1,7,14,15]. Such bladder neck injury in men is generally a longitudinal anterior rupture that extends from the prostatic urethra to the bladder neck. This injury has been described as secondary to a tearing force from the puboprostatic ligaments being pulled apart, most

Download English Version:

<https://daneshyari.com/en/article/4267966>

Download Persian Version:

<https://daneshyari.com/article/4267966>

[Daneshyari.com](https://daneshyari.com)