



Original Article

Prostate-specific antigen kinetics after stereotactic body radiotherapy as monotherapy or boost after whole pelvic radiotherapy for localized prostate cancer



Hun Jung Kim*, Jung Hoon Phak, Woo Chul Kim

Department of Radiation Oncology, Inha University Hospital, Inha University School of Medicine, Incheon, South Korea

ARTICLE INFO

Article history:

Received 10 September 2015

Received in revised form

18 September 2015

Accepted 22 September 2015

Available online 2 November 2015

Keywords:

Cyberknife

Prostate cancer

Prostate-specific antigen kinetics

Prostate-specific antigen nadir

Stereotactic body radiotherapy

ABSTRACT

Purpose: Stereotactic body radiotherapy (SBRT) has emerged as an effective treatment for localized prostate cancer. However, prostate-specific antigen (PSA) kinetics after SBRT has not been well characterized. The purpose of the current study is to assess the kinetics of PSA for low- and intermediate-risk prostate cancer patients treated with SBRT using Cyberknife as both monotherapy and boost after whole pelvic radiotherapy (WPRT) in the absence of androgen deprivation therapy.

Methods: A total of 61 patients with low- and intermediate-risk prostate cancer treated with SBRT as monotherapy (36.25 Gy in 5 fractions in 32 patients) and SBRT (21 Gy in 3 fractions in 29 patients) boost combined with WPRT (45 Gy in 25 fractions). Patients were excluded if they failed therapy by the Phoenix definition or had androgen deprivation therapy. PSA nadir and rate of change in PSA over time (slope) were calculated and compared.

Results: With a median follow-up of 52.4 months (range, 14–74 months), for SBRT monotherapy, the median PSA nadir was 0.31 ng/mL (range, 0.04–1.15 ng/mL) and slopes were -0.41 ng/mL/mo, -0.17 ng/mL/mo, -0.12 ng/mL/mo, and -0.09 ng/mL/mo, respectively, for durations of 1 year, 2 years, 3 years, and 4 years postradiotherapy. Similarly, for SBRT boost after WPRT, the median PSA nadir was 0.34 ng/mL (range, 0.04–1.44 ng/mL) and slopes were -0.53 ng/mL/mo, -0.25 ng/mL/mo, -0.14 ng/mL/mo, and -0.09 ng/mL/mo, respectively. The median nadir and slopes of SBRT monotherapy did not differ significantly from those of SBRT boost after WPRT. Benign PSA bounces were common in 30.4% of all cohorts, and the median time to PSA bounce was 12 months (range, 6–25 months).

Conclusions: In this report of low- and intermediate-risk prostate cancer patients, an initial period of rapid PSA decline was followed by a slow decline, which resulted in a lower PSA nadir. The PSA kinetics of SBRT monotherapy appears to be comparable to those achieved with SBRT boost with WPRT.

Copyright © 2015 Asian Pacific Prostate Society, Published by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Prostate cancer is the most common cancer and the second leading cause of death among men in the United States,¹ and its incidence rates in Korea, although relatively lower than those in western nations, continue to increase.² At present, intensity-modulated radiotherapy (IMRT) is the standard of care in radical radiotherapy for prostate cancer.³ In IMRT delivery, accuracy is critically important as steep dose gradients may increase the risk of

geographical miss.⁴ Cyberknife (Accuray Inc., Sunnyvale, CA, USA) is one of the tools used for hypofractionated stereotactic body radiotherapy (SBRT) and real-time image guidance to account for intrafraction prostatic motion. Advanced techniques of Cyberknife allow high doses of radiation to be delivered precisely to the target while sparing the surrounding healthy tissue.⁵

Accumulating recent clinical evidence has demonstrated that the α/β ratios of prostate cancer may be around 2 Gy, which is lower than that of the surrounding normal tissue.^{6,7} The hypofractionated radiotherapy schema may improve the biochemical control of prostate cancer without increasing toxicities associated with late-responding tissue.⁶ Analogously, hypofractionation with Cyberknife has demonstrated excellent efficacy and toxicity profiles as both monotherapy and boost after whole pelvic radiotherapy (WPRT) for localized prostate cancer.^{8–14}

* Corresponding author. Department of Radiation Oncology, Inha University Hospital, Inha University School of Medicine #7-206, Shinheung-dong 3 Ga, Jung-ku, Incheon, 400-711, South Korea.

E-mail address: cancerovercome@gmail.com (HJ Kim).

Prostate-specific antigen (PSA) is a well-established biomarker for prostate cancer and available for monitoring response to treatment. In patients without androgen deprivation therapy (ADT), analysis of PSA kinetics after treatment may reveal the biologic effect of radiation on prostate cancer. The changes of PSA after radical prostatectomy, external beam radiotherapy, and brachytherapy have been researched extensively.¹⁵ A lower PSA nadir and a rapid decline in PSA after treatment have been related to improved clinical outcome.^{16–19} Unfortunately, kinetics of PSA decline following SBRT using Cyberknife remains poorly understood, and there are only a few reports from western countries.^{20,21} It is necessary to elucidate the kinetics of SBRT in Asian population. The purpose of the current study is to assess the kinetics of PSA for low- and intermediate-risk prostate cancer patients treated with SBRT using Cyberknife as both monotherapy and boost after WPRT in the absence of ADT.

2. Materials and methods

2.1. Patient characteristics

From 2008 to 2014, 61 patients newly diagnosed with low- and intermediate-risk (National Comprehensive Cancer Network definition) localized prostate cancer treated with SBRT using the Cyberknife robotic radiosurgery system (Accuray Inc.) were enrolled in this retrospective analysis. The treatment modality was chosen by the patient after extensive discussion of various treatment options, including surgery, SBRT, and IMRT. Of 61 patients, 32 were treated with SBRT as monotherapy and 29 received SBRT boost after WPRT. All patients had histologically confirmed primary adenocarcinoma of the prostate. None of these patients had received any other local or systemic primary treatment of prostate cancer. Prior transurethral resection of the prostate for urinary symptom relief was allowed. Patients were stratified according to 2014 National Comprehensive Cancer Network risk stratification guidelines.²² This study was approved by the Ethical Committee for Clinical Trials of our institution, and the retrospective data were prospectively collected from our institutional database.

In order to assess PSA kinetics in response to radiotherapy alone, patients were excluded if they failed therapy by the Phoenix definition.²³ All included patients had at least 1 year of follow-up. PSA bounce was defined as an absolute increase in PSA level of 0.2 ng/mL from the previous level, followed by a subsequent decrease.²⁴ Toxicity was documented at follow-up visits using the Radiation Therapy Oncology Group scale.

2.2. SBRT treatment planning and delivery

Four or more gold fiducial markers were implanted transperineally into the prostate. After 7 days, patients underwent magnetic resonance imaging and thin-cut computed tomography scan. Fused computed tomography and magnetic resonance images were used for treatment planning. The prostate, seminal vesicles, rectum, bladder, penile bulb, and bowel were contoured. The clinical target volume (CTV) included the prostate and proximal seminal vesicles. The planning target volume (PTV) equaled the CTV expanded by 3 mm posteriorly and 5 mm in all other dimensions. A prescription dose of 36.25 Gy, delivered in five fractions, was prescribed to the PTV. The prescription dose covered at least 95% of the PTV, normalized to the 75–85% isodose line [median homogeneity index of 1.27 (range, 1.24–1.41)]. The rectal dose–volume goals were as follows: <50% of the rectal volume receiving 50% of the prescribed dose, <20% receiving 80% of the dose, <10% receiving 90% dose, and <5%

receiving 100% of the dose. Treatments were given over 5 consecutive days. All SBRT treatment plans were generated on MultiPlan (version 2.2.0; Accuray Inc.).

2.3. WPRT and SBRT boost treatment planning and delivery

The prostate gland, seminal vesicles, and area of radiographic extracapsular extension were defined as CTV1. CTV2 included the external iliac nodes, internal iliac nodes, presacral nodes, and obturator nodes following the Radiation Therapy Oncology Group consensus.¹⁵ PTV1 was extended 7 mm beyond the CTV1 in all directions, except in the posterior direction, wherein it was extended 5 mm. PTV2 was extended 7 mm in all directions. The prescription dose of WPRT was 45 Gy, which was administered in 25 fractions. A minimum of 95% of the prescription dose was assured to cover 100% of the PTV. All WPRT treatment plans were generated on a Varian Eclipse treatment planning system (version 8.8.6; Varian Medical Systems, Palo Alto, CA, USA).

In SBRT boost planning, the prostate gland, seminal vesicles, and area of radiographic extracapsular extension were defined as the CTV, which was the same as that of the WPRT treatment plans. The PTV extended 5 mm beyond the CTV in all directions, except in the posterior direction, wherein it was extended 3 mm. A prescription boost dose of 21 Gy, delivered in three fractions, was prescribed to the PTV. The prescription dose covered at least 95% of the PTV, normalized to the 75–85% isodose line; median homogeneity index of 1.29 (range, 1.21–1.43)]. The rectal dose–volume goals were as follows: <50% of the rectal volume receiving 50% of the prescribed dose, <20% receiving 80% of the dose, <10% receiving 90% dose, and <5% receiving 100% of the dose. Treatments were given over 3 consecutive days.

2.4. Statistical analysis

To eliminate the effect of differing follow-up durations between SBRT monotherapy and SBRT boost after WPRT, we calculated the rate of change in PSA over time intervals from the completion of radiotherapy to 1 year, 2 years, 3 years, and 4 years post-treatment. The slope of PSA decline (ng/mL/mo) was calculated as the regression coefficient in a linear regression model for each individual.²⁵ The *t* test was performed to compare slopes of PSA and other mean values and analysis of variance in continuous variables.

3. Results

All patients completed the treatment. Sixty-one patients with a median 52.4-month (range, 14–74 months) follow-up were analyzed. The pretreatment median PSA level was 7.68 ng/mL (range, 3.45–19.50 ng/mL). Patient characteristics are summarized in Table 1.

Fig. 1 shows PSA changes over time, with a different rate of PSA decline for each time interval since the end of radiotherapy. To investigate the PSA kinetics after radiotherapy, the rate of PSA decline (slope) was calculated for four intervals following radiotherapy (0–1 year, 0–2 years, 0–3 years, and 0–4 years). The slope for all cohorts was maximal in the 1st year, but tapered off quickly in the following years, with median values of -0.43 ng/mL/mo, -0.22 ng/mL/mo, -0.13 ng/mL/mo, and -0.09 ng/mL/mo for durations of 1 year, 2 years, 3 years, and 4 years after radiotherapy, respectively. Distribution of the slope for SBRT monotherapy did not differ from that of the SBRT boost and WPRT through the years (Table 2). Owing to relatively homogenous inclusion criteria, there were no significant differences in the comparison of the rate of PSA decline by the Gleason score (≤ 6 vs. 7) and pretreatment PSA (≤ 10 vs. >10).

Download English Version:

<https://daneshyari.com/en/article/4274090>

Download Persian Version:

<https://daneshyari.com/article/4274090>

[Daneshyari.com](https://daneshyari.com)