



Original article – urothelial cancer

## Effectiveness of different diagnostic tools for upper urinary tract urothelial carcinoma

Yi-Sheng Tai<sup>a</sup>, I-Ni Chiang<sup>b, c, \*</sup>, Chao-Yuan Huang<sup>b</sup>, Huai-Chin Tai<sup>b</sup>, Yeong-Shiau Pu<sup>b</sup><sup>a</sup> Department of Urology, National Taiwan University Hospital Yun-Lin Branch, Yun-Lin, Taiwan<sup>b</sup> Department of Urology, National Taiwan University Hospital and College of Medicine, Taipei, Taiwan<sup>c</sup> Institute of Biomedical Engineering, College of Medicine and College of Engineering, National Taiwan University, Taipei, Taiwan

## ARTICLE INFO

## Article history:

Received 14 February 2014

Received in revised form

13 May 2014

Accepted 28 July 2014

Available online 26 September 2014

## Keywords:

cytology

intravenous urography

retrograde pyelography

upper urinary tract urothelial carcinoma

ureterorenoscopy

## ABSTRACT

**Objective:** The results of urinalysis, radiographic studies, urinary cytology examinations, and ureterorenoscopy (URS) biopsies, as well as the results of histopathology can be used to establish a diagnosis of upper urinary tract urothelial carcinoma (UTUC).

**Materials and Methods:** We enrolled 99 patients who underwent radical nephroureterectomy (RNU) during the period 2003–2007. A total of 65 random urine and 83 URS washing cytology examinations, 48 intravenous urography (IVU), 59 retrograde pyelography (RP), and 81 URS biopsy results were available prior to RNU and were compared with the pathological grades and stages of these surgical specimens.

**Results:** Ninety-three UTUCs were found among the 99 RNU specimens. Initial presentations and urinalysis results could not predict tumor stages. The patient with preoperative pyuria was significantly associated with high-grade UTUC (75.0% vs 52.6%,  $p = 0.031$ ). Random urine and URS washing cytology results could not predict tumor grades or stages. The sensitivity of 3-day random urine cytology was significantly better than 2-day and 1-day examinations ( $p = 0.002$  and  $p = 0.019$ , respectively). The abnormal findings in IVU and RP accounted for 89.4% and 100%, respectively. Non-enhancement of images was significantly associated with high tumor grading ( $p = 0.01$ ). URS biopsy ( $n = 72$ ) was positive for malignancy in 52 patients (69.3%). Biopsy grade had a significant correlation with surgical tumor grade ( $\kappa = 0.649$ ) and high-grade biopsy results were significantly associated with invasive tumor stage (pT2–T4) ( $p = 0.004$ ).

**Conclusion:** Combining random urine cytology for 3 nonconsecutive days, upper urinary tract images, and URS biopsies provided an accurate diagnosis of UTUC. This study found that preoperative pyuria in urinalysis, non-enhancement in IVP or RP, and high-grade tumor in URS biopsy could predict high-grade tumor in RNU specimens.

Copyright © 2015, Taiwan Urological Association. Published by Elsevier Taiwan LLC.

Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

### 1. Introduction

The incidence of upper urinary tract urothelial carcinoma (UTUC) is high in Taiwan.<sup>1</sup> Worldwide, UTUC, including carcinomas of the renal pelvis and ureter, only accounts for 4–5% of all urothelial cancer.<sup>2,3</sup> However, in Taiwan, the ratio of renal pelvic urothelial carcinoma (UC) to ureteral UC, and to bladder UC is 1.2:1:6.7, which means UTUC is close to 25%.<sup>4</sup> Previous studies showed that 31.2–34.1% of UTUC patients developed metachronous bladder UC after radical nephroureterectomy (RNU) and bladder cuff resection, the gold standard for treatment of UTUC.<sup>5,6</sup>

A diagnosis of upper urinary tract tumor is based on results of urinalysis, urine cytology examination, upper urinary tract

imaging, endoscopic inspection, and biopsy for histopathological analysis.<sup>7,8</sup>

We retrospectively reviewed patients receiving RNU and bladder cuff resection because of suspected upper urinary tract tumor after the ureterorenoscopy (URS) examinations, and investigated the sensitivities of urinalysis, random urine cytology, upper urinary tract washing cytology, intravenous urography (IVU), retrograde pyelography (RP), URS inspection, and biopsy in the detection of UTUC in Taiwan. We also correlated these diagnostic tools with the pathological staging and grading of UTUC to determine the effectiveness of these examinations.

### 2. Methods

From 2003 to 2007, 99 patients received RNU and bladder cuff resection after URS examination in our institution. None of the patients had a history of bladder or upper urinary tract UC. All of

\* Corresponding author. Department of Urology, National Taiwan University Hospital, Room 11, Number 7, Chung Shan South Road, Taipei 10016, Taiwan.

E-mail address: [inichiang@gmail.com](mailto:inichiang@gmail.com) (I.-N. Chiang).

the patients submitted a urine specimen for dipstick and microscopic analysis prior to URS.

Microscopic hematuria was defined as red blood cell counts >5 under high-power field and pyuria was defined as white blood cell >5 under high-power field. Proteinuria was defined as urine concentration of protein of  $\geq 30$  mg/dL.

Other surveillance techniques included random urine cytology, IVU, RP, computed tomography (CT), magnetic resonance image (MRI), URS washing cytology, and URS biopsy. Positive findings of random urine cytology and washing cytology included at least one specimen with suspicious malignant cells or atypical cells. The abnormal findings of the image studies included obstructive uropathy, urinary tract filling defect, and visible tumors in the radiographic examinations.

The combination of these surveillance tools depended on the patients' conditions and the physicians' choices. Of the 99 patients, 99 urinalysis results, 65 random urine cytology results, 48 IVU results, 59 RP results, 81 CT results, 17 MRI results, 99 URS examinations, 83 URS washing cytology results, and 81 biopsy results were available.

The  $\chi^2$  test was used to analyze associations between diagnostic tools and pathological parameters. The interobserver agreement between URS and RUN pathological results was evaluated using the Cohen's  $\kappa$  method. A  $p$  value <0.05 was considered statistically significant.

### 3. Results

Of the 99 patients who received RNU and bladder cuff resection due to diagnosis or suspicion of UTUCs, six had RNU specimens without UC. Two of the patients had URS biopsy specimens that revealed superficial low-grade UC without lamina propria invasion. Two of the patients had renal cell carcinomas. The other two patients had biopsy specimens that showed mesenchymal polyps and papilloma, respectively.

Ninety-three patients had pathologically proven UC in the upper urinary tract, including 49 women (52.7%) and 44 men (47.3%). The median age of all UTUC patients was 69 years (interquartile range, 58–74 years). UC was found on the left side in 47 patients (50.5%), on the right side in 44 patients (47.3%), and on both sides in two patients (2.2%). UC was located only at the renal pelvis in 40 patients (43.0%), only at the ureter in 31 patients (33.3%), and at both renal pelvis and ureter in 22 patients (23.7%).

A total of 93 urinalysis results, 62 random urine cytology results, 47 IVU results, 54 RP results, 78 CT results, 16 MRI results, 93 URS examinations, 77 URS washing cytology results, and 75 biopsy results were available in the cohort of 93 UTUC patients.

#### 3.1. Initial presentations and urinalysis

Initial presentations are listed in Table 1. The most common symptom was gross hematuria in 71 patients (76.3%). Eleven patients (11.5%) visited urology clinics because abnormal findings had been disclosed at hospitals elsewhere, including microscopic hematuria, renal function impairment, anemia or hydronephrosis, and ultrasonographic evidence of a renal mass.

Urinalysis was obtained prior to URS studies and showed microscopic hematuria in 67.7% ( $n = 63$ ), pyuria in 38.7% ( $n = 36$ ), and proteinuria in 34.4% ( $n = 32$ ) of patients (Table 1). Initial presentations and urinalysis results could not predict tumor staging. Interestingly, patients with pyuria had a significantly higher percentage of high-grade UC (75.0% vs 52.6%,  $p = 0.031$ ).

**Table 1**

Initial presentations and urinalysis in upper urinary tract urothelial carcinoma patients.

	Results	$n$ (%)
Initial presentation ( $n = 93$ )	Gross hematuria	71 (76.3)
	Flank pain or soreness	22 (23.7)
	Abdominal discomfort	5 (5.3)
	Abnormal findings in health examination <sup>a</sup>	11 (11.5)
	Body weight loss	3 (3.2)
Urinalysis ( $n = 93$ )	Recurrent UTI	2 (2.2)
	Microscopic hematuria	63 (67.7)
	Pyuria	36 (38.7)
	Proteinuria	32 (34.4)

UTI = urinary tract infection.

<sup>a</sup> Abnormal findings in health examination included microscopic hematuria, hydronephrosis, renal mass, elevated creatinine, and anemia.

#### 3.2. Cytology

Random urine cytology and URS washing cytology results were available for 62 patients and 77 patients, respectively. The sensitivity of random urine cytology and URS washing cytology was 58.1% and 58.4%, respectively (Table 2). Thirty-nine patients submitted random 3-day urine cytology specimens, nine patients submitted 2-day results, and 14 patients submitted 1-day results. The sensitivity of 3-, 2-, and 1-day collection of random urine cytology results was 74.4%, 33.3%, and 28.6%, respectively. Three-day random urine cytology was significantly better than 2- and 1-day urine cytology ( $p = 0.002$  and  $p = 0.019$ , respectively); however, there was no difference between 2- and 1-day urine cytology sensitivity ( $p = 0.809$ ).

The contributions of high pathological grade and invasive tumor stage stratified by results of random urine cytology and upper urinary tract washing urine cytology are listed in Table 2. There were no associations between cytology collecting methods or predictions of tumor grading and staging.

#### 3.3. Upper urinary tract imaging

Of the 47 patients with IVU prior to URS examinations, 19.1% had hydronephrosis, 19.1% had non-enhancement of upper urinary tract, 51.1% had urinary tract filling defects, and 10.6% had no definite abnormal findings. A total of 54 patients received RP following URS studies. The procedures showed hydronephrosis in 7.4%, non-enhancement of upper urinary tract in 3.7%, urinary tract filling defects in 88.9%, and no definite abnormal finding in any of the patients.

The contributions of pathological results in IVU and RP characteristics are demonstrated in Table 3. Non-enhancement in the upper urinary tract group had significant superiority over the no definite abnormal finding group in predicting high tumor grading ( $p = 0.01$ ). The results of CT were available in 78 patients and showed simply hydronephrosis, suspected upper urinary tract tumors, and no definite abnormal findings in seven (9.0%) patients, 70

**Table 2**

Urine cytological and pathological results.

	Cytology results, $n$ (%)	High-grade UC (%)	Invasive tumor (%)
Random urine cytology ( $n = 62$ )	Negative 26 (41.9)	50.0	$p = 0.121$ 42.3
	Positive 36 (58.1)	69.4	$p = 0.96$ 41.7
URS washing cytology ( $n = 77$ )	Negative 32 (41.6)	46.9	$p = 0.053$ 34.4
	Positive 45 (58.4)	68.9	$p = 0.1$ 53.3

UC = urothelial carcinoma; URS = ureterorenoscopy.

Download English Version:

<https://daneshyari.com/en/article/4276333>

Download Persian Version:

<https://daneshyari.com/article/4276333>

[Daneshyari.com](https://daneshyari.com)