



## Original Research

## Prognostic value of Tissue Transition Projection 3D transparent wall CT reconstructions in bowel ischemia



Marco Moschetta<sup>a, \*</sup>, Arnaldo Scardapane<sup>b</sup>, Michele Telegrafo<sup>b</sup>, Nicola Maria Lucarelli<sup>b</sup>,  
Valentina Lorusso<sup>b</sup>, Giuseppe Angelelli<sup>b</sup>, Amato Antonio Stabile Ianora<sup>b</sup>

<sup>a</sup> DETO – Department of Emergency and Organ Transplantation, University of Bari Medical School, Piazza Giulio Cesare 11, 70124 Bari, Italy

<sup>b</sup> DIM – Interdisciplinary Department of Medicine, Section of Diagnostic Imaging, Aldo Moro University of Bari Medical School, Piazza Giulio Cesare 11, 70124 Bari, Italy

## HIGHLIGHTS

- CT represents the main imaging technique for diagnosing bowel ischemia.
- Bowel dilatation represents a CT sign indicating poor prognosis.
- We assess the role of TTP 3D reconstruction for evaluating loop dilatation.
- TTP 3D reconstructions are a rapid and automatic tool in this field.
- A prognostic value could be proposed for TTP 3D imaging.

## ARTICLE INFO

## Article history:

Received 3 May 2016

Accepted 29 August 2016

Available online 31 August 2016

## Keywords:

Computed tomography

MDCT

Bowel ischemia

Bowel dilatation

## ABSTRACT

**Background:** Multi-detector computed tomography (MDCT) represents the gold standard in patients with acute abdomen syndrome and suspected bowel ischemia. It provides a correct diagnosis and contributes to appropriate treatment planning. This study aims to evaluate the role of 3D Tissue Transition Projection (TTP) transparent wall CT reconstruction for detecting the degree of bowel dilatation and to correlate this finding with the aetiology and prognosis in patients affected by mesenteric infarction.

**Materials and methods:** Forty-seven patients affected by bowel infarction due to vascular obstruction (arterial in 66% of cases, venous in 34%) were assessed by MDCT examination searching for the degree of bowel dilatation (subdivided into 4 groups: entire small bowel (SB);  $\geq 50\%$  of SB;  $< 50\%$  of SB; large bowel only). Two blinded radiologists evaluated TTP 3D transparent wall and multi-planar reconstructions. Chi square test was used to correlate CT findings with the disease course and the mortality rate. Cohen's kappa statistics was used in order to assess inter-observer agreement.

**Results:** The overall mortality rate was 64%, with a 90% value for arterial forms and 10% in case of venous infarctions. The entire SB ( $n = 10$ ) or a  $\geq 50\%$  SB dilatation ( $n = 16$ ) correlated with poor prognosis in all cases ( $p < 0.05$ ); a  $< 50\%$  SB dilatation ( $n = 16$ ) correlated with good prognosis in 87.5% of cases ( $p < 0.05$ ). A large bowel only dilatation ( $n = 5$ ) did not show a significant prognostic value ( $p = 0.13$ ). Almost perfect agreement between the two readers was found ( $k = 0.84$ ).

**Conclusion:** MDCT offers different reconstruction software for diagnosing bowel ischemia. 3D TTP transparent wall reconstructions represent a rapid and automatic tool for identifying loop dilatation, which significantly correlates with an arterial aetiology and poor prognosis.

© 2016 Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd.

## 1. Introduction

Acute bowel ischemia represents an abdominal emergency accounting for nearly 1% of patients presenting with acute abdomen syndrome. It occurs when blood flow to the bowel loops is

\* Corresponding author.

E-mail address: [marco.moschetta@gmail.com](mailto:marco.moschetta@gmail.com) (M. Moschetta).

compromised because of mesenteric arterial hypo-perfusion, impaired venous drainage or bowel obstruction [1–5]. In fact, the causes of intestinal ischemia can be occlusive or non occlusive. Occlusive causes are due to the embolic or thrombotic occlusion of arterial or venous vessels and account for about 80% of all cases of intestinal ischemia [6,7]. Non occlusive causes account for about 20–30% of all cases of intestinal ischemia.

Bowel ischemia could represent a transient and reversible event if diagnosed in an early phase; however, it may lead to intestinal infarction that requires surgical or interventional management [6]. Despite the new available imaging tools, mortality rate from intestinal infarction still remains very high, with reported value ranging from 60% to 90% [8–10].

Multi-detector computed tomography (MDCT) represents the main imaging technique that must be promptly performed in all patients with acute abdomen syndrome and suspected bowel ischemia. In fact, MDCT can provide a correct diagnosis and contributes to appropriate treatment planning being able to define the nature of the disease and the extent of anatomical damage [6,7].

Several CT signs have been described in this field and recently they have also been correlated to the prognosis of the disease giving an important prognostic value to this imaging tool.

Bowel loop dilatation represents one of these signs; it has been related to an increased capillary permeability within bowel walls and usually reflects an advanced stage of disease. Besides, bowel loop dilatation is typical of the arterial forms and characterized by high mortality rates [7].

The introduction of multi-detector CT devices has highly improved the diagnostic and prognostic potential for detecting CT signs of bowel ischemia with the possibility of using different image reconstruction software starting from isotropic voxels obtained by CT scanning.

Tissue Transition Projections (TTP) represent a particular sub-type of Surface Shaded Volume Rendering (SS-VRT) that can be used for this purpose [11–15]. So far, no experience has been reported in this field.

This study aims to evaluate the role of 3D TTP transparent wall CT reconstruction for detecting the degree of bowel dilatation and to correlate this finding with the aetiology and prognosis in patients affected by mesenteric infarction.

## 2. Material and methods

### 2.1. Patients

We retrospectively reviewed 47 patients with bowel infarction who were examined at our department between January 2012 and May 2013. Patients were identified through a search of our electronic archive system using terms such as ischemia, infarction and intestinal necrosis. All 47 selected cases were affected by occlusive ischemia: 31 (66%) of the arterial and 16 (34%) of the venous type. Patients included 26 men (55%) and 21 women (45%), with age ranging from 46 to 86 years (mean age 69.3). Thirty of the 47 patients (64%) had died, and 17 (36%) were alive 6 months after the diagnosis. Only 30 patients (64%) underwent surgery; of these, 18 (60%) died and 12 (40%) survived. The definite diagnosis was provided by surgery, with histology of the resected intestinal segment in 30 patients. In the remaining 17 cases, the diagnosis was based on radiological and clinical data and on the disease course. All patients were examined in an emergency setting due to sudden onset of acute abdomen.

### 2.2. CT protocol

The CT scanners used were a 320-slice MDCT device (Aquilion

One, Toshiba Medical System, Nasu, Japan; protocol: detector collimation  $320 \times 0.5$  mm, increment 0.5, rotation time 0.5 s, mean kVp/mAs: 120/200) and a 16-slice MDCT (TSX- 101A- Aquilion 16, Toshiba Medical System, Tochigi, Japan; protocol: detector collimation  $16 \times 1$  mm; increment 0.8 mm; rotation time 0.5 s; mean kVp/mAs 120/250). In all cases, scans were acquired before and after the intravenous injection of contrast material (120–140 ml injected at a rate of 3–3.5 ml/s), with image acquisition in the arteriolar phase, generally with a delay of 30–40 s from the beginning of contrast-agent administration, and in the venous phase, with a delay of 60–90 s from the beginning of contrast-agent administration. The data sets were retrospectively analyzed on a workstation (HP XW 6400) equipped with image reconstruction software (Vitrea 4.1, Vital Images, Minneapolis, MN, US). 3D TTP transparent wall software and multiplanar (MPR) reconstructions were used.

### 2.3. Image analysis

3D TTP transparent wall reconstructions and MPR images were assessed by two blinded radiologists with more than 5 years' experience in the field of abdominal CT imaging and CT image reconstruction, searching for the following parameters:

- bowel-loop dilatation, defined as a small-bowel diameter  $>2.5$  cm and large-bowel diameter  $>8$  cm; basing on the degree of bowel dilatation, patients were subdivided into 4 groups:
  - entire small bowel (SB) in the case of both jejunal and ileal involvement;
  - $\geq 50\%$  of SB in the case of predominant jejunal or ileal involvement associated with some ileal or jejunal dilated loops, respectively;
  - $< 50\%$  of SB in the case of exclusive involvement of some jejunal or ileal loops;
  - large bowel only;
- the morphology and patency of the coeliac trunk, superior and inferior mesenteric arteries, superior mesenteric vein and portal trunk, in order to define the arterial or venous nature of the vascular occlusion causing ischemia.

3D TTP transparent wall reconstructions were used for evaluating the degree of bowel loop dilatation while MPR images for establishing the aetiology of the disease.

Post-processing time was approximately 10 min.

### 2.4. Statistical analysis

The sample population was initially analyzed using descriptive statistics and considering age, gender, aetiology of ischemia (arterial or venous), disease course (dead or living patient) and calculating mortality rates. The groups of patients classified basing on bowel loop dilatation as detected by 3D TTP reconstructions were correlated with treatment outcomes (dead vs living) and with the aetiology of the disease (arterial vs. venous infarctions). Statistical significance of this information was verified by applying the chi-square test and calculating *p* values. The obtained data were considered to be statistically significant at a *p* value  $\leq 0.05$ . Cohen's kappa statistics was used in order to assess inter-observer agreement. A *k* value of more than 0.81 was considered to represent almost perfect agreement and values of 0.61–0.80 and 0.41–0.60 to represent substantial and moderate agreement, respectively. All calculations were performed using NCSS2007\_ statistical software.

Download English Version:

<https://daneshyari.com/en/article/4285278>

Download Persian Version:

<https://daneshyari.com/article/4285278>

[Daneshyari.com](https://daneshyari.com)