



Review

An analysis of surgical anatomy of the gastric fundus in bariatric surgery: Why the gastric pouch expands? A point of technique



Radwan Kassir ^{a,*}, Pierre Blanc ^b, Patrice Lointier ^c, Olivier Tiffet ^a, Christophe Breton ^b, Imed Ben Amor ^d, Antonio Iannelli ^d, Jean Gugenheim ^d

^a Department of Bariatric Surgery, CHU Hospital, Jean Monnet University, Saint Etienne, France

^b Department of Bariatric Surgery, Clinique Chirurgicale Mutualiste de Saint Etienne, France

^c Department of Digestive Surgery, Clinique de la Chataigneraie, Beaumont, France

^d Department of Bariatric Surgery, Archet 2 Hospital, University Hospital of Nice, Nice, France

HIGHLIGHTS

- It is essential to completely release the Fundus in bariatric surgery in order to create a narrow gastric pouch.
- Dissection during bypass prevents any “rocking” movement from above downwards and from lateral to medial.
- This minimal dissection is compensated by further dissection at the level of the left pillar.
- The greater tuberosity has no electrical activity and therefore no mechanical activity.
- This would appear logical as this area does not contain food but generally air.

ARTICLE INFO

Article history:

Received 7 August 2014

Received in revised form

21 August 2014

Accepted 31 August 2014

Available online 16 September 2014

Keywords:

Gastric fundus

Surgical anatomy

Bariatric surgery

Hiatus hernia

Diaphragmatic pillar

ABSTRACT

In bariatric surgery, it is essential to completely release the Fundus in order to create a narrow gastric pouch. The upper part of the fundus is located above the omental bursa and is therefore retro-peritoneal. In order to release this completely, not only does the arterial supply to the fundus need to be divided to visualise the left diaphragmatic pillar, but the right attachment beginning at the left diaphragmatic pillar and running towards the fundus needs to be divided. This minimal dissection is compensated by further dissection at the level of the left diaphragmatic pillar and traction on the stomach from right to left during the final division stapling division process. The surgeon still has the impression of having released the posterior aspect of the Fundus, exposing the pillar of the diaphragm, although in fact part of the Fundus still remains adherent to the diaphragm and is therefore not released.

© 2014 Surgical Associates Ltd. Published by Elsevier Ltd. All rights reserved.

1. Introduction

In principle, the anatomy of the stomach is not complicated. Although knowledge of the anatomy of the gastric Fundus is extremely useful in bariatric surgery, this is only very rarely clearly described. The aim of this article is to describe the anatomical features of the gastric Fundus which may explain some operation

difficulties, some “disappointing” postoperative radiological appearances and finally, which allow to discuss the surgical strategy for the treatment of severe obesity.

2. Technical point

It is essential to completely release the Fundus in bariatric surgery in order to create a narrow gastric pouch both in sleeve gastrectomy and in gastric bypass. Dissection may be incomplete in gastric bypass if the posterior gastric dissection which is minimal does not pass to the right of the posterior gastric vessels (Fig. 1). This minimal dissection is generally compensated by

* Corresponding author. Department of Obesity Surgery, CHU Hospital, Jean Monnet University, Avenue Albert Raimond, 42270 Saint Etienne, France.

E-mail address: Radwankassir42@hotmail.fr (R. Kassir).




Fig. 1. Technical point. Laparoscopic view.

further dissection at the level of the left pillar and traction on the stomach from right to left during the final division stapling division process (Fig. 2). In revision surgery (after a ring or anti-reflux surgery) this release is impossible. The surgeon still has the impression of having released the posterior aspect of the Fundus, exposing the pillar of the diaphragm, although in fact part of the Fundus still remains adherent to the diaphragm and is therefore not released. This is due to the inflammation (created by the foreign body or creation of a valve) passing above the omental bursa and abolishing the avascular dissection plane. This explains why gastric pouches may be too wide (incomplete dissection) and the increased complication rate after revision bariatric surgery (when the tissues are more fragile) (Figs. 3 and 4) [1]. A too wide gastric pouch may explain insufficient weight loss as this part of the stomach may dilate over time because of the lack of electrical and mechanical activity at this level. There is always a risk after an initial procedure of having to perform a second one in the long term, as bariatric surgery is “counter to nature” and nature always seeks to rectify itself. It is therefore a form of surgery which changes over time and the anatomy explains why surgical management of obesity should begin with correct dissection of the fundus, either with a sleeve gastrectomy or gastric bypass and not routinely with a ring as a first line choice, particularly in massive obesity when the risk of failure is greater as more kilograms need to be lost [2–3].

Malrotation abnormalities are extremely rare, although these must be looked for on opacified preoperative gastrointestinal imaging [4]. This is also used to investigate for abnormalities at the gastro-esophageal junction (hiatus hernia). The upper part of the fundus is located above the omental bursa and is therefore retroperitoneal (Fig. 5). In order to release this completely, not only does the arterial supply to the Fundus need to be divided to visualise the left pillar but the right attachment beginning at the left diaphragmatic pillar and running towards the fundus needs to be divided. If the fundus is divided and stapled, the staple lengths must not be too long at this point. The consensus conference statement recommends 2.5 mm staples for the first surgery and at least 4.8 mm staples for revision surgery (when the tissues are thicker). Four to six 60 mm cartridges should be available in the operating theatre for a sleeve gastrectomy [5]. Longer staples may be required for the antrum (green cartridges). The staples may be shorter for gastric bypass when two to four 60 mm cartridges should be available in the operating theatre (horizontal division 6 cm from the cardia) [6].

3. Discussion

The stomach is visible from the 4th week of pregnancy. The dorsal part (the future greater curvature) develops more quickly during growth than the ventral part (the future lesser curvature). It

Download English Version:

<https://daneshyari.com/en/article/4286353>

Download Persian Version:

<https://daneshyari.com/article/4286353>

[Daneshyari.com](https://daneshyari.com)