



Original research

Perioperative care in elderly patients undergoing thyroid surgery



Salvatore Guarino^{*}, Carla Di Cosimo, Carlo Chiesa, Alessio Metere, Valerio Di Bella, Angelo Filippini, Laura Giacomelli

Department of Surgical Sciences, "Sapienza" University of Rome, Viale Regina Elena 324, 00161 Roma, Italy

ARTICLE INFO

Article history:

Received 15 May 2014

Accepted 15 June 2014

Available online 23 August 2014

Keywords:

Thyroid surgery
Elderly

ABSTRACT

The features of western world population are rapidly changing. The increment geriatric population obliges clinicians to implement specific recommendations and guidelines to manage these patients.

In the field of thyroid surgery, when indications are represented by benign conditions, surgeons and endocrinologists tend to avoid surgery for the increased perioperative risks in the over 70 year old population.

We reviewed our experience in thyroid surgery in geriatric patients within the environment of a "week surgery unit". This unit was conceived to offer a highly specialized setting for thyroid patients needing short stay after surgery.

Results showed that the surgical outcomes were comparable to the ones from third surgery in young patients.

The week surgery approach is the best and safest formula to offer to the geriatric population needing thyroid surgery.

© 2014 Surgical Associates Ltd. Published by Elsevier Ltd. All rights reserved.

1. Introduction

The definition of geriatric population may strongly vary in medical literature. Different authors used 65, 70 or 75 years as cut off [1]. Western world population is rapidly changing. According to the USA Census Bureau, the increased life expectancy will lead to an approximately 80 millions of elderly population by 2050.

The prevalence of thyroid nodules increases with age; almost 50% of patients ≥ 65 years demonstrate nodules on ultrasound examination [2].

It has been estimated that 90% of thyroid glands in women over the age of 70 will contain nodules and 80% of the glands of men over the age of 80 will be nodular [3].

The incidence of thyroid cancer also increases with age [4,5]. Elderly patients often present with more aggressive forms of

thyroid cancer, larger tumours, more extensive local growth, or distant metastases [6–8].

The increasing number of the geriatric population obliges clinicians to establish guidelines and protocols to standardize the optimal care for these patients.

Several studies have confirmed that, in general, elderly is not a contraindication to elective thyroid surgery. Nevertheless the prevailing approach among surgeons has long been to avoid elective surgery on the elderly, as they tend to have higher rates of comorbidity, longer postoperative stays, and poorer long-term outcomes [9].

Because of an elevated risk for perioperative morbidity among elderly patients undergoing surgical procedures, indications for thyroidectomy in this population are often restricted to overt compressive symptoms or a strong suspicion for malignancy [10].

In the era of the "fast track" applied to surgical patients, paramount importance is given to the optimization of the perioperative care in order to avoid unnecessary money and time waste aiming to obtain the best cost-effectiveness formula.

In this scenario where different combinations of short stay hospitalization have been attempted in thyroid surgery [11–13], hereby we report our experience of "week surgery" approach to thyroid surgery applied also to the geriatric population.

Abbreviations: USA, United States of America; IV, intravenous; NSAID, non-steroidal anti-inflammatory drugs; ASA, American society of anaesthesiologists; FNA, fine needle aspiration; ICU, intensive care unit; TDS, Ter Die Sumendum.

^{*} Corresponding author.

E-mail addresses: salvatore.guarino@gmail.com (S. Guarino), carla.dicosimo@gmail.com (C. Di Cosimo), carlochiesa8@gmail.com (C. Chiesa), alessio.metere@gmail.com (A. Metere), valeriodibella@hotmail.it (V. Di Bella), angelo.filippini@uniroma1.it (A. Filippini), laura.giacomelli@uniroma1.it (L. Giacomelli).

2. Methods

In our Department of Surgical Science approximately 80 thyroid procedures per year are performed.

Since January 2011 a short stay “week surgery” Unit was established to accommodate patients undergoing thyroid and parathyroid procedures. This unit was conceived to offer a service of excellence to thyroid patients within a highly specialized medical environment and in order to reduce costs of unnecessary hospitalizations in regular inpatients wards.

2.1. Week surgery pathway

2.1.1. Preoperative assessment

Patients seen in the Outpatient Clinic with an established indication for surgery were given an appointment for the pre-assessment clinic within 15 days. The patient attending the pre-assessment unit was seen by one of the junior member of the Surgical Team and by a senior Anaesthetist after having performed blood test (full blood count, biochemistry, thyroid function test), chest X-ray, Pulmonary function test. Those tests were aimed to reveal any relevant condition that might have raised the surgical or anaesthetic risk. Vocal cords were routinely assessed by indirect laryngoscopy to detect pre-existing dysfunctions. Some of the patients were sent to have pre-operative Q-Elastosonography of the thyroid nodules as predictor of malignancy [14].

2.1.2. Surgery

Patients were admitted on the same day of the surgery having been starved from the night before to solid food and to clear fluid till two hours before surgery. This would have typically happened at the beginning of the week (either on a Monday or a Tuesday). Total or subtotal thyroidectomies were performed in a standard fashion without the use of energy devices under general anaesthetic [15]. Extreme importance was given to the recognition and preservation of parathyroid glands, of the external branch of the superior laryngeal nerves and of the recurrent nerves. One suction drains were left bilaterally in the thyroid lodge. Closure of the cervicotomy was typically performed with subcuticular absorbable suture [16]. Thyroid specimen with cancer was sent for biomolecular study in order to detect BRAF mutations as predictor for possible target therapy [17].

2.1.3. Post-operative care

Patients undergoing surgery in the morning were normally let eating and drinking on the same day few hours after the end of the surgical procedure. As required analgesia was prescribed using either IV paracetamol or NSAID tablets. Calcium supplements and Vitamin D3 were not routinely prescribed. In case of acute hypocalcaemic crisis, IV Calcium gluconate was administered intravenously. Calcium level in the blood was tested everyday starting from the morning after surgery. The suction drain was removed on day 2 post op.

2.1.4. Discharge criteria

Patients were deemed dischargeable, typically on day 2 or day 3 after surgery:

- in absence of complications (hypocalcaemic crisis, haematoma, cardio-respiratory complications)
- with a level of blood calcium >7 mg/dl
- pain controlled with oral medication
- patients compliant to be discharged

Those patients that did not fulfil the discharge criteria by the Friday of the same week, were transferred to the normal surgical elective ward as the “week surgery” unit would have closed by the Friday evening. Follow up and post surgical care were arranged alongside with the endocrinologist colleagues to detect the best medical formula [18].

3. Our experience

Since the implementation of the week Surgery in February 2011, 167 patients were admitted and underwent thyroid surgery. Of those 15 were above 70 year old. Of the over-70 year old 4 were male and 11 were female. The indications for surgery within the geriatric population were: cancer, papillary proliferation, compressive symptoms, uncontrolled hyperthyroidism (relapses/non responders) (Table 1).

We examined the outcomes of the geriatric population undergoing “week thyroid surgery”.

The group of over 70 year old patients was quite heterogeneous in terms of co-morbidity. Their operative risk was stratified using the American Association of Anaesthetists score (ASA) (Table 2).

The outcomes considered were:

- intraoperative complication rate (difficult intubation, intra-operative cardiac events, need for postoperative intensive care)
- short term postoperative complication rate (hypocalcaemia, cardiorespiratory complications, infections, haematoma, early recurrent nerve palsy)
- long term postoperative complication rate (permanent hypocalcemia, permanent recurrent nerve palsy)
- need for reoperation
- need to stay beyond the day 3 after surgery
- 30 day readmission rate
- 30 day mortality rate

4. Results

All of the patients underwent total thyroidectomy. Of the three cancer cases, all of them had the lymph-node dissection of the central compartment, and only one had associated also a radical neck dissection for malignant lymph-nodes on the preoperative cytology. The majority of geriatric patients were in the group of ASA 2 for the presence of comorbidity such as diabetes, hypertension and dyslipidaemia.

Four patients were scored ASA 3 for symptoms due to previous ischaemic cardiac and cerebral events. Only one patient was scored ASA 4 for the co-existence of ischaemic cardiac disease and chronic pulmonary condition. This 73 year old patient, that was affected by a single thyroid nodule resulting thy 5 on the FNA, was declared high risk for surgery. After discussion, decision was taken to carry on the procedure with a possible need for postoperative ICU bed, but this was not necessary and the patient had an uneventful recovery on the regular week surgery unit.

Examining the outcomes of the geriatric population undergoing thyroid surgery in our institution (Table 3) no long term

Table 1
Showing the indications for thyroidectomy in the geriatric patients.

Indication	Number	%
Thy 3	5	33%
Compressive symptoms	4	26%
Uncontrolled hyperthyroidism	3	20%
Cancer	3	20%

Download English Version:

<https://daneshyari.com/en/article/4286585>

Download Persian Version:

<https://daneshyari.com/article/4286585>

[Daneshyari.com](https://daneshyari.com)