



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Transient epiphyseal lesion of the femoral head after traumatic hip dislocation: A case report[☆]



Yusuke Kubo^a, Takuaki Yamamoto^{b,*}, Goro Motomura^a, Satoshi Kido^c,
Kazuyuki Karasuyama^a, Kazuhiko Sonoda^a, Yukihide Iwamoto^d

^a Department of Orthopaedic Surgery, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan

^b Department of Orthopaedic Surgery, Faculty of Medicine, Fukuoka University, 7-45-1 Nanakuma, Jonan-ku, Fukuoka 814-0180, Japan

^c Department of Orthopaedic Surgery, Yamaguchi Red Cross Hospital, 53-1 Yahatababa, Yamaguchi 753-8519, Japan

^d Department of Orthopaedic Surgery, Kyushu Rosai Hospital, 1-3-1, Kuzuharatakamatsu, Kokuraminami-ku, Kitakyushu 800-0296, Japan

ARTICLE INFO

Article history:

Received 28 February 2016

Received in revised form 2 April 2016

Accepted 3 May 2016

Available online 9 May 2016

Keywords:

Hip dislocation

Bone marrow edema

Insufficiency fracture

ABSTRACT

INTRODUCTION: We experienced a rare case in which magnetic resonance imaging (MRI) showed a transient epiphyseal lesion of the femoral head four months after traumatic hip dislocation. To our knowledge, there have been no previously published reports on the development of such transient lesions after traumatic hip dislocation involving no abnormalities just after dislocation.

PRESENTATION OF CASE: We report a 22-year-old man who showed a transient epiphyseal lesion of the femoral head after traumatic hip dislocation. On MRI performed two days after dislocation, no bony injuries were observed around the hip joint. Four months after dislocation, the patient suddenly experienced right hip pain without any new trauma or injury. A low-intensity band convex to the articular surface was apparent above the epiphyseal scar on T1-weighted imaging, and bone marrow edema was observed around the band lesion on short-tau inversion recovery imaging. Following a two-month period of non-surgical conservative therapy, the patient's hip pain resolved and the low-intensity band was no longer observed on follow-up MRI.

DISCUSSION: Although the detailed pathogenesis of this transient changes was unclear, we speculate that prolonged rest after traumatic hip dislocation may contribute to bone insufficiency, resulting in an insufficiency fracture of the femoral head.

CONCLUSION: This study suggests that transient epiphyseal lesions of the femoral head may occur in patients with a history of traumatic hip dislocation associated with an adapted long-term rest.

© 2016 The Author(s). Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Traumatic hip dislocation may have a number of sequelae, one of which is osteonecrosis of the femoral head (ONFH) [1–4]. The incidence of ONFH after traumatic dislocation has been reported to range from 10 to 40%, and this complication generally causes hip pain between 12 and 24 months after traumatic dislocation [5–10]. Magnetic resonance imaging (MRI) is commonly used to diagnose ONFH, demonstrating characteristic low-intensity bands on T1-weighted imaging [11,12]. It has been reported that the diagnosis of ONFH may be confidently made if an abnormal signal is observed on MRI performed three months after the injury [8].

On the other hand, a previous study reported the appearance of transient bone marrow changes in the femoral head on MRI early after traumatic hip dislocation [8]. Although the cause of these transient changes was unclear, the authors postulated that the early changes reflected bone bruises or microfractures due to the injury itself. However, to our knowledge, there have been no published reports of transient lesions of the femoral head after traumatic hip dislocation showing no abnormalities just after dislocation. We herein report a rare case in which MRI showed a transient epiphyseal lesion four months after traumatic hip dislocation.

2. Presentation of case

A 22-year-old Japanese man dislocated his right hip joint posteriorly while snowboarding and was treated with closed reduction within three hours. He had no history of hip disease or no other risk factors contributing to bone fragility, including steroids, alcohol abuse, smoking, renal and liver transplant or systemic lupus erythematosus. On MRI performed two days after dislocation, no

[☆] Investigation performed at the Department of Orthopaedic Surgery, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan.

* Corresponding author.

E-mail address: yamataku@ortho.med.kyushu-u.ac.jp (T. Yamamoto).




Fig. 1. Short-tau inversion recovery (STIR) sequence image 2 days after the traumatic dislocation of the hip joint. Coronal STIR sequence image obtained 2 days after dislocation shows no bony injuries or bone marrow edema around the hip joint.




Fig. 2. Anteroposterior radiograph after the onset of right hip pain. An anteroposterior radiograph obtained after the onset of right hip pain shows an irregular contour at the outside of the femoral head, around which focal bone loss is observed (white arrows).

bony injuries or bone marrow edema were observed around the hip joint (Fig. 1). For one week after the injury, the patient was encouraged to avoid weight-bearing activities. Two weeks later, he was discharged from the hospital. He subsequently stayed home for two months without leaving his home, using crutches while indoors. Three months after the injury, he returned to work at a laundry, where he was required to lift heavy laundry loads and stand all day.

Four months after the injury, the patient suddenly experienced right hip pain without any new trauma or injury. On examination, the range of motion of the right hip was slightly limited (110° of flexion, 0° of extension, 40° of external rotation and 20° of internal rotation). An anteroposterior radiograph obtained four months after the injury showed an irregular contour along the outside of the femoral head, around which focal bone loss was observed (Fig. 2). A low-intensity band convex to the articular surface was apparent above the epiphyseal scar on T1-weighted imaging (Fig. 3a), and bone marrow edema changes were observed around the band lesion on short-tau inversion recovery (STIR) imaging (Fig. 3b). Moreover, gadolinium-enhanced MRI showed the band and surrounding area to be enhanced, accompanied by diffuse bone marrow edema (Fig. 3c), and single-photon emission computed tomography/computed tomography (SPECT/CT) with Tc-99 m hydroxymethylene diphosphonate demonstrated the presence of increased uptake involving the entire femoral head (Fig. 4). Based on these findings, the patient was suspected to have an insufficiency fracture of the femoral head. After a two-month period of non-surgical conservative therapy, including the avoidance of weight-bearing activities, the patient's hip pain resolved and the low-intensity band previously seen on T1-weighted imaging was no longer observed on follow-up MRI performed two months after the onset of pain (Fig. 5).

3. Discussion

To our knowledge, only one previous study has shown the occurrence of transient bone marrow signal changes on MRI after hip dislocation. Poggi et al. performed MRI examinations of 14 hips following traumatic dislocation and found transient bone marrow changes in the femoral head in five cases (36%) [8]. However, since




Fig. 3. a–c Magnetic resonance images (MRI) performed 4 months after the injury following the onset of right hip pain. (a) A coronal T1-weighted image shows a low-intensity band convex to the articular surface above the epiphyseal scar (black arrows). (b) A coronal short-tau inversion recovery (STIR) sequence image shows bone marrow edema around the band (white arrowheads). (c) On gadolinium-enhanced magnetic resonance image MRI, the band and surrounding area were enhanced (white arrows), accompanied by diffuse bone marrow edema (white arrowheads).

Download English Version:

<https://daneshyari.com/en/article/4288240>

Download Persian Version:

<https://daneshyari.com/article/4288240>

[Daneshyari.com](https://daneshyari.com)