



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Pneumothorax following shoulder arthroscopy under combined regional and general anaesthesia—A case report



Oskar Leander-Olsson, Anna Borglund-Hemph, Jan G. Jakobsson*

Department of Anaesthesia & Intensive Care, Institution for Clinical Science, Karolinska Institutet, Danderyds University Hospital, Stockholm, Sweden

ARTICLE INFO

Article history:

Received 25 April 2016

Received in revised form 5 May 2016

Accepted 6 May 2016

Available online 11 May 2016

Keywords:

Case report

Upper extremity block

Shoulder arthroscopy

General anaesthesia

Pneumothorax

ABSTRACT

INTRODUCTION: Pneumothorax is a most rare intraoperative event but jeopardise safety. Risk factors for its occurrence should be acknowledged and it should be considered in cases of unexpected deterioration of oxygenation that not respond to standard management to improve oxygenation.

PRESENTATION OF CASE: We describe a patient with mild obesity and recent shoulder trauma scheduled for elective arthroscopy under combined interscalene block and general anaesthesia in the beach chair position. The block was performed with ultrasound guidance in accordance with the departmental routines and before the induction of general anaesthesia. The patient experienced low saturation shortly after administration of general anaesthesia. A thoracic ultrasound was performed in order to rule out pneumothorax and recruitment manoeuvres was done based on the suspicion of atelectasis. Surgery was commenced under pressure controlled ventilation and saturation was kept around 90 at FiO₂ 1.0 after recruitment manoeuvres. An initial postoperative lung X-ray on the operating table showed subcutaneous emphysema but no pneumothorax. A postoperative CT-thorax, patient still intubated, showed an extensive pneumothorax that was treated with pulmonary drainage. Patient had subsequently an uncomplicated postoperative course.

DISCUSSION: We can not state “whom to blame”, the interscalene block, intubation and pressures assisted ventilation, recruitment manoeuvres, or surgery? In this paper we discuss possible causes that should raise the suspicion around pneumothorax in patients exhibiting unexpected low saturation perioperatively.

CONCLUSION: Compromised oxygenation during surgery that not respond to recruitment should raise suspicion about pneumothorax.

© 2016 The Author(s). Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Pneumothorax, air entrainment into the pleura space compromise oxygenation and may also impact venous return and thus cause circulatory instability. The degree of impaired oxygenation may vary. Pneumothorax should however be considered in patients showing unexpected deterioration in oxygenation and risk factors for its occurrence should be acknowledged. Lung auscultation and frontal X-ray not sufficient to exclude a pneumothorax. We describe a patient with pneumothorax shown on postoperative CT-scan of the thorax following shoulder arthroscopic rotator cuff repair performed under combined interscalene plexus block and general anaesthesia in a patient with multiple risk factors.

2. Case report

A 72 year old man with BMI 38.4, sleep apnoea and night time CPAP, gout and benign prostatic hyperplasia scheduled for surgery on his left shoulder because of rotator cuff syndrome. He had previously been operated for spinal stenosis and had also had arthroscopic surgery in both shoulders twelve years ago. He was assessed as being vital and healthy, a previous echocardiogram from January 2015 had shown normal heart function but with a minor aortic insufficiency. During the waiting time for surgery he fell in a stairway and hit his right shoulder giving him a fracture in fossa glenoidale and partial ruptures in subscapularis and supraspinatus. These injuries lead to frequent luxations of the shoulder which the patient painfully could relocate himself. Because of the trauma and more intense symptoms from the right shoulder he planned surgery on his left shoulder got exchanged for suturing the rotator cuff on his right shoulder instead. Upon arrival to the surgery department the patient was in a good mood. The patient was circulatory stabile. Respiratory wise he had a saturation of 92% measured by standard pulseoximetry, earlier in the

* Corresponding author.

E-mail address: Jan.jakobsson@ki.se (J.G. Jakobsson).

morning at the ward it was at 93%. The low saturation was thought of to be due to atelectasis from the overweight. He was assessed as having a difficult airway. The major part of his overweight was on his upper-body giving him a wide and very short neck and a small chin. The ultrasound probe for the plexus block (interscalene) gave a good view of the nerves allowing for needle placement in-plane. The most difficult part of the block was the patient's skin being hard to puncture with the ultrasound needle and thus a pre-puncture had to be made with a larger, sharper needle. The needle was easily spotted and the local anaesthetic was administered with good effect. After the block was administered the patient was preoxygenated with low levels of pressure support increasing the saturation to 100%. The patient received general anaesthesia with rapid sequence induction and intubation with a video laryngoscope (STORZ) without any desaturation.

Initially there was no problem with ventilation, the patient was on pressure control with volume guarantee and we were able to reduce to minimal flow on the ventilator with good saturation. Shortly after the start of surgery the nurse anaesthetist called to notify that the patient saturation had dropped to 91% and that the ventilator needed high top pressures to get sufficient tidal volumes. It was assessed that the patient had a shallow anaesthetic depth and was not breathing in synchrony with the ventilator. He was switching over to manual ventilation and adequate tidal volumes was easily restored. Manual lung requirement was performer and the end-tidal sevoflurane increased. He was put back on pressure controlled ventilation with volume guarantee and ventilation was stabilised. After some further time oxygen saturation once more started to decrease and FiO₂ was increased. Lung auscultation revealed breath sounds bilaterally but better on patients left. The tracheal tube which was secured relatively deep, on 26 cm, was adjusted to 23 cm resulting in improved breath sounds but with no major improvement in saturation. A new lung recruitment manoeuvre, this time with the ventilator (pressure control, breathing frequency put on 5/min, and PEEP gradually raised to a top pressure of 30 cmH₂O) was performed. This improved compliance from 30 to 65 ml/cm H₂O and again improved saturation but just briefly. The patient now had a saturation around 90% and FiO₂ 1.0 and no clear breath sound were heard on right side. Surgery was completed within 30 min without further deterioration of patient's respiratory status. Thoracic ultrasound and then chest X-ray were performed immediately post-surgery, patient still on operating table and intubated on the suspicion of a pneumothorax. The emergent ultra-sound was inconclusive and X-ray showed extensive subcutaneous emphysema but did not confirm any clear pneumothorax. The patient could not be taken off the ventilator due to the high oxygen demand and was transported still anaesthetised and intubated for a CT-scan. The thoracic CT confirmed a large pneumothorax. Patient was transferred to the ICU and a traditional chest-drain was inserted. Oxygenation rapidly improved and the patient could after a couple of hours be extubated and had a further uncomplicated course. The drain was removed after 2 days.

3. Discussion

Our patient had an unexpected intraoperative deterioration in oxygenation that did not respond adequately to lung recruitment. The suspicion of pneumothorax was raised but following adjustment of the tube position and intraoperative lung auscultation it was initially rejected. Frontal ultra-sound as well as an acute frontal X-ray failed to confirm the diagnosis that was later confirmed on CT.

Intraoperative pneumothorax is a rare event but should be acknowledged in situation of unexpected decrease in oxygenation that did not respond to lung recruitment. We can not state "whom to blame"; there are several plausible explanations for the devel-

opment of pneumothorax in the present case. The most prevalent factors include: needle injuries associated with the performance of the upper extremity block, external pressure on the upper pleura dome associated with the shoulder arthroscopic procedure, thermal forces induced by the use of diathermia, baro-trauma caused by airway pressure associated to intubation and ventilation, or lung recruitment. Patient factors include: smoking habits, lung disease (COPD, emphysema, cancer), prior history of pneumothorax, history of recent thorax trauma and rare diseases such as alpha-1-antitrypsin deficiency should also be taken into account.

4. Upper extremity block and the risk of pneumothorax

Pneumothorax is one of side effects that may be seen, albeit rarely, associated with interscalene block. The use of ultrasound help visualising the needle tip reducing but not eliminating the risk [1,2]. Mandim et al. suggested that a high pleura dome e.g. caused by smoking may contribute to needle injury. Montoro et al. suggest that merely short loss of needle tip visualisation could be a possible explanation [3]. There are several papers reporting a reassuringly low incidence associated to ultra sound guided blocks especially when performed of trained anaesthesiologists [4,5]. Bhatia et al. summarised the 3 major factors that increase risk of pneumothorax associated to ultra-sounded guided upper extremity block; **operator factors**, loss of visualisation of the needle tip, **technical factors**, probe and plane and hydro-visualisation, and **patient factors such as** lung disease, smoking, emphysema and obesity [6]. Abell and Barrington published a case report in 2014 data collected from the International Registry of Regional Anesthesia (IRORA), a multi-centre registry that systematically and uniformly collects information from patients who receive peripheral nerve blockade [7,8]. They estimated the risk of pneumothorax associated with ultrasound-guided supraclavicular block to be 0.4 per 1000 but with a huge 95% confidence interval, 0.01–2.3 per 1000 blocks. They also addressed needle visualisation in terms of the relative importance in reducing the risk of pneumothorax. Neal recently published an updated systematic review around the topic of ultrasound guidance for patient safety. The review showed ultrasound guidance to have a favourable effect in reducing the risk of local anaesthesia toxicity, as well as reducing but not eliminating the risk of pneumothorax [9].

5. Subcutaneous emphysema, pneumomediastinum and pneumothorax related to shoulder arthroscopy, the surgical procedure

There are rare reports of pulmonary complications in arthroscopy of the shoulder where the complication has been suggested to possibly be related to the arthroscopic procedure itself. Lee et al. described two cases of subcutaneous emphysema and tension pneumothorax [10]. Lau went on to described a patient that developed subcutaneous emphysema and pneumo-mediastinum associated with arthroscopic surgery of the shoulder and raised the suspicion that it may have had an association to the intraarticular shaving [11]. Dietzel and Cuillo described 4 cases of pneumothorax following a shoulder arthroscopy under general anaesthesia [12]. All four patients had risk factors but a relation to the surgical procedure could not be excluded. Calvisi et al. described a case of pneumothorax associated with arthroscopy performed under local regional anaesthesia [13]. They emphasise the importance of adequate intended equipment such as pumps and suction devices in order to minimising tissue air entrapment. Pneumothorax has also been seen associated to breast reconstruction [14]. Possibly related to pleura trauma associated to stretch forces, diathermia or accidental needle injury during placement of the local anaesthesia [15].

Download English Version:

<https://daneshyari.com/en/article/4288248>

Download Persian Version:

<https://daneshyari.com/article/4288248>

[Daneshyari.com](https://daneshyari.com)