



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

An intra-cerebral abscess in a patient with Eisenmenger syndrome: An unusual case

Allan Hall^{a,*}, Mark A.J. White^b, Pasquale Gallo^b^a Institute of Neurological Sciences, Queen Elizabeth University Hospital, Glasgow, United Kingdom^b Department of Clinical Neurosciences, Western General Hospital, Edinburgh, United Kingdom

ARTICLE INFO

Article history:

Received 25 November 2015

Received in revised form 17 January 2016

Accepted 18 January 2016

Available online 25 January 2016

Keywords:

Intracerebral abscess

Eisenmenger syndrome

ABSTRACT

INTRODUCTION: We present an unusual case where a large intra-cerebral abscess with severe complications was treated successfully with medical management alone in a patient with Eisenmenger syndrome. **PRESENTATION OF CASE:** A 40 year old patient with Eisenmenger syndrome presented with a seven day history of headache, neck pain and intermittent vomiting and fever. The only neurological examination finding was blurred vision. MRI revealed a large left occipital lobe abscess, which subsequently ruptured into the lateral ventricle with associated ventriculitis and hydrocephalus. This complicated abscess was successfully treated with intravenous antibiotics alone, with complete resolution of the abscess and hydrocephalus on MRI at 13 months post-diagnosis.

DISCUSSION: Patients with congenital cyanotic heart conditions, like Eisenmenger syndrome, are at increased risk of developing intra-cerebral abscesses. Effectively managing large abscesses with associated intra-ventricular rupture and hydrocephalus in these patients without any form of surgical intervention, including aspiration, is extremely rare.

CONCLUSION: Patients with congenital cyanotic heart disease who develop large intra-cerebral abscesses with severe complications, which would normally warrant surgical intervention, have the potential to be successfully treated with antibiotics alone with excellent outcome.

© 2016 The Authors. Published by Elsevier Ltd. on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Patients with congenital cyanotic heart disease, like Eisenmenger syndrome, are at an increased risk of developing intra-cerebral abscesses. It is common practice for abscesses larger than 2 cm to undergo some form of surgical intervention, ranging from aspiration to excision of the abscess. However it is rare for large abscesses with associated intra-ventricular rupture and hydrocephalus to be successfully treated with intravenous antibiotics alone.

We present a rare case of a 40 year old patient with Eisenmenger syndrome who developed a 4 cm intra-cerebral abscess with rupture into the lateral ventricle and associated hydrocephalus. The patient was successfully treated with IV antibiotics alone with excellent outcome.

2. Presentation of case

A 40-year-old female with Eisenmenger syndrome, secondary erythrocytosis, and tetralogy of fallot was admitted with a seven day history of headache, neck pain and intermittent vomiting and fever. There was also blurred vision however no visual field defects were detected upon examination. There was no history of seizures or loss of consciousness, and there was no upper or lower limb weakness either in the history or at presentation.

On examination the patient had both central and peripheral cyanosis with a loud P2 ejection systolic murmur at the aortic area and a pansystolic murmur at the tricuspid area. The patient had no speech deficit and her neurological examination, including cranial nerve examination, showed no deficits. Her observations at presentation were GCS 15; blood pressure 120/70; heart rate of 74 and oxygen saturation of 77% (the patient's normal saturations are 70% due to her Eisenmenger syndrome).

The patient's medication at presentation was ferrous sulphate 200 mg OD (once daily), warfarin, amiodarone 200 mg OD, furosemide 20 mg OD, paroxetine 40 mg OD, folic acid 400 micrograms OD, sildenafil 20 mg TDS (three times daily), and co-codamol 30/500 QDS (four times daily).

The patient subsequently underwent a CT head with contrast, which showed an intra-cerebral abscess within the left occipital

* Corresponding author at: Institute of Neurological Sciences, Queen Elizabeth University Hospital, Glasgow G51 4TF, United Kingdom.

E-mail address: allanhall@nhs.net (A. Hall).




Fig. 1. (a) MRI with contrast on day of admission which shows abscess within the left occipital lobe. (b) FLAIR MRI on day 33 post-admission, which shows reduction in abscess size however there is now diffuse enhancement of the lateral ventricular ependyma. (c) MRI with contrast at 13 months post-admission. Note there is complete resolution of the abscess.

lobe abutting the occipital horn of the lateral ventricle. The abscess measured 40×40 mm in the axial plane. A baseline MRI was undertaken on the day of admission, which confirmed the lesion was an abscess (Fig. 1a). The patient was started on ceftriaxone and metronidazole after initial discussion with microbiology. However the patient's clinical state deteriorated 5 days post admission and a contrast-enhanced MRI was undertaken, which showed rupture of the abscess into the lateral ventricle. It was concluded the abscess had ruptured into the lateral ventricle by the combination of deterioration in the patient's clinical status and a diffusely enhanced ventricular ependyma with hydrocephalus on the MRI. Following a cardiology and anaesthetic review the patient was deemed unfit for surgery due to her Eisenmenger syndrome and therefore a biopsy of the abscess, surgical evacuation or ventricular drainage was unable to be performed under general anaesthetic.

Local anaesthetic aspiration or ventricular access device could have been considered as a life saving measure with its own inherent difficulties, including patient positioning. After discussion with microbiology the Ceftriaxone was changed to meropenam, and vancomycin was added. The

IV course of meropenam, metronidazole and vancomycin continued for 4 weeks. The patient continued to have intermittent episodes of pyrexia and therefore vancomycin was subsequently switched to linezolid. In addition meropenam was switched to co-amoxiclav. The patient completed 10 weeks of antibiotics as an in-patient and was discharged on co-amoxiclav, linezolid and metronidazole, which she continued for a further 2 weeks. In addition to the day of admission MRI's were undertaken at 18, 24, 33 and 61 days post-admission. The MRIs, collectively, showed an overall trend of a gradual reduction in the abscess size with an initial enlargement of ventricular involvement (Fig. 1b), followed by a subsequent reduction.

The patient was kept under regular day ward review following discharge. She reported no further episodes of pyrexia, and her follow up MRI at 6 months post-admission showed continued resolution of the abscess and normal ventricular size. The latest follow-up MRI revealed complete resolution of the abscess at 13 months post diagnosis (Fig. 1c).

This work has been reported in line with the CARE criteria (<http://www.care-statement.org/>).

Download English Version:

<https://daneshyari.com/en/article/4288575>

Download Persian Version:

<https://daneshyari.com/article/4288575>

[Daneshyari.com](https://daneshyari.com)