



Contents lists available at ScienceDirect

## International Journal of Surgery Case Reports

journal homepage: [www.casereports.com](http://www.casereports.com)

# Favorable outcome of a total hip arthroplasty with insufficient bone coverage of the roof reinforcement ring



Fernando M. Judas\*, Francisco M. Lucas, Ruben L. Fonseca

Orthopedics Department, Coimbra Hospital and University Centre (CHUC), Faculty of Medicine, University of Coimbra, Praceta Mota Pinto, 3030 Coimbra, Portugal

## ARTICLE INFO

### Article history:

Received 5 March 2015

Received in revised form

16 September 2015

Accepted 17 September 2015

Available online 30 September 2015

### Keywords:

Developmental dysplasia of the hip

Total hip arthroplasty

Roof reinforcement ring

Conical stem

## ABSTRACT

**INTRODUCTION:** Total hip arthroplasty in patients with developmental dysplasia of the hip can be a complex procedure due to acetabular and proximal femoral deformities.

**PRESENTATION OF CASE:** A 59-year-old male patient underwent a total hip arthroplasty for the treatment of end-stage dysplastic osteoarthritis. A roof reinforcement ring, a cemented polyethylene cup, and a cementless stem were used. A portion of the superior rim of the ring was uncovered by the host bone. Morsellized autogenous femoral-head graft was impacted to fill the space between the superior rim of the ring and the superior part of the dysplastic acetabulum. At the follow-up after 5-years, the patient had no complaints and was very satisfied with the operation result. The hip radiograph revealed no signs of instability of the acetabular component, and no bone graft resorption.

**DISCUSSION:** Favorable results were described using metal rings and conical femoral stems for the treatment of the developmental dysplasia of the hip. The superior rim of the metal ring should be against host bone for 60% of its support. Despite the suboptimal implantation of the ring compromising, apparently, mechanical stability of the arthroplasty, the outcome was favorable.

**CONCLUSION:** This result can be supported by the good fixation of the metal ring to the pelvis with screws, the adequate orientation of both components of the total hip arthroplasty, and the bone graft incorporation.

© 2015 The Authors. Published by Elsevier Ltd. on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The outcomes of maltreated or neglected developmental dysplasia of the hip are highly detrimental. Degree of subluxation and time left untreated are the two main factors that contribute to severity of the sequelae of this disorder [1].

Total hip arthroplasty (THA) is the procedure of choice for most patients with symptomatic end-stage osteoarthritis secondary to hip dysplasia [2]. Ideally, the acetabular cup is covered completely by host bone and should span the distance between the teardrop and the superolateral margin of the acetabulum [3].

Standard cementless femoral stems with a rectangular cross-section can be difficult to use for the treatment of dysplastic osteoarthritis, due to the femoral proximal deformities. A cementless stem with a conical shape can be indicated [4].

In this report a THA was performed for the treatment of a hip osteoarthritis secondary to developmental dysplasia. An acetabular roof reinforcement ring with a cemented polyethylene cup, and a

conical stem were implanted. A superolateral portion of the metal ring was uncovered by host-bone, due to an inadequate operative technique.

The purpose of this paper was to show the outcome of the THA at 5-years follow-up, namely the mechanical behavior of the acetabular component.

## 2. Case report

A 59-year-old male patient underwent a total hip arthroplasty for the treatment of end-stage dysplastic osteoarthritis, in 2010. He presented a severe hip pain, functional impairment, the limb-length discrepancy was 30 mm, and was Thendelenburg-type gait. The pelvic radiograph revealed dysplasia of the right acetabulum according Hartofilakidis classification [5], coxa valga, and pelvic obliquity (Fig. 1). The right femoral head presented marginal osteophytes and was large and flattened in shape. Using a standard posterior approach, a roof reinforcement ring and a cemented polyethylene cup were implanted. The femur was reconstructed with a cementless femoral stem with conical shape.


An appropriate size ring was inserted with a slight press-fit into the acetabulum, and secured using cancellous bone screws in order to achieve satisfactory mechanical fixation to the pelvis.

\* Corresponding author.

E-mail addresses: [fernandojudas@gmail.com](mailto:fernandojudas@gmail.com), [fernandojudas@iol.pt](mailto:fernandojudas@iol.pt) (F.M. Judas), [francisco.m.lucas@gmail.com](mailto:francisco.m.lucas@gmail.com) (F.M. Lucas), [ruben.fonseca@msn.com](mailto:ruben.fonseca@msn.com) (R.L. Fonseca).



**Fig. 1.** Preoperative anteroposterior radiograph of the pelvis showing advanced right hip osteoarthritis, dysplasia of the acetabulum and coxa valga, and pelvic obliquity.



**Fig. 2.** Postoperative THA radiograph at follow-up period of 1-year. The superolateral portion of the roof reinforcement ring is not covered by the native acetabulum. It is possible to observe the incorporation of impacted morsellized autograft placed between the rim of the metal ring and the superior part of the dysplastic acetabulum.

A polyethylene cup was then secured with cement in the desired position. Thus, optimum anteversion and inclination is possible, regardless of the position of the ring.

However, a portion of the rim of the metal ring was uncovered by host bone. Then, the removed femoral head was morsellized (using a rongeur) on table, and bone graft was carefully impacted to fill the space between the uncovered border of the ring, and the superolateral margin of the native acetabulum. A conical stem was easily implanted with a correct femoral anteversion.

The postoperative course showed no problems with respect to the hip. Daily activity was quickly resumed. At 6-months after surgery the patient presented an asymptomatic hip with no limping, and approximation of equalization of limb-length. At 1-years

follow-up the hip radiograph revealed a satisfactory orientation of the prosthesis, a small medialization of hip's center of rotation, and some heterotopic ossifications. The bone autograft was incorporated (Fig. 2).

At 5-years follow-up, the patient continued to present a stable and asymptomatic hip and expressed high degree of satisfaction with surgery result. The hip radiograph showed unchanged and stable seating of the acetabular component, and stable fixation of the femoral stem by bone ingrowth, no measurable subsidence or radiolucent lines around the stem (Fig. 3). Heterotopic ossifications can be observed and were classified as stage I according to the classification of Brooker [6]. The autograft presented no signs of bone resorption.

Download English Version:

<https://daneshyari.com/en/article/4288644>

Download Persian Version:

<https://daneshyari.com/article/4288644>

[Daneshyari.com](https://daneshyari.com)