



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Unusual case of life threatening subcutaneous hemorrhage in a blunt trauma patient

Ashraf F. Hefny^{a,*}, Laith N. Kaka^b, El Nazeer A. Salim^a, Nabil N. Al Khoury^c^a Department of Surgery, Al Rahba Hospital, Abu Dhabi, United Arab Emirates^b Department of Neurosurgery, Al Rahba Hospital, Abu Dhabi, United Arab Emirates^c Department of Anesthesia and Critical Care, Al Rahba Hospital, Abu Dhabi, United Arab Emirates

ARTICLE INFO

Article history:

Received 13 April 2015

Received in revised form 23 August 2015

Accepted 23 August 2015

Available online 29 August 2015

Keywords:

Injury

Shock

Subcutaneous

ABSTRACT

INTRODUCTION: Hemorrhage is the most common cause of shock in injured patients. Bleeding into the subcutaneous plane is underestimated cause of hypovolemic shock.**PRESENTATION OF CASE:** Unrestrained male driver involved in a rollover car crash. On examination, his pulse rate was 144 bpm, blood pressure 80/30 mmHg, and GCS was 7/15. His right pupil was dilated but reactive. Back examination revealed severe contusion with friction burns and lacerations. A Focused Assessment Sonography for Trauma (FAST) was performed. No free intraperitoneal fluid was detected. CT scan of the brain has shown right temporo-parietal subdural hematoma and extensive hematoma in the deep subcutaneous soft tissues of the back. Decompressive craniotomy and evacuation of the subdural hematoma was performed. On the 4th postoperative day, three liters of dark brown altered blood was drained from the subcutaneous plane.**DISCUSSION:** The patient developed severe hypovolemic shock and our aim was to identify and control the source of bleeding during the resuscitation. The source of bleeding was not obvious. Severe shearing force in blunt trauma causes separation between the loose subcutaneous tissues and the underlying relatively immobile deep fascia. This is known as post-traumatic closed degloving injury. To our knowledge this is the first reported case in the English Literature with severe subcutaneous hemorrhage in blunt trauma patients without any previous medical disease.**CONCLUSION:** Bleeding into the subcutaneous plane in closed degloving injury can cause severe hypovolemic shock. It is important for the clinicians managing trauma patients to be aware this serious injury.© 2015 The Authors. Published by Elsevier Ltd. on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Road Traffic Collisions (RTC) is major health and economic burden all over the world. The main victims of RTC are the young productive population. The patients usually suffer multiple injuries that differ according to the biomechanics of the collision [1]. Care of the patient with shock can be one of the most challenging issues in trauma patients. Hemorrhage is the most common cause of shock in injured patients [2]. Bleeding into the subcutaneous plane is underestimated cause of hypovolemic shock [3]. Herein we are report a case of young man who was involved in RTC and developed severe hypovolemic shock. It was due to closed degloving injury of his back skin with extensive subcutaneous hemorrhage.

2. Case report

A 19-year-old unrestrained male driver was involved in a rollover car crash. He was ejected around 30 m away from the car. In the Emergency Department, his pulse rate was 144 bpm, blood pressure 110/70 mmHg, and respiratory rate was 44/min. His GCS was 7/15 and the right pupil was dilated but reactive to light (4 mm in diameter). Chest examination revealed decreased air entry on the right side. Abdomen was soft and pelvis was clinically stable. Back examination revealed severe contusion mainly on the right side of the back of the chest and left flank with friction burns and lacerations (Fig. 1). No past history of chronic medical diseases.

The patient was immediately intubated. Chest X-ray has shown right sided hemopneumothorax and a chest tube was inserted. Around 200 ml of blood was drained.

Suddenly, the patient becomes hypotensive with blood pressure of 80/30 mmHg. A Focused Assessment Sonography for Trauma (FAST) was performed. No free intraperitoneal fluid was detected. The IVC was flat and the heart was hyper contractile. No pericardial effusion was detected. Patient has neither pelvic nor long

* Corresponding author at: Department of Surgery, Al Rahba Hospital, Abu Dhabi Clinical Assistant Professor, Trauma Group, Department of Surgery, CMHS, Al-Ain P.O. Box 18532 Main Building of Post Office, Al-Ain, UAE, Fax: +971 3 7672067.

E-mail addresses: ahefny@uaeu.ac.ae, hefnyashraf@hotmail.com (A.F. Hefny).



Fig. 1. Severe contusion and friction burn over the back of the patient.




Fig. 2. CT scan of the brain shows right temporo-parietal subdural hematoma (arrow).

bone fracture. The patient received three units of packed red blood cells (RBCs) and four liters of crystalloids during the early phase of resuscitation.

A decision was taken to transfer the patient for rapid CT scan of the brain and cervical spine on the way to operating theater for craniotomy and possible diagnostic laparoscopy. CT scan of the brain has shown right temporo-parietal subdural hematoma (Fig. 2).

Right fronto-temporo-parietal decompressive craniotomy and evacuation of the subdural hematoma was performed. The patient received further four units of packed RBCs during the operation. On table, a repeated abdominal ultrasound has shown minimal free fluid in Morrison pouch and pelvis. A diagnostic laparoscopy has shown a right side retroperitoneal hematoma with minimal hemoperitoneum. Increased swelling in the right flank was noticed at the end of the operation. Trauma CT scan has shown stable fracture of the 1st, 3rd and 5th thoracic vertebrae, fracture of the right transvers process of all lumbar vertebrae, right sided retroperitoneal hematoma, and extensive hematoma in the deep subcutaneous soft tissues of the back (Fig. 3). No contrast extravasation



Fig. 3. Sagittal reformatted CT scan of the abdomen shows extensive hematoma between the subcutaneous fat and the paraspinal muscles (arrow).

was detected. In the Intensive Care Unit (ICU) the patient received three units of packed RBC and he remains hemodynamically stable.

On the 4th postoperative day, a huge soft fluctuant swelling all over the back was noticed (Video 1). Ultrasound scan has shown cystic swelling involving the whole back with depth of about 7 cm (Fig. 4). Percutaneous suction drainage was performed. Three liters of dark brown altered blood was drained from the subcutaneous plane. Replacement of the skull bone flap with dural graft reconstruction was performed on the 13th post-operative day. Post-operative period was uneventful and the subcutaneous swelling resolved completely.

3. Discussion

It is important in managing injured patient to recognize the presence of the shock state and to initiate the treatment during identification of the possible cause. The initial evaluation and treatment of our patient followed the Advanced Trauma Life Support Program guidelines [2].

The patient developed hypovolemic shock and our aim was to identify and control the source of bleeding during the resuscitation. Even after the decompressive craniotomy and diagnostic laparoscopy, the source of bleeding was not obvious. Trauma CT scan with IV contrast was performed to identify the source of bleeding and possible angio-embolization if needed.

Post traumatic extensive hematoma in the deep subcutaneous soft tissues is a rare condition [4].

Download English Version:

<https://daneshyari.com/en/article/4288831>

Download Persian Version:

<https://daneshyari.com/article/4288831>

[Daneshyari.com](https://daneshyari.com)