

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Case report of nail bed injury after blunt trauma; what lies beneath the nail?



Geraldine L. Nanninga*, Kevin de Leur, Anne Loes van den Boom, Mark R. de Vries, Tessa M. van Ginhoven

Department of Surgery, Reinier de Graaf Groep, Delft, The Netherlands

ARTICLE INFO

Article history:

Received 28 June 2015

Received in revised form 25 August 2015

Accepted 25 August 2015

Available online 30 August 2015

Keywords:

Blunt trauma

Nail bed

Distal Phalanx

Fingertip injuries

ABSTRACT

INTRODUCTION: A stable, mobile and sensate fingertip is of paramount importance to perform daily tasks and sense dangerous situations. Unfortunately, fingertips are easily injured with various extents of soft tissue damage. Delayed and inadequate treatment of nail bed injuries may cause substantial clinical problems. The aim is to increase awareness about nail bed injuries among physicians who often treat these patients.

PRESENTATION OF CASE: We present a 26-year-old male with blunt trauma to a distal phalanx. Conventional radiographs showed an intra-articular, multi-fragmentary fracture of the distal phalanx. At the outpatient department the nail was removed and revealed a lacerated nail bed, more than was anticipated upon during the first encounter at the emergency department.

DISCUSSION: Blunt trauma to the fingertip occurs frequently and nail bed injuries are easy to underestimate. An adequate emergency treatment of nail bed injuries is needed to prevent secondary deformities and thereby reduce the risk of secondary reconstruction of the nail bed, which often gives unpredictable results.

CONCLUSION: However, adequate initial assessment and treatment are important to achieve the functional and cosmetic outcomes. Therefore awareness of physicians at the emergency department is essential.

© 2015 The Authors. Published by Elsevier Ltd. on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Fingertips are an essential part of the human body and are used to explore our surroundings and provide danger warning signs. In addition, stable, mobile and sensate fingertips are needed to perform activities of daily living. With the recent increase in the use of touchscreen in smartphones, tablets and remote controls, the fingertip is needed more than ever for those delicate movements. In addition the fingertip is important for the overall function of the hand and cosmetic impairment can have an enormous social impact [1,2].

Fingertip and nail bed injuries are seen at all ages, with a peak incidence in 4–30 years old patients. Patients are most often initially treated by family doctors or physicians at the emergency department [1,3,4]. Chang et al. reported that 10% of all the accidents encountered in the emergency department involve the hand. In case of fingertip injuries the nail bed is damaged in 15–24% of the cases [3]. Fingertip injuries are defined as any injury to the distal

phalanx, including soft tissue, nail plate, nail bed, tendon and bone [1,4]. In most cases a combination of injured structures is present.

Fingertip injuries are caused by sharp or blunt trauma. Sharp trauma (amputations), for example with knives or cutting machines, is less common and the extent of the injury, especially involvement of the nail bed, is relatively easy to identify. In contrast, more often, injuries are due to blunt trauma (e.g. crush injuries by doors, heavy machinery or during sports) [5]. This type of injury results in compression of soft tissue, mainly the nail bed, between the nail and the underlying bone in addition to the external forces. The extent of soft tissue damage can easily be underestimated and often the nail bed is injured.

Nail bed injuries may vary from subungual hematomas, lacerations and avulsions of matrix tissue. Currently there is an ongoing debate about whether or not to remove the nail and inspect the nail bed after blunt trauma. Inspection of the nail bed is important in order to appreciate the extent of the injury. However, not every needs to be removed. If lacerated, it is important that the nail bed will be repaired adequately and with attention to detail as inadequately treated nail bed injuries lead to loss of function and pain, which have an enormous impact on the patient [1,6]. It is recommended that in repairing the nail bed a loupe magnification and the use of micro (vascular) instruments are used. Sometimes early

* Corresponding author at: Department of Surgery, Reinier de Graaf Groep, P.O. Box 5011, 2600 GA Delft, The Netherlands.

E-mail address: geraldinenanninga@gmail.com (G.L. Nanninga).

intervention by a dedicated hand surgeon is necessary to achieve better functional and aesthetic outcomes [1].

The aim of this report is illustrate the clinical problem of nail bed injuries by means of a case description. In addition, we will provide guidelines for physicians encountering blunt fingertip injuries highlighting the importance of early identification and treatment of nail bed injuries.

2. Case report

A 26-year-old healthy male presented himself at the Emergency Department (ED) with an injured fingertip after blunt trauma sustained during a field hockey game. Inspection of the fourth digit of the right hand revealed isolated fingertip injury. Both proximal and middle phalanx were unaffected. A laceration of two centimeters was identified at the volar/ulnar side of the fingertip. The nail was partially loose, and the nail bed seemed more or the less intact. There was no loss of sensation and all movements in the distal interphalangeal joint were possible, but limited due to pain (Fig. 1A). Plain radiographs showed an intra-articular, multi-fragmentary fracture of the distal phalanx with a congruent distal interphalangeal joint (Fig. 1B, C). Further examination under local “oberst” anesthesia confirmed no problems with flexion and extension of both the proximal and distal interphalangeal joints. As a first treatment, the laceration was sutured with transcutaneous interrupted stiches after which the fingertip regained a normal aspect. Antibiotics were prescribed, because of the extended nature of the fracture and soft tissue damage. The patient was scheduled for an appointment at the outpatient department the next day. This physician identified the injury as a “high risk injury” for nail bed involvement based on the laceration and the fracture. The patient was directly scheduled for a nail bed repair in the operation room the same day. After removing the nail, the extent of the nail bed injury became apparent and was much greater than

anticipated by the physician on the by the physician on the ER department the day before. Multiple lacerations were identified and the nail bed was repaired with interrupted sutures. Prolene 6.0 was used as suture material (Fig. 1D). The nail was replaced and fixated by proximal and distal sutures with Ethilon 5.0 (Fig. 1E) to function as a splint and natural cover for 2 weeks. Postoperative radiographs showed a nearly anatomical reposition of the intra-articular multi-fragmentary fracture of the distal phalanx. After 2 weeks the nail and the sutures were removed. Thereafter, the fingertip was protected by a small splint. During follow-up no post-operative complications were encountered. The nail bed recovered completely and the nail plate seemed to develop without any deformities (Fig. 1F). Seven months after the initial trauma there are no complaints about pain or loss of function during daily activities.

3. Discussion

Patients with blunt trauma of the fingertip are mostly encountered by family doctors and emergency physicians [1,3]. Although many injuries require relative simple treatments, nail bed injuries are easily overlooked. If treated inadequately secondary deformities will occur if explained below. It remains difficult to assess the extent of the injury in swollen, painful and bloody fingertips, especially in children as they are less cooperative and less suitable for local anesthetic ring block to aid inspection [6]. Furthermore, an intact nail makes it impossible to assess the damage of the nail bed without removing the nail [6]. In case of nail bed hematomas covering more than 50% of the nail bed without evidence of a fracture and intact nail and nail edges, it is sufficient to evacuate the hematoma through a cauterized hole and leave the nail in place. However, bear in mind that nail bed lacerations may be present in this clinical situation the resulting scar tissue of an unrepaired nail bed laceration can cause deformity and make the patient prone to repetitive infections. The nail plate should be removed in the pres-



Fig. 1. A = Clinical picture of the initial trauma; B + C = X-ray of the initial trauma; D = peroperative image of the nail bed; E = peroperative image of the nail; F = postoperative image after a 7-month follow-up.

Download English Version:

<https://daneshyari.com/en/article/4288835>

Download Persian Version:

<https://daneshyari.com/article/4288835>

[Daneshyari.com](https://daneshyari.com)