



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

A unique complication of radiofrequency therapy to the tongue base

Chrysostomos Tornari^{a,*}, Gentle Wong^a, Asit Arora^a, Bhik Kotecha^a^a Royal National Throat, Nose and Ear Hospital, UK

ARTICLE INFO

Article history:

Received 23 September 2014
 Received in revised form 7 December 2014
 Accepted 27 December 2014
 Available online 8 January 2015

Keywords:

Snoring
 Radiofrequency
 Ablation
 Tongue
 Thyroglossal
 Cyst

ABSTRACT

INTRODUCTION: Radiofrequency ablation treatment of the tongue base can be used either alone or as part of a multilevel approach in the treatment of snoring. This involves the generation of thermal energy around the circumvallate papillae of the tongue. Potential complications include ulceration, dysphagia, haematoma and abscess formation.

PRESENTATION OF CASE: We present the case of a 50-year-old patient who developed an anterior neck swelling following a second application of radiofrequency ablation therapy to the tongue base for snoring. This was secondary to an infection of a previously undiagnosed thyroglossal cyst. The patient made a full recovery following intravenous antibiotic therapy and ultrasound-guided needle aspiration.

DISCUSSION: Thyroglossal tract remnants are thought to be present in seven percent of the adult population. An infection in a thyroglossal tract cyst has not previously been reported following radiofrequency ablation of the tongue base. Given the relatively high complication rate of tongue base radiofrequency ablation in some series, this complication may be under-recognised.

CONCLUSION: An infected thyroglossal tract cyst should be suspected in patients with anterior neck swellings following radiofrequency ablation therapy to the tongue base. We advise caution when performing this procedure on patients with known thyroglossal tract remnants though there is insufficient evidence to suggest that this procedure is contraindicated.

© 2015 The Authors. Published by Elsevier Ltd. on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-SA license (<http://creativecommons.org/licenses/by-nc-sa/4.0/>).

1. Introduction

Marked advances have been made in surgery for snoring and obstructive sleep apnoea, particularly in radiofrequency ablation. This technique utilizes thermal energy from high frequency alternating current to cut and coagulate tissue. It can be used alone, in conjunction with other management strategies such as laser assisted uvulopalatoplasty and mandibular advancement devices, or to assist continuous positive airway pressure (CPAP) usage [1]. Snoring surgery typically requires a multilevel approach to the upper airway and radiofrequency treatment is well suited to this, allowing concurrent applications to the soft palate, base of tongue, and resection of redundant palatopharyngeus mucosa [1,2].

The thyroglossal duct begins at the embryological origin of the thyroid gland, the foramen caecum. This is located at the tongue base between the median lingual swelling (anteriorly) and the copula (posteriorly). The thyroid gland descends through the anterior neck to its final location and its duct usually atrophies before birth. A patent thyroglossal tract can cause a thyroglossal cyst which usu-

ally presents in childhood but can often present after the age of 50 [3].

Radiofrequency treatment of the tongue base is applied in the central posterior third of the tongue. This can cause ulceration, dysphagia, and – more rarely – haematoma or abscess formation which can cause airway compromise [4]. We present an unusual case of radiofrequency ablation therapy to the tongue base causing an infected thyroglossal cyst in a previously undiagnosed and asymptomatic patient. This is, to our knowledge, the first reported case of its kind.

2. Presentation of case

A 50-year-old female patient was referred to the ENT clinic due to snoring. A previous septoplasty for nasal obstruction had improved the nasal airway and the patient had allergic rhinitis without polyposis that was treated by a regular nasal steroid spray. There was no history of a neck lump.

A sleep study showed an apnoea-hypopnoea index of 2.0 and a flow-limitation index of 4.3 with a BMI of 33.3 and an Epworth Sleepiness Score of two. Subsequent sleep nasendoscopy demonstrated the level of obstruction to be at the tongue base with no epiglottic involvement, i.e. grade five [5].

Conservative therapy was not possible as the patient did not have adequate dentition to support a mandibular advancement splint. Therefore, tongue base radiofrequency ablation was com-

Abbreviations: ENT, ear, nose and throat; CT, computed tomography.

* Corresponding author at: Royal National Throat, Nose and Ear Hospital, 330 Grays Inn Road, London, WC1X 8DA, UK. Tel.: +44 2034567890.

E-mail address: t.tornari@gmail.com (C. Tornari).

<http://dx.doi.org/10.1016/j.ijscr.2014.12.039>

2210-2612/© 2015 The Authors. Published by Elsevier Ltd. on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-SA license (<http://creativecommons.org/licenses/by-nc-sa/4.0/>).



Fig. 1. Clinical photographs of the infected thyroglossal cyst on postoperative day 22. Anterior views are shown with neutral tongue position (a) and tongue protrusion (b). The dressing indicates the location of ultrasound-guided needle aspiration.

menced. Six applications of six watts were delivered to the tongue base at the typical location during the first treatment [1]. The patient subsequently reported an approximately 25% improvement in her symptoms.

Two months later, a second-stage procedure was performed with no immediate complications and the patient was discharged from hospital on the evening of surgery. Two days post-operatively, she developed a right submandibular swelling that spread to the right side of the face prompting attendance at a local accident and emergency department. Though there was mild dysphagia, there was no airway compromise and the patient was discharged with oral Co-Amoxiclav for a presumed post-operative infection. A GP appointment was arranged following completion of the seven day course of antibiotics as the neck swelling persisted, though the facial swelling improved. A second course of oral antibiotics was prescribed and, two weeks post-operatively, the patient was discussed with the on call ENT team who arranged urgent review.

ENT review on the 16th postoperative day showed continuing submandibular swelling, redness and discomfort with no dysphagia or difficulty in breathing (Fig. 1). Oropharyngeal examination and fiberoptic nasolaryngoscopy did not demonstrate any tongue base abnormality (Fig. 2). Subsequent review, on the 20th postoperative day, revealed no improvement and the patient was admitted for further investigation.

A CT scan of the neck demonstrated an infected thyroglossal cyst without any tongue base or deep neck collections (Fig. 3) and a subsequent ultrasound-guided needle aspiration reduced the size



Fig. 3. CT scan of the neck on postoperative day 21. An axial, post-contrast CT scan of the neck at the level of C3. The asterisk denotes the position of a 2.7 cm simple cyst with a thin enhancing wall consistent with an infected thyroglossal cyst. Chrysostomos Tornari and Bhik T Kotecha

of the cyst. This improved the patient's symptoms sufficiently to allow discharge four days after admission. The patient remains under review and has not suffered any recurrence.

3. Discussion

In this case, the patient suffered from simple snoring due to tongue base collapse. Radiofrequency treatment of the tongue base was performed according to our standard technique (indicated in Fig. 2) [1]. Compared to the turbinates and palate – two other sites that are commonly treated with radiofrequency therapy – the tongue base carries the highest rate of moderate and severe complications (6.2%) i.e. serious infection requiring drainage or other significant airway compromise [4]. However, there is a considerable variation, ranging from 0% to 32%, in the reported incidence of mild and moderate complications. The reason for this variability is not known [4].

There are reports of post-operative infections that required drainage following tongue base radiofrequency ablation [4]. However, there is no report of this procedure causing a thyroglossal cyst infection. Thyroglossal cysts are the most common congenital neck cysts and the most common cause of midline neck swellings in the paediatric population though, in terms of age, there is a bimodal distribution for their identification with similar incidence in children and adults [6,7]. They can arise anywhere along a thyroglossal tract remnant and it is thought that such remnants are present in seven




Fig. 2. Radiofrequency ablation of the tongue base. The radiofrequency ablation probe is shown (a) along with a fiberoptic nasendoscopic photograph of the tongue base on postoperative day 23. Asterisks indicate the location of radiofrequency applications according to the operative note.

Download English Version:

<https://daneshyari.com/en/article/4289159>

Download Persian Version:

<https://daneshyari.com/article/4289159>

[Daneshyari.com](https://daneshyari.com)