



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Ruptured spontaneous splenic artery aneurysm: A case report and review of the literature

Aisha Abdulrahman^a, Alaa Shabkah^a, Mazen Hassanain^b, Murad Aljiffry^{a,*}^a Department of Surgery, College of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia^b Department of Surgery, College of Medicine, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Article history:

Received 9 June 2014

Received in revised form 7 August 2014

Accepted 24 August 2014

Available online 2 September 2014

Keywords:

Splenic artery

Aneurysm

Rupture

Spontaneous

Idiopathic

ABSTRACT

INTRODUCTION: Splenic artery aneurysm is a rare condition, however, potentially fatal. The importance of splenic artery aneurysm lies in the risk for rupture and life threatening hemorrhage.

PRESENTATION OF CASE: This is a case of a ruptured splenic artery aneurysm in a 58-year-old lady. She presented with hypovolemic shock and intra-peritoneal bleeding. Diagnosis was confirmed by CT angiography and she was managed by operative ligation of the aneurysm with splenectomy and distal pancreatectomy.

DISCUSSION: The literature pointed the presence of some risk factors correlating to the development of splenic artery aneurysm. In this article we discuss a rare case of spontaneous (idiopathic) splenic artery aneurysm and review the literature of this challenging surgical condition.

CONCLUSION: Splenic artery aneurysm needs prompt diagnosis and management to achieve a favorable outcome, high index of suspicion is needed to make the diagnosis in the absence of known risk factors.

© 2014 The Authors. Published by Elsevier Ltd. on behalf of Surgical Associates Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

1. Introduction

Splenic artery aneurysm is a rare condition with a prevalence of 1%.¹ It accounts for about 60% of all splanchnic arterial aneurysms. It only follows aortic and iliac arteries aneurysms as the third most common intra-abdominal aneurysm.¹ The significance of splenic artery aneurysm lies in the potential risk for rupture and life threatening hemorrhage which occurs in 10% of cases with a mortality rate of 10–25% in non-pregnant patient and up to 70% during pregnancy.²

A true aneurysm is different from pseudoaneurysms. In a true aneurysms, all the layers of the vessel are intact with a thinned and dilated wall. On the other hand, a pseudoaneurysms or false aneurysms result from a tear in the vessel wall with subsequent formation of a peri-arterial hematoma and is usually secondary to a local inflammatory process such as pancreatitis.³

To date the literature pointed the presence of some risk factors correlating to the development of splenic artery aneurysm such as fibromuscular dysplasia, collagen vascular diseases, female gender, history of multiple pregnancies and portal hypertension.⁴ We describe a case of spontaneous (idiopathic) splenic artery aneurysm

whom presented uniquely without any previously known risk factor.

2. Presentation of case

58-year-old female, presented to ER compiling of epigastric abdominal pain and nausea for one day. There were no associated symptoms nor there was a history of previous similar episodes. Her past medical history includes cavernous sinus venous thrombosis for which she was on lifelong warfarin anticoagulation and she did not have any previous surgeries. Initial physical examination was unremarkable. Her initial lab test showed a hemoglobin of 13.6 g/dl and INR of 6.6 IU. Abdominal ultrasound did not show any obvious abnormality. The patient was admitted for a possible upper GI endoscopy if symptoms did not resolve after correction of the coagulopathy. Twelve hours following admission the patient complained that her pain suddenly became severe. On examination her heart rate became 110/min and the blood pressure dropped to 90/50 mmHg. Her abdomen became distended with mild general tenderness and her repeated hemoglobin was 6.9 g/dl. Patient was resuscitated and stabilized then shifted to radiology suit. CT angiogram showed large amount of free intra peritoneal fluid and a splenic artery aneurysm with intravenous contrast extravasation consisting of intra peritoneal hemorrhage (Figs. 1 and 2).

The patient was appropriately resuscitated with blood and plasma transfusions and booked for an emergency exploration.

* Corresponding author at: Department of Surgery, College of Medicine, King Abdulaziz University Hospital, P.O. Box 80215, Jeddah 21589, Saudi Arabia.

Tel.: +966 012 6401000x18209; fax: +966 012 6408347.

E-mail address: dr.aljiffry@gmail.com (M. Aljiffry).



Fig. 1. CT abdomen showing intra peritoneal hematoma.



Fig. 2. CT abdomen showing splenic artery aneurysm and contrast extravasation.

In the operating room a midline laparotomy incision was performed. Upon entering the peritoneal cavity a moderate amount of blood and clots were encountered. After proper exposure, a large hematoma was seen in the lesser sac bulging through the




Fig. 3. Excised specimen showing the splenic artery (artery clamp through) and the site of the aneurysm rupturing into the pancreas close to the hilum.

gastro-colic omentum. The supra-celiac aorta was identified by dissecting through the gastro-hepatic ligament to assure the availability of vascular control. Exploration of the lesser sac hematoma revealed a ruptured aneurysm involving the distal part of the splenic artery close to the hilum of the spleen, with a total amount of evacuated blood of 3 L. The splenic artery was ligated proximally, followed by a splenectomy with distal pancreatectomy to excise the aneurysm that was inseparable from the pancreatic tail (Fig. 3). As the patient was in hypovolemic shock a damage control approach was taken to expedite the transfer of the patient to the ICU for hemodynamic management and overall support. We believed that a splenectomy was the procedure of choice to accomplish the control of the bleeding in this patient, rather than a splenic tissue preserving procedure. The patient had uneventful postoperative course and discharged home on the 7th postoperative day. The histopathology report confirmed the diagnosis of a true splenic artery aneurysm.

3. Discussion

Splenic artery aneurysm was first described by Beaussier in 1770.⁵ The incidence of splenic artery aneurysm in large autopsy series found to be between 0.01% and 0.2%. The incidence

Table 1
Summary of all reported cases of spontaneous (idiopathic) splenic artery aneurysm.

Year	Author	Age (years)	Sex	Presentation	Method of diagnosis	Procedure	Outcome
1990	Lie et al. ¹⁶	71	M	Abdominal pain/UGIB	Angiography	Distal pancreatectomy	Survived
1994	Willis et al. ¹⁷	86	F	UGIB	Not reported	Ligation of the aneurysm	Survived
2000	Deshpand et al. ¹⁸	62	F	Abdominal pain	Intra-operative	Distal pancreatectomy	Survived
		27	M	Shock	Intra-operative	Splenectomy	Died
2001	Jamsheer et al. ¹⁹	30	M	Shock	CT scan	Splenectomy	Survived
2005	Tsokos et al. ²⁰	33	M	Shock	Autopsy	Not done	Died
2006	Afridi et al. ²¹	18	F	Abdominal pain/shock	Intra-operative	Splenectomy	Survived
2007	Mattick ²²	47	M	Abdominal pain/shock	Intra-operative	Splenectomy	Survived
2009	Betal et al. ²³	58	M	Epigastric pain/shock	CT scan	Splenectomy	Survived
2009	Zubaidi ²⁴	42	F	LUQ pain/hematemesis	CT scan	Splenectomy/embolization of proximal aneurysm	Survived
2009	Sezgin et al. ¹¹	22	F	LUQ pain	Intra-operative	splenectomy	Survived
2009	Ousadden et al. ¹⁴	36	F	hematemesis	Intra-operative	Splenectomy	Survived
2013	Katirci et al. ²⁵	46	F	LUQ pain	CT scan	Splenectomy	Survived
2014	Oakley et al. ²⁶	33	M	Generalized body pain/shock	Intra-operative	Splenectomy	Survived
		48	M	Abdominal pain/shock	Intra-operative	Splenectomy	Survived

Download English Version:

<https://daneshyari.com/en/article/4289243>

Download Persian Version:

<https://daneshyari.com/article/4289243>

[Daneshyari.com](https://daneshyari.com)