

Aortic arch aneurysm repair using the Najuta stent graft in a challenging compromised seal zone

Naoki Toya, MD,^a Kota Shukuzawa, MD,^a Soichiro Fukushima, MD,^a Masamichi Momose, MD,^a Tadashi Akiba, MD,^b and Takao Ohki, MD,^c *Chiba and Tokyo, Japan*

We report the case of a 67-year-old patient with an anatomically complex aneurysm of the aortic arch treated by fenestrated thoracic endovascular aortic repair with subclavian-carotid extrathoracic bypass. We used the Najuta thoracic stent graft, which was approved for use in January 2013 in Japan and successfully excluded the aneurysm. Our case shows that the Najuta stent graft procedure is a feasible treatment if open repair is unsuitable for cases of aortic arch aneurysm with a challenging compromised seal zone. (*J Vasc Surg Cases* 2016;2:21-4.)

A major issue in thoracic endovascular aortic repair (TEVAR) of aortic arch aneurysm with a short proximal neck is coverage of the origin of arch branches to achieve an adequate proximal landing zone.¹ However, in aortic arch aneurysm with a short proximal neck, stent graft coverage of all arch branches, including the innominate artery, is required. A fenestrated stent graft in the aortic arch could achieve a better sealing zone in the challenging compromised proximal seal zone. Hybrid fenestrated TEVAR using the Najuta thoracic stent^{2,3} graft has been successfully performed. Here, we present a similar case from our facility and relevant literature review. The consent of the patient was obtained to publish this report.

CASE REPORT

A 67-year-old woman had undergone replacement of an ascending thoracic aorta at another institution in 2014 for treatment of an acute type A dissection caused by a cardioplegia needle after aortic valve replacement for an aortic valve disease. She had also undergone right axillofemoral (Ax-F) bypass because of intraoperative right external iliac artery injury during surgery for ovarian cancer. In February 2015, follow-up computed tomography (CT) revealed a rapidly growing aneurysm in the distal arch involving the left subclavian artery (LSCA) with a short proximal neck and extreme proximal and innominate artery angulation (*Fig 1, A*). The patient also had an anomalous origin of the left vertebral artery from the aortic arch. Because of careful imaging of the right

vertebral artery up to the basilar artery, we concluded that sufficient posterior cerebral circulation would be maintained if the left vertebral artery was closed without surgical revascularization.⁴ Because of two prior sternotomies, it would be a high surgical risk, and a redo operation was unsuitable for this patient; she was treated by fenestrated TEVAR using the Najuta thoracic stent graft (Kawasumi Laboratories, Inc, Tokyo, Japan; *Fig 1, B and C*) and a debranching bypass.

The procedure was performed under general anesthesia. First, an 8-mm heparin-bonded expanded polytetrafluoroethylene (Propaten; W. L. Gore & Associates, Flagstaff, Ariz) extrathoracic bypass (Ax-F graft-left common carotid artery [LCCA]-LSCA bypass) was performed. The proximal LCCA was ligated (*Fig 1, D*).

The left femoral artery was exposed, and a 6F guiding sheath was inserted from the Ax-F bypass graft into the innominate artery (*Fig 1, D*). A 0.032-inch Radifocus guidewire (Terumo, Tokyo, Japan) was then pulled through from the right subclavian artery to the left femoral artery. The proximal LSCA was occluded with an Amplatzer vascular plug (St. Jude Medical, St. Paul, Minn) to prevent a type II endoleak (*Fig 1, D*). The Najuta was delivered with a 22F J-shaped sheath maintained under continuous strain by traction at both wire ends (body floss technique).^{2,5-7} Because of a severe angulation and tortuosity of the ascending aorta and innominate artery due to prior ascending aorta replacement, the delivery of the system through the arch was difficult by only a body floss technique. Therefore, the guiding sheath through the Ax-F bypass was deeply inserted into the origin of the innominate artery, and it supported the delivery of the stent graft. The Najuta was then delivered and deployed at a zone 0 proximal site with delicate positional adjustment of the fenestration of the Najuta to the innominate artery. In this case, the stent graft had two fenestrations (*Fig 1, B and C*). The first fenestration was opened from the first marker to the second marker, and the second fenestration was opened between the second marker and the third marker (*Fig 2, A*). The third marker was accurately put on the distal end of the innominate artery, and the fenestration was fitted for the branch automatically without rotation (*Fig 2, B*). The LCCA, LSCA, and left vertebral artery were excluded. We exchanged the wire from a 0.032-inch Radifocus wire to a Lunderquist extrastiff wire (Cook Medical, Bloomington, Ind). In addition, the Relay stent graft (Bolton Medical, Barcelona, Spain) was deployed at a zone 1 proximal site. The proximal

From the Division of Vascular Surgery, Department of Surgery,^a and the Department of Surgery,^b The Jikei University Kashiwa Hospital, Chiba; and the Division of Vascular Surgery, Department of Surgery, The Jikei University School of Medicine, Tokyo.^c

Author conflict of interest: none.

Correspondence: Naoki Toya, MD, 163-1Kashiwashita, Kashiwa-shi, Chiba 277-8567, Japan (e-mail: toyanaoki@gmail.com).

The editors and reviewers of this article have no relevant financial relationships to disclose per the Journal policy that requires reviewers to decline review of any manuscript for which they may have a conflict of interest.

2352-667X

Copyright © 2016 The Authors. Published by Elsevier Inc. on behalf of the Society for Vascular Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<http://dx.doi.org/10.1016/j.jvsc.2016.02.004>




Fig 1. **A**, Preoperative contrast-enhanced computed tomography (CT) showed a rapidly growing aneurysm in the distal arch involving the left subclavian artery (LSCA) with a short proximal neck and right axillofemoral (Ax-F) bypass (arrow). *a*, Aneurysm; *b*, Dacron tube graft; *c*, innominate artery. **B**, The Najuta fenestrated stent graft is a customized fenestrated device comprising a self-expandable stainless steel Z-stent and an expanded polytetrafluoroethylene graft. **C**, A fenestration of the Najuta to the innominate artery (arrow). **D**, Surgical strategy (schema); white arrow, guiding sheath; black arrow, delivery sheath of the Najuta.




Fig 2. The first fenestration was opened from the first marker to the second marker, and the second fenestration was opened between the second marker and the third marker. **A**, Before deployment (arrows, markers). **B**, After deployment (arrows, markers).

bare stent of the Relay stent graft was inserted into the prosthetic graft of the ascending aorta.

Completion angiography showed complete exclusion of the aneurysm and confirmed preservation of the innominate artery and debranching bypass flow. The patient's postoperative course was uneventful. Six months after the procedure, postoperative

CT revealed that the aneurysm had been completely excluded (Fig 3, A and B). She recovered without any complications.

DISCUSSION

The hybrid repair of arch lesions for zone 0 is feasible in patients unsuitable for open repair.⁸ On the other hand,

Download English Version:

<https://daneshyari.com/en/article/4305508>

Download Persian Version:

<https://daneshyari.com/article/4305508>

[Daneshyari.com](https://daneshyari.com)