

Lack of the cerebral peduncle involvement in a series of adult supratentorial AVM: A diffusion tensor imaging study

Bing Fu^a, Jizong Zhao^{a,*}, Bo Wang^b, Mingqi Yang^a, Long Xu^a, Yan Zhuo^b

^a Department of Neurosurgery, Beijing Tiantan Hospital, Capital Medical University, Beijing 100050, China

^b State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese Academy of Sciences, No. 15 Datun Road, Beijing 100101, China

ARTICLE INFO

Article history:

Received 24 February 2010

Received in revised form 17 August 2010

Accepted 1 September 2010

Keywords:

Arteriovenous malformation

Cavernous malformation

Cerebral peduncle

Diffusion tensor imaging

ABSTRACT

Congenital as arteriovenous malformation (AVM) is, most patients with AVM would be asymptomatic until adults. During the past 2 years, 23 cases of adult supratentorial AVM patients had DTI after admission. The region of interest was placed in the cerebral peduncle. Their FA value and fiber number was compared with those of cavernous malformation (CM) and tumor (glioma and meningioma). In the AVM group, there was no significant difference in FA of the cerebral peduncle (ipsilateral 0.758 ± 0.055 versus contralateral 0.755 ± 0.049 ; $P > 0.05$) and fiber number (319.6 ± 82.9 versus 304.7 ± 89.1 ; $P > 0.05$). In the CM group, FA of the cerebral peduncle on ipsilateral side (0.711 ± 0.092) was significantly lower than that of contralateral side (0.768 ± 0.043) ($P < 0.01$). Similar result was in fiber number of the CM group (251 ± 82.1 versus 307.3 ± 77.0 ; $P < 0.05$). In tumor group, FA of ipsilateral side (0.713 ± 0.084) was lower than that of contralateral (0.751 ± 0.052) without significant difference. There was no significant difference in fiber number between ipsilateral and contralateral sides in the tumor group (308.9 ± 112.4 versus 287.9 ± 62.4). Unlike non-AVM lesions (CM and tumor), FA value and fiber number of the ipsilateral cerebral peduncle is less influenced in the AVM group. The lack of the cerebral peduncle involvement indicates that there is plasticity of white matter in AVM.

© 2010 Elsevier Ireland Ltd. All rights reserved.

Cerebral arteriovenous malformation (AVM) is usually thought to be congenital. There were few reports that AVM is de novo [9] or regresses spontaneously [3]. Hemorrhage, headache, and seizure are the three common presentations in cerebral AVM [36]. But most patients with AVM would be asymptomatic until adults.

Fractional anisotropy (FA) has been used to characterize water diffusion anisotropy of brain in diffusion tensor imaging (DTI). FA depends on organization and myelination of nerve tract [7,8]. Thus, DTI measurements of the cerebral peduncle may be used to characterize microstructure abnormalities in AVM and non-AVM lesions. In the present study, we aimed at investigating the cerebral peduncle involvement in AVM and non-AVM lesions using DTI.

During the past 2 years, there were 51 cases of adult with supratentorial lesions, including AVM, cavernous malformation (CM) and tumor (glioma and meningioma), who had DTI after admission. The mean age of the AVM group ($n = 23$) was 33.6 years (range: 21–51 years). Cases of CM ($n = 14$; mean age 37.4 years; range: 20–57 years) and tumors ($n = 14$, 10 cases of glioma and 4 cases of meningioma; mean age 46.1 years; range: 24–63 years) were used as control. The ratio of male to female was 15:8, 9:5, and 10:4,

respectively. All lesions were solitary without involvements of the corpus callosum and the midbrain. Except two patients (one in the AVM group and one in CM) who could not accept surgical risks, 49 patients underwent operation with informed consent, whose diagnoses were confirmed by histology.

On a Siemens 3.0 Tesla MRI Scanner, the diffusion-weighted data were acquired using a single-shot spin-echo diffusion sensitized echo-planar imaging (EPI) sequence with 12 encoding directions, a diffusion sensitization of $b = 1000 \text{ s/mm}^2$, TR of 3 s, and TE of 93 ms. EPI image distortion artifacts were reduced using GRRAPA with acceleration factor or k -space undersampling of $R = 2$. The slice thickness was 4 mm with 22 axial slices covering the whole brain (foramen magnum to vertex), a field of view of $220 \text{ mm} \times 220 \text{ mm}$, and an image matrix of 128×128 that matched sequences described above. The number of image averages was 4. The total DTI acquisition time was approximately 2.8 min.

All of the image processing was performed using the software MedINRIA (<http://www-sop.inria.fr/asclepios/software/MedINRIA>). Regions of interest (ROI) were selected according to FA maps. First, we determined the image with the highest FA value of the cerebral peduncle in the axial plane FA maps. Then a 3×3 voxels ROI was placed in the cerebral peduncle to produce the highest average FA value (Fig. 1). The number of fibers through the ROI was calculated using the software. The FA threshold value was 0.20 to keep tracking based on the connectivity of the neighborhood.

* Corresponding author. Tel.: +86 010 65113440; fax: +86 010 65113440.
E-mail address: zhaojz@public.bta.net.cn (J. Zhao).




Fig. 1. Selection of the ROI in FA map. Angiography demonstrates a case of arteriovenous malformation (AVM) in the left parieto-occipital lobe (on the left). Region of interest (ROI) (in red) is placed in the cerebral peduncles in fractional anisotropy (FA) map (on the right). The FA was 0.798 in the ipsilateral cerebral peduncle and 0.799 in the contralateral cerebral peduncle.

All data are presented as means \pm SD and compared using *t* test. Differences were considered statistically significant at a *P* value <0.05 .

In the AVM group, the complaints were epilepsy (11 cases), headache (8 cases) and neurological deficit (5 cases). In the CM group, the complaints included neurological deficit (7 cases), headache (3 cases), epilepsy (3 cases) and asymptomatic (1 case). In the tumor (glioma and meningioma) group, patients exhibited neurological deficit (6 cases), headache (4 cases), epilepsy (4 cases) and asymptomatic (1 case). The maximum diameter of lesions was 33.9 ± 11.3 mm (range: 18–55 mm) in the AVM group, 32.2 ± 11.2 mm (14–60 mm) in the CM group and 40.9 ± 15.3 mm (22–67 mm) in the tumor group.

In the AVM group, there was no statistically difference between mean FA of the ipsilateral cerebral peduncle (0.758 ± 0.055) and that of the contralateral cerebral peduncle (0.755 ± 0.049) ($P > 0.05$). In the tumor group, mean FA was 0.713 ± 0.084 of the ipsilateral cerebral peduncle and 0.751 ± 0.052 of the contralateral cerebral peduncle ($P > 0.05$). In the CM group, the FA measured in the ipsilateral cerebral peduncle (0.711 ± 0.092) was significantly ($P < 0.01$) lower as compared with the contralateral cerebral peduncle (0.768 ± 0.043). A significant reduction was also observed in the ipsilateral FA of non-AVM lesions (the CM group and the tumor group) as compared to that of AVM ($P < 0.05$). But there were no statistically differences in the contralateral FA between groups (Fig. 2).

The potential effects of age, sex, side and lesion location on the DTI measurements were also investigated. All patients (mean age 38.1 ± 11.9 years, $n = 51$) were divided into four groups according to their age (<31 years, 31–40 years, 41–50 years, and >50 years). Mean FA of the contralateral cerebral peduncle was 0.771 ± 0.044 ($n = 14$), 0.744 ± 0.049 ($n = 19$), 0.771 ± 0.042 ($n = 8$) and 0.752 ± 0.055 ($n = 10$), respectively. There was no significant difference between groups ($P > 0.05$). The mean FA of the contralateral cerebral peduncle was 0.760 ± 0.043 in male ($n = 34$) and 0.752 ± 0.057 in female ($n = 17$) without statistically difference ($P > 0.05$). Whether the side was the left (0.753 ± 0.050 , $n = 30$) or the right (0.763 ± 0.045 , $n = 21$) did not affect the FA value of the contralateral cerebral peduncle (Fig. 3). There were 8 cases of AVM and 4 cases of CM that involved temporal lobe. The FA of the cerebral peduncle was 0.786 ± 0.048 on ipsilateral side and 0.744 ± 0.056 on

contralateral side in the temporal lobe AVM group. In the temporal lobe CM group, FA was 0.727 ± 0.022 and 0.770 ± 0.015 , respectively. There was significant difference between bilateral side in the temporal lobe CM ($P < 0.05$), but not in the temporal lobe AVM ($P > 0.05$). By comparing the results of ipsilateral side, the difference between the temporal lobe AVM and CM was statistically significant ($P < 0.05$). No contralateral side difference was found between ($P > 0.05$) the temporal lobe AVM and CM (Fig. 3).

In the AVM group, the number of fibers through the ROI was 319.6 ± 82.9 on ipsilateral side and 304.7 ± 89.1 on contralateral side ($P > 0.05$). In the CM group, fiber number of ipsilateral side (251 ± 82.1) was significantly lower than that of contralateral side (307.3 ± 77.0) ($P < 0.05$). In tumor, fiber number was 308.9 ± 112.4 and 287.9 ± 62.4 without significant difference between bilateral side ($P > 0.05$) (Fig. 4).

In literatures, DTI had been used in the evaluation of AVM surgery and radiotherapy. DTI could demonstrate the relationship of nerve tracts with AVM [12,17,33]. Some nerve tracts were less




Fig. 2. FA of the cerebral peduncle in the AVM, cavernous malformation (CM) and tumor group. In the AVM group, FA of the cerebral peduncle is 0.758 ± 0.055 on ipsilateral side and 0.755 ± 0.049 on contralateral side. In the CM group, FA is 0.711 ± 0.092 and 0.768 ± 0.043 . In the tumor group, FA is 0.713 ± 0.084 and 0.751 ± 0.052 . There is significantly difference between bilateral side in the CM group ($P < 0.01$), but not in AVM or tumor ($P > 0.05$).

Download English Version:

<https://daneshyari.com/en/article/4345446>

Download Persian Version:

<https://daneshyari.com/article/4345446>

[Daneshyari.com](https://daneshyari.com)