

## Early music exposure modifies GluR2 protein expression in rat auditory cortex and anterior cingulate cortex

Feng Xu, Rui Cai, Jinghong Xu, Jiping Zhang, Xinde Sun\*

*College of Life Sciences, Institute of Cognitive Neuroscience, East China Normal University, Shanghai 200062, China*

Received 15 March 2007; received in revised form 29 April 2007; accepted 1 May 2007

### Abstract

GluR2, a major subunit in AMPA receptor, plays an important role in brain functional activity. We studied the effect of music exposure during development on the expression level of GluR2 proteins in the auditory cortex (AC) and anterior cingulate cortex (ACC) of SD rats. Rats were divided into three groups, Music1 (exposed to Nostalgia) group, Music2 (exposed to Wishmaster) group, and control (no music exposure) group. For music exposure groups, rats were exposed to music from postnatal day (PND) 14, and the expression levels of GluR2 proteins were determined at PND 28, 42 and 56. For the control group, the expression levels of GluR2 proteins were determined at PND 1, 3, 5, 7, 9, 11, 14, 21, 28, 42, and 56. Results showed an age-dependent expression of GluR2 proteins in control rats. In AC, exposure to Music2 dramatically increased the expression of GluR2, while exposure to Music1 had no effect. In ACC, we found remarkable discrepancies in time-dependent expression of GluR2 between music exposed rats and control rats. These results indicate that exposure to music can modify the expression level of GluR2 protein in AC and ACC.

© 2007 Elsevier Ireland Ltd. All rights reserved.

**Keywords:** Rat; Auditory cortex; Anterior cingulate cortex; Music exposure; GluR2; Western blotting

Glutamate receptors (GluRs) mediate most of the excitatory neurotransmission in the central nervous system (CNS). They also participate in plastic changes in the efficacy of synaptic transmission underlying learning, memory and formation of neural networks during development [12,17]. AMPA receptor, one type of GluRs, consists of four subunits (named GluR1–GluR4). It mediates fast excitatory neurotransmission at a majority of synapses in the CNS. Previous studies have shown that the distribution and density of GluR1–GluR4 exhibited regional expression patterns in adult rat brain. The expression pattern also changes during developmental stages. In the cerebral cortex, the expression patterns of GluR1, GluR3 and GluR4 mRNAs differ among layers, while GluR2 mRNAs are even [15].

GluR2 subunit has a relatively lower  $\text{Ca}^{2+}$  permeability compared with other subunits. Glutamate-induced  $\text{Ca}^{2+}$  entry occurs through three kinds of channels: NMDA receptor channel,  $\text{Ca}^{2+}$ -permeable AMPA receptor channel, and voltage-dependent  $\text{Ca}^{2+}$  channel [6]. Any changes in the channels will ultimately influence the  $\text{Ca}^{2+}$  influx. It has been demonstrated that exces-

sive  $\text{Ca}^{2+}$  entry under pathological conditions leads to neuronal cell death [10]. Because GluR2 regulates  $\text{Ca}^{2+}$  permeability, we speculate that GluR2 may participate in the regulation of synaptic plasticity.

Studies on rats have shown that music exposure during pregnancy resulted in elevated neurogenesis in hippocampus and enhanced spatial learning ability [9]. Auditory deprivation or auditory experience also affects cortical plasticity [2,13]. Therefore, sound or music does have impacts on brain plasticity. However, it remains unclear whether music exposure can modify the expression of GluR2 subunit. In the present study, we investigate the music effects on the expression level of GluR2 protein in auditory cortex (AC) and anterior cingulate cortex (ACC) in Sprague-Dawley (SD) rats.

Sprague-Dawley rats were used in this study. The rats were raised in a sound-attenuated room with a background noise level less than 45 dB SPL re 20  $\mu\text{Pa}$ . The inside room was maintained at a temperature of  $22 \pm 3^\circ\text{C}$  and on a light/dark cycle with 12 h light throughout the experiment.

In experiment 1, 44 rats were used. The GluR2 protein expression in AC and ACC were measured at postnatal days (PNDs) 1, 3, 5, 7, 9, 11, 14, 21, 28, 42, and 56. Each sampling age included four rats.

\* Corresponding author. Tel.: +86 21 62232775.  
E-mail address: [xdsun@bio.ecnu.edu.cn](mailto:xdsun@bio.ecnu.edu.cn) (X. Sun).




Fig. 1. A series of coronal sections of the rat brain. The extent of separated areas of auditory cortex (AC) region (a) and anterior cingulate cortex (ACC) region (b) are shown in grey.

In experiment 2, 45 SD rats were divided into three groups: Music1 (exposed to Nostalgia) group, Music2 (exposed to Nightwish) group and control group. Each group included 15 rats. The Music1 or Music2 exposure (70 dB SPL) was started at postnatal day 14 (PND14) in a sound attenuated room throughout the dark period for 12 h per day. After music exposure, the rats were put back in the sound-attenuating room in which the control rats were raised. The control rats were not exposed to music throughout the experiment. For each group, the expression of GluR2 protein in AC and ACC was measured at PND28, 42 and 56, with five rats at each sampling age.

SD rats were deeply anaesthetized with injection of sodium pentobarbital (50 mg/kg BW). Immediately after decapitation, the brains were obtained. The right and left AC regions (−3.30 to −6.30 mm anterior to bregma, according to Paxinos and Watson 1998; also according to the blood vessels in brain surface) and ACC regions (3.70 to −1.40 mm anterior to bregma) were separated (Fig. 1), and put in buffer equals to 10 times the volume (137 mmol/l NaCl, 20 mmol/l Tris, 1.5 mmol/l  $\text{Na}_3\text{VO}_4$ , 1% NP-40, 10% glycerol, freshly added with 1 mmol/l PMSF, 10  $\mu\text{g}/\text{ml}$  aprotinin, 0.2  $\mu\text{g}/\text{ml}$  leupeptin). Centrifugation was done at 16,000 rpm for 10 min after homogenization. The supernatants were then collected. All the above procedures were done at 4 °C. Concentration was measured by Bradford methods: the standards contained a range of 0–100  $\mu\text{g}$  protein (BSA) in 100  $\mu\text{l}$  volume, and were added to 5 ml Coomassie Brilliant Blue G-250 (100 mg G-250 in 50 ml 95% ethanol, and 100 ml 85% (w/v) phosphoric acid). After 3 min incubating, the absorbance at 595 nm was measured. The standard curve is:  $Y = 555.56X - 0.04$ . Samples were boiled at 100 °C in the presence of sample buffer (included 250 mM Tris-HCl, pH 6.8, 4% sodium dodecyl sulfate (SDS), 1%  $\beta$ -mercaptoethanol, 1% bromophenol blue, and 20% glycerol) for 5 min, and the final concentration of sample protein was adjusted to 5  $\mu\text{g}/\mu\text{l}$ . Samples were preserved at −80 °C.

Proteins (15  $\mu\text{l}$  of each sample) were separated on a 7.5% SDS-polyacrylamide gel and transferred to a nitrocellulose membrane (PALL). Blocked with 5% non-fat dry milk dissolved in 1% TBST for 2 h, the immobilized proteins were incubated at 4 °C for 12 h with primary antibody (goat-anti-rat GluR2 antibody, 1:200, Santa Cruz; goat actin antibody, 1:2000, Santa Cruz). After being washed with 1% TBST, membranes were incubated with secondary antibody (rat-anti-goat IgG HRP, 1:1000, Jacksonimmuno) for 2 h at room temperature, and then washed again. Proteins were visualized by using enhanced chemiluminescence reagents (Pierce) for 5 min fol-




Fig. 2. The specificity and the band location of protein loading. The molecular weight of GluR2 is about 108 kDa.

lowed by exposure to medical X-ray film (Kodak) for 20 min. Blots were scanned, and analyzed by using Scanband software.

The band intensity for each sample was calculated as total gray value. The values were then normalized to the actin gray value. The statistical significance was determined by *t*-test in SigmaPlot software.

The specificity and the band location of GluR2 protein loading was shown in Fig. 2. Expression of GluR2 proteins in AC (Fig. 3a, Table 1) and ACC (Fig. 3b, Table 1) exhibited age-dependent increasing trend. Evidence showed that excitatory synaptic response mainly depended on NMDA receptors in infant, and the role of NMDARs showed age-related decrease

Table 1  
GluR2 protein expression level in auditory cortex and anterior cingulate cortex from PND7 to PND56 (nmol/mg)

Postnatal days	Auditory cortex (n = 4)	Anterior cingulate cortex (n = 4)
1	79.68 ± 3.11	70.02 ± 11.10
3	92.58 ± 12.64	128.52 ± 13.08
5	127.70 ± 18.40	160.65 ± 10.66
7	154.88 ± 23.33	182.98 ± 6.91
9	201.04 ± 36.78	223.65 ± 7.49
11	251.21 ± 41.53	273.30 ± 10.74
14	272.94 ± 54.19	376.26 ± 58.52
21	428.41 ± 64.32	598.17 ± 107.26
28	485.31 ± 69.84	740.27 ± 140.00
42	603.19 ± 150.29	936.96 ± 122.25
56	889.78 ± 131.70	1503.66 ± 303.05

Download English Version:

<https://daneshyari.com/en/article/4349420>

Download Persian Version:

<https://daneshyari.com/article/4349420>

[Daneshyari.com](https://daneshyari.com)