

PICK1 is not a susceptibility gene for schizophrenia in a Japanese population: Association study in a large case–control population

H. Ishiguro^{a,b,*}, M. Koga^{a,b}, Y. Horiuchi^{a,b}, T. Inada^c, N. Iwata^d, N. Ozaki^e,
H. Ujike^f, T. Muratake^g, T. Someya^g, T. Arinami^{a,b}

^aDepartment of Medical Genetics, Doctoral Program in Social and Environmental Medicine, Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki 305-8575, Japan

^bCore Research for Environmental Science and Technology (CREST), Japan Science and Technology Agency, Kawaguchi-shi, Saitama 332-0012, Japan

^cDepartment of Psychiatry, Teikyo University School of Medicine, Chiba Medical Center, Anesaki 3426-3, Ichihara-shi, Chiba 299-0111, Japan

^dDepartment of Psychiatry, Fujita Health University School of Medicine, Toyoake, Aichi 470-1192, Japan

^eDepartment of Psychiatry, School of Medicine, Nagoya University, Nagoya, Aichi 466-8550, Japan

^fDepartment of Neuropsychiatry, Okayama University, Graduate School of Medicine, Dentistry & Pharmaceutical Sciences, 2-5-1 Shikata-cho, Okayama 700-8558, Japan

^gDepartment of Psychiatry, Niigata University Graduate School of Medical and Denatal Sciences, Niigata 951-8510, Japan

Received 12 January 2007; accepted 8 February 2007

Available online 20 February 2007

Abstract

The protein interacting with C-kinase 1 (PICK1) has been implicated in the susceptibility to schizophrenia. PICK1 interacts with enzymes and receptors that play roles in the pathogenesis of schizophrenia via glutamatergic dysfunction. Recently, two studies reported associations between schizophrenia and two *PICK1* gene polymorphisms, rs3952 in Chinese and Japanese populations and rs2076369 in a Japanese population. We attempted to confirm these associations in a case–control study of 1765 Japanese patients with schizophrenia and 1851 Japanese control subjects. Neither polymorphism was associated with schizophrenia (rs3952, $p = 0.755$; rs2076369, $p = 0.997$). A haplotype block with these polymorphisms spanning the 5' region of the *PICK1* gene showed high linkage disequilibrium in the Japanese population ($D' = 0.98$, $r^2 = 0.34$); however, neither haplotype was significantly associated with schizophrenia. We conclude that the common haplotypes and polymorphisms of the *PICK1* gene identified thus far are unlikely to contribute to genetic susceptibility to schizophrenia in the Japanese population.

© 2007 Elsevier Ireland Ltd and the Japan Neuroscience Society. All rights reserved.

Keywords: Schizophrenia; PICK1; Japanese; Glutamate; Case–control study

1. Objective of the study

The protein interacting with C-kinase 1 (PICK1) is also known as protein kinase C- α -binding protein (PRKCABP). PICK1 contains a PDZ domain and a conserved arfaptin homology (AH) domain. The PDZ domain of rat Pick1 interacts with splice variants of AMPA receptor subunits (Dev et al., 1999; Lu and Ziff, 2005; Sossa et al., 2006), and PICK1 appears to modulate synaptic

transmission by regulating AMPA receptors in brain (Xia et al., 1999). AMPA receptors can improve cognitive function in patients with schizophrenia (Coyle et al., 2002). PICK1 also colocalizes and interacts with dopamine transporter (DAT) (Torres et al., 2001), and metabotropic glutamate receptor 7 (GRM7) (Boudin et al., 2000; Dev et al., 2001; Enz and Croci, 2003). More recently, Fujii et al. (2006) found that PICK1 interacts with serine racemase, which is a D-serine synthesizing enzyme that may play a role in the pathogenesis of schizophrenia. Taken together, the functional and structural features of PICK1 suggest that it may play a role in schizophrenia, because glutamatergic and dopaminergic dysfunction are thought to be involved in one of the most important neural pathways underlying vulnerability to the disease.

* Corresponding author at: Department of Medical Genetics, Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki 305-8575, Japan. Tel.: +81 29 853 3352; fax: +81 29 853 3333.

E-mail address: hishigur@md.tsukuba.ac.jp (H. Ishiguro).

In situ hybridization analysis revealed that *PICK1* gene expression was decreased in dorsolateral prefrontal cortex (DLPFC) of patients with schizophrenia (Beneyto and Meador-Woodruff, 2006). In contrast, Dracheva et al. (2005) could not find a significant change in *PICK1* gene expression in DLPFC. Therefore, evaluation of possible effects of antipsychotics on *PICK1* expression or a transgenic animal study may be required to understand a role of the gene underlying schizophrenia.

Despite inconclusive results, genetic studies have yielded evidence that *PICK1* plays a role as a susceptibility gene in schizophrenia. *PICK1* contains 13 exons spanning approximately 18 kb and is located on chromosome 22q12.13-q13.2, where linkages for schizophrenia have been suggested (Gill et al., 1996; Kalsi et al., 1995). Further, two case-control studies of single nucleotide polymorphisms (SNPs), rs3952 and rs2076369 in *PICK1* indicated an association between *PICK1* and schizophrenia in Asian populations (Fujii et al., 2006; Hong et al., 2004). These SNPs are located in the 5' region (introns 3 and 4) of *PICK1* with 7054 bp distance in one haplotype block that covers most *PICK1* of the region both in Asian and European populations. There are less informative markers in the 3' region of *PICK1*, according to the HapMap database (<http://www.hapmap.org/>). The haplotype block contains the region encoding the functionally important PDZ domain of *PICK1*. In addition, because the haplotype block might contain the promoter region of *PICK1*, the association may explain reported differences in *PICK1* expression in schizophrenia (Beneyto and Meador-Woodruff, 2006).

The previously reported association studies composed 400 Japanese and 485 Chinese subjects. It should be noted that the minor allele of rs3952 was different one between those two populations. The NCBI database showed a similar genotype distribution of rs3952 in Japanese and European populations but provides no information for Chinese population. Larger population studies may solve a stratification problem in general Asian population. Because the most straightforward way to evaluate a genetic association is to perform analyses with sufficient statistical power to reveal the association, we attempted to confirm the association between *PICK1* and schizophrenia in a large Japanese population in the present study.

2. Materials and methods

2.1. Subjects

All subjects were of Japanese descent and were recruited from the main island of Japan. A total of 1851 unrelated patients with schizophrenia (mean

age \pm S.D., 48.9 \pm 14.5 years; 1020 men and 831 women) were diagnosed according to the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV). Control subjects were 1765 mentally healthy unrelated subjects (mean age \pm S.D., 49.0 \pm 14.3 years; 944 men and 821 women) with no self-reported family history of mental illness within second-degree relatives. The present study was approved by the Ethics Committees of the University of Tsukuba, Niigata University, Fujita Health University, Nagoya University, Okayama University and Teikyo University, and all participants provided written informed consent.

2.2. Genotyping

DNA was extracted from blood samples. We genotyped two SNP markers, rs3952 in intron 3 and rs2076269 in intron 4, in *PICK1*. The SNPs were genotyped by TaqMan assay. Predesigned TaqMan SNP genotyping assays, C_2487464_10 for rs3952 and C_2487476_10 for rs2076369, were selected from the Applied Biosystems database (<http://www.appliedbiosystems.com/>). The TaqMan reaction was performed in a final volume of 3 μ l consisting of 2.5 ng genomic DNA and Universal Master Mix (Eurogentec, Seraing, Belgium), and genotyping was performed with an ABI PRISM 7900HT Sequence Detection System (Applied Biosystems, Foster City, CA, USA).

2.3. Statistical analysis

Hardy–Weinberg equilibrium, linkage disequilibrium and allelic/haplotype frequencies, as well as an association between SNP or haplotype and schizophrenia, were evaluated with the Haploview software program (<http://www.broad.mit.edu/mpg/haploview/>). Permutation tests were also performed to calculate corrected *p*-values for multiple testing with the Haploview software. Genotype-based association was tested with Cochran–Armitage test for trends. Statistical significance was accepted at $p < 0.05$.

3. Results

Genotypic/allelic distributions of the two *PICK1* SNPs among the subject groups are shown in Tables 1 and 2. Distributions of the rs3952 and rs2076369 SNPs did not differ from Hardy–Weinberg equilibrium ($p = 0.15$ and 0.11 , respectively). No genotype/allelic association with schizophrenia was detected for rs3952 ($p = 0.54/p = 0.48$ [corrected by multiple comparison: 0.76]), and rs2076369 ($p = 0.91/p = 0.90$ [corrected by multiple comparison: 1.00]). With respect to schizophrenia subtype, no association was found between either population with organized or disorganized schizophrenia versus controls (Table 3). These SNPs were in linkage disequilibrium from each other in both groups of controls ($D' = 0.98$, $r^2 = 0.34$) and of the patients ($D' = 0.99$, $r^2 = 0.34$). Haplotype frequencies did not differ significantly between the schizophrenia and control groups (Table 4).

Table 1
Distribution of the polymorphisms in the *PICK1* gene

SNP marker	Controls			Schizophrenics			<i>p</i> -Value
	AA	AG	GG	AA	AG	GG	
rs3952	46.3% ^a	42.0% ^a	11.7% ^a	44.6% ^b	43.8% ^b	11.6% ^b	0.299
rs2076369	18.0% ^c	47.4% ^c	34.5% ^c	17.7% ^b	47.4% ^b	34.9% ^b	0.748

^a $n = 1850$, ^b $n = 1765$, ^c $n = 1851$.

Download English Version:

<https://daneshyari.com/en/article/4352571>

Download Persian Version:

<https://daneshyari.com/article/4352571>

[Daneshyari.com](https://daneshyari.com)