
Contents lists available at ScienceDirect

Ecological Modelling

journal homepage: www.elsevier.com/locate/ecolmodel



Erratum

SEVIER

Erratum to "Global sensitivity analysis of an end-to-end marine ecosystem model of the North Sea: Factors affecting the biomass of fish and benthos" [Ecol. Model. 273 (2014) 251–263]



David J. Morris*, Douglas C. Speirs, Angus I. Cameron, Michael R. Heath

Department of Mathematics and Statistics, University of Strathclyde, 26 Richmond Street, Glasgow G1 1XH, UK




Fig. 2. Results of Morris sensitivity analysis for suspension feeding benthos. Labels for the ten most influential factors are shown.

The publisher regrets that Figs. 2–7 were incorrectly reproduced. The correct figures are presented below. The publisher would like to apologise for any inconvenience caused.

http://dx.doi.org/10.1016/j.ecolmodel.2014.02.010 0304-3800/© 2014 Elsevier B.V. All rights reserved.

DOI of original article: http://dx.doi.org/10.1016/j.ecolmodel.2013.11.019.

^{*} Corresponding author. Tel.: +44 141 548 3804; fax: +44 141 548 3345.

E-mail address: david.morris@strath.ac.uk (D.J. Morris).




Fig. 3. Results of Morris sensitivity analysis for carnivorous/scavenging benthos. Labels for the ten most influential factors are shown.




Fig. 4. Results of Morris sensitivity analysis for pelagic fish larvae. Labels for the ten most influential factors are shown.

Download English Version:

https://daneshyari.com/en/article/4375979

Download Persian Version:

https://daneshyari.com/article/4375979

Daneshyari.com