



Short communication

Differentiation of two South African otter species (*Aonyx capensis* and *Lutra maculicollis*) from spraint based on partial *CytB* primer sets



M. Thabang Madisha^{a,b,*}, Damian Ponsonby^c, Ute Schwaibold^c,
Antoinette Kotzé^{a,d}, Raymond Jansen^b, Helene Brettschneider^a,
Desiré L. Dalton^{a,d}

^a National Zoological Gardens of South Africa, P.O. Box 754, Pretoria, 0001, South Africa

^b Department of Environmental, Water and Earth Sciences, Tshwane University of Technology, Private Bag X680, Pretoria, 0001, South Africa

^c School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Private Bag 3, Wits 2050, South Africa

^d Genetics Department, University of the Free State, PO Box 339, Bloemfontein 9300, South Africa

ARTICLE INFO

Article history:

Received 24 March 2015

Received in revised form 4 May 2015

Accepted 4 May 2015

Available online 11 May 2015

Keywords:

CytB

Degraded DNA

DNA primers

Otter species

Spraint

ABSTRACT

Accurate species identification based on visual cues can be challenging due to morphological similarities and the cryptic nature of certain species. Thus a more conclusive method of identification is required, namely DNA barcoding. This is the case regarding two South African otter species, Cape Clawless otter (*Aonyx capensis*) and the spotted necked otter (*Lutra maculicollis*). Due to the cryptic nature of these animals faecal samples, known as spraints, are the easiest way of confirming the presence of the animal in an area. In this study, we compared results obtained for universal and partial *CytB* primer sets on collected spraint and tissue control samples. Universal *CytB* primers revealed a low percentage of amplified otter species from faecal samples (species specific amplification success of 10.9%) whereas, the partial *CytB* primer set resulted in successful amplification of 45 out of 55 (82%) samples. We were thus able to positively differentiate between the two otter species using the partial *CytB* primer set developed in this study. The ability to accurately identify species using partial DNA will be beneficial in understanding numerous aspects of the behaviour and ecological importance of animals in their environment.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Accurate species identification based on visual cues can often be difficult due to cryptic morphological features of species. In addition, morphological identification keys often rely on a certain life stage or gender of an animal to be used correctly (Hebert et al., 2003; Bonito, 2009; Chaves et al., 2012). In the case of two cryptic, co-occurring South African otter species, namely the Cape Clawless otter (*Aonyx capensis*) and the spotted necked otter (*Lutra maculicollis*), faeces (spraint) is used as a means of determining presence, space use and diet (Arden-Clarke, 1986; Van Niekerk et al., 1998; Perrin and Carugati, 2000; Angelici et al., 2005). However, due to similar diets the spraint can most often be visually similar, making it difficult to accurately identify the species occurring in a given area. In addition, erosion of spraint through exposure to environmental

* Corresponding author at: National Zoological Gardens of South Africa, P.O. Box 754, Pretoria, 0001, South Africa. Tel.: +27 12 339 2802.
E-mail address: thabang@nzdg.ac.za (M.T. Madisha).

conditions leads to the physical breakdown of the spraint changing its appearance making it even more difficult to visually differentiate between the two species of otters based solely on the spraint (Rowe-Rowe, 1992). This can be problematic when trying to understand the movement patterns of the species, which is important for conservation purposes. This would, for example, influence the design of green corridors in urban environments, as the two species have different habitat requirements.

The South African Red Data Book (Friedmann and Daly, 2004) has listed *A. capensis* as 'least concern' due to habitat loss because of infrastructure development as a main threat. *L. maculicollis* is listed as 'near threatened' with indication of human induced habitat loss through agriculture, infrastructure development, exploitation/direct loss (persecution trapping as fish competitors and livestock predators) and accidental mortality caused by fish traps (Friedmann and Daly, 2004). *L. maculicollis* is listed as a protected species in National Environmental Management: Biodiversity Act (NEMBA, 2004), suggesting it is an indigenous species of high conservation value or national importance that requires national protection.

The development of a reliable method to differentiate these two otter species makes it easier to conduct a rough estimation of the occurrence rate of each species in an area, as well as help to better define their preferred habitats. Recent studies have shown how the use of genetic barcoding can be effectively used as a means of identifying species based on mitochondrial DNA (mtDNA) sequences (Hebert et al., 2003; Flynn et al., 2005; Sass et al., 2007; Corse et al., 2010; Chaves et al., 2012). Although this is a relatively easy and informative technique and; mtDNA is abundant in cells, most studies use non-degraded DNA such as that from tissue and blood samples. In the case of elusive animals it is often difficult to obtain tissue and blood samples, and alternative non-invasive samples can be used by collecting material left behind by the animal including hair, shed skin, egg shells and even faeces (Rowe-Rowe, 1992; Prigioni et al., 2005; Waits and Paetkau, 2005). The use of these non-invasive samples has furthered research on a variety of elusive, rare and endangered species which previously could not be studied due to various reasons such as ethics, safety concerns and even unavailability of good samples (Taberlet et al., 1997; Palomares et al., 2002; Medina-Vogel and Gonzalez-Lagos, 2008). Considering the obvious advantage of these non-invasive samples, there are also several drawbacks. Otter faecal samples do contain epithelial cells from the intestinal mucosa as excreta are pushed through the intestines the quality of the sample is low as there is a combination of the animal's DNA and DNA from prey eaten by the animal such as fish, small rodents, crabs, birds, and frogs (Dallas et al., 2000). This may be problematic as general barcoding of faecal samples used to identify unknown species tend to amplify the DNA of prey more than the target species as the primers used are usually universal and non-specific (Dallas et al., 2000). Another problem that arises from using faecal samples is the manner in which weather conditions can lead to degradation of genetic material (Hung et al., 2004). It requires technique optimisation to be able to use small amounts of DNA and target the amplification of short fragments. Despite this, degraded faecal samples have been used in previous studies to acquire genetic material (Meusnier et al., 2008). Considering this, the aim of this study is to describe the amplification of a short fragment (< 150 bp) of the mitochondrial Cytochrome b (*CytB*) gene, isolated and amplified from faecal samples, and discuss its usefulness in genetically differentiating between the Cape clawless otter (*A. capensis*) and the Spotted necked Otter (*L. maculicollis*) found in South Africa. We have selected *CytB* as it is commonly used as a species-level marker and particularly so in mammalian biosystematics (Bradley and Baker, 2001; Pfunder et al., 2004).

2. Materials and methods

This study forms part of a larger project on the ecology of otters along river systems in the Gauteng province of South Africa, which required surveying of rivers banks for collecting otter spraints to conduct a detailed dietary analysis. The Jukskei River, Klip River, Hennops River, Crocodile River, Mooi River and Pienaars River along with their associated tributaries were the focus rivers of this study (Fig. 1). During the survey of the riverbank we aimed to collect fresh spraint with jelly (gelatinous secretion of anal scent glands), however the age of the sample and amount of jelly varied. Spraint found was collected with an inverted plastic zip-lock bag so as to prevent contamination of the sample. The bag was then reverted, sealed and allocated a unique identification number, and samples were immediately stored at -10°C for further processing. The QIAGEN® QIAamp® DNA Stool Mini Kit (QIAGEN, 2010) was used for DNA extraction from otter spraint following the manufactures protocol for isolation of DNA from stool for human DNA analysis. A total of 55 samples were collected from 10 river systems.

2.1. Universal *CytB* primers

Initially, universal primers; L14724: CGAAGCTTGATATGAAAAACCATCGTTG and H15149: AAAGTCAGCCCCCTCAGAATGATATTGTCCTCA (Hsieh et al., 2003) were used to amplify approximately 400 bp of the *CytB* gene region, thereafter sequencing of this fragment was used to genetically assign collected samples to either of the two otter species found in South Africa. A total of two reference tissue samples, one each from the otter species; a road killed Cape clawless otter (*A. capensis*) from Knysna and spotted necked otter (*L. maculicollis*) found drowned on Grahamstown were used as controls.

Amplicons for sequencing were generated using the Dreamtaq PCR mastermix (Thermo Scientific, USA) at a final reaction volume of 25 μL . The reaction consisted of 12.5 μL of Dreamtaq mastermix (2x), 9.5 μL double distilled water (ddH_2O), 10 pmol forward and reverse primers (Integrated DNA technologies, Coralville, IA, USA) and 1 μL (20 ng) of template DNA. The PCR conditions for the universal primers were as follows: one cycle of denaturation at 94°C for 5 min (min) followed by

Download English Version:

<https://daneshyari.com/en/article/4379477>

Download Persian Version:

<https://daneshyari.com/article/4379477>

[Daneshyari.com](https://daneshyari.com)