



2nd International Symposium on Aquatic Products Processing and Health  
ISAPPROSH 2015

## Microalgae *Dunaliella salina* (Teodoresco, 1905) Growth using the LED Light (Light Limiting Dioda) and Different Media

Shifa Helena<sup>a\*</sup>, Muhammad Zainuri<sup>b</sup>, Jusup Suprijanto<sup>b</sup>

<sup>a</sup>Graduate Student of Marine Science, University of Diponegoro, Tembalang, Semarang, 50275 Indonesia

<sup>b</sup>Faculty Fisheries and Marine Sciences, University of Diponegoro, Tembalang Semarang, 50275 Indonesia.

### Abstract

The purpose of this research was to analyze the growth of *Dunaliella salina* (Teodoresco, 1905) with different media types and based light source of LED as well as obtaining the optimum media and light source that generate the highest growth of *D. salina*. The data was analyzed using ANOVA. The material used was *D. salina* which cultured on 3 L bottles, with three treatments and three replications. The treatments given were respectively Walne Jepara, Walne Lampung, and Za added with NPK, which were then illuminated by using red LED and blue LED. The density of *D. salina* cells were determined using a light microscope with a Neubauer chamber. The highest result of *D. salina* density is  $8.504 \times 10^4 \text{ cell} \cdot \text{mL}^{-1}$  which produced by the red LED treatment in the media of Walne Jepara. The results of blue LED treatment showed that *D. salina* density was  $5.768 \times 10^4 \text{ cell} \cdot \text{mL}^{-1}$  with Walne Jepara.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the science and editorial board of ISAPPROSH 2015

**Keywords:** Culture growth; *Dunaliella salina* (Teodoresco, 1905); LED; Walne Pro-analyze; Walne technic

### 1. Introduction

*Dunaliella salina* (Teodoresco, 1905) is a microalgae and their cell was highly responsive to osmotic changes permitting arapid changes in the cell shape. Reproduction of *D. salina* is asexual flagellated cells that may produce

\* Corresponding author. Tel.: +62 857 4220 2010.  
E-mail address: shifahelena31@gmail.com

thick walled cysts, which allow survivorship under an environmental stress. The accumulation of carotenoids in *D. salina* allowed the intensity of light as the main stimulus for  $\beta$ -carotene references by Lamers et al. (2010). Utilization of *D. salina* was quite diverse ranging from health food as it has been marketed in developed countries (Chang et.al., 2011). *D. salina* in response to stressful conditions such as high light intensity, temperature and salt concentration causes the buildup of various secondary metabolites such as lycopene,  $\beta$ -carotene, lutein and zeaxanthin (Lamers et.al., 2012). Microalgae are expensive to produce, although many efforts are under way addressed to achieve cost-efficient modes for mass cultivation of these organisms. Different systems have been designed for the growth and handling of microalgae on a large scale references by Zainuri et.al., (2008). The objective of this research was to analyze the growth of *D. Salina* at different media types based on light and source of LED (light-emitting dioda), as well as obtaining the optimum media and light source that generate the highest growth of *D. salina*.

Medium Walne is a basic medium that is often used in the cultivation of *D. salina*. However, this basic medium has the shortcomings include the nutrient content that is still able to accelerate the growth of *D. salina*. Pro-analyze walne Jepara of fertilizer is very expensive so if they are used for mass scale by the public will be detrimental to farmers (Andersen, 2005). Therefore, the aim of the research was to looking for an alternative nutrient thus saving fertilizer in cultivation of *D. salina*. The first treatment that is done to meet the nutrient requirement is the provision of ZA fertilizer containing ammonium sulfate or  $(\text{NH}_4)_2\text{SO}_4$  and non-technical walne.

The microalgae cultivation facilities typically use sea water enriched with nutrients, especially carbon, nitrate and phosphate (Fu et.al., 2012). *D. salina* need a complete nutrient composition can affect the concentration of biomass production and nutrient content of microalgae. The research to increase the growth of microalgae is to control the content of both macro and micro nutrients in the cultivation environment (Harisson, 2005).

## 2. Material and methods

The material used was *D. salina* from the brackish water aquaculture in Jepara in Central Java, Indonesia and then it cultured in 3 L bottles, with three treatments and three replications. The method used was Completely Randomized Design (CRD) in time. The purpose of the Completely Randomized Design in time was to determine the effect of treatment on the provision in improving water quality (Creswell, 2010).

The treatment given were as follows: type media the first was Walne pro-analyze from brackish water from Jepara, Central Java, Indonesia, second Walne techniques from brackish water from Lampung in North Sumatra Indonesia and the last was media ZA (ammonium sulfate) added NPK (Nitrogen Phosphorous Potassium). The factor two was were then illuminated by using red LED and blue LED.

Cells were determined by direct counting, using a light microscope (magnification 40 $\times$ ) with a Neubauer chamber basic Hemocytometer Usage. The Neubauer chamber was a thick crystal slide with the size of a glass slide (30 mm  $\times$  70 mm and 4 mm thickness).

Culture of *D. salina* can be done when it reaches the peak of the population. Population peaks can be seen from the color change in the culture medium and the total population based on growth patterns. According to some sources, harvesting was done at the time of *D. salina* was located at the end of the exponential phase, approximately on 7 d. The culture has reached a population peak was precipitated first by lethal aeration. Then the solids obtained by using a centrifuge, and then weighed to determine the biomass produced. *D. salina* then has weighed, dried at cool room temperature, around 21  $^{\circ}\text{C}$  for about 4 d, from each sample.

The research was carried out in the Laboratory of Marine Biology, University of Diponegoro, Semarang, Indonesia. The data were analyzed using SPSS factorial analysis of variance (ANOVA two ways).

## 3. Result and discussions

The results showed that on difference media types and based light source given different real influence on the pattern of growth of *D. salina*. The result of growth of *D. salina* were highest of  $850.4 \times 10^{-4} \text{ cell} \cdot \text{mL}^{-1}$ , in using a media Walne Jepara and red LED. The highest results of blue LED was of  $576.8 \times 10^{-4} \text{ cell} \cdot \text{mL}^{-1}$ , Walne Jepara.

Test statistics on cell density data *D. salina* showed that the normal kinds of data spread, homogenous and are additive. Data source of light and the media is very significant or significant to the growth of *D. salina*. The highest

Download English Version:

<https://daneshyari.com/en/article/4383613>

Download Persian Version:

<https://daneshyari.com/article/4383613>

[Daneshyari.com](https://daneshyari.com)