



Short communication

Increased density facilitates plant acclimation to drought stress in the emergent macrophyte *Polygonum hydropiper*

Feng Li^{a,b,c}, Yonghong Xie^{a,b,*}, Chen Zhang^{a,b}, Xinsheng Chen^{a,b}, Bingbing Song^d,
Youzhi Li^{a,b}, Yue Tang^{a,b}, Jiayu Hu^{a,b,e}

^a Key Laboratory of Agro-ecological Processes in Subtropical Region, The Chinese Academy of Sciences, Hunan 410125, China

^b Dongting Lake Station for Wetland Ecosystem Research, Institute of Subtropical Agriculture, Changsha 410125, China

^c State Key Laboratory of Lake Science and Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Science, Nanjing 210008, China

^d Hunan Province Environmental Monitoring Center, Changsha 410019, China

^e University of the Chinese Academy of Sciences, Beijing 100049, China

ARTICLE INFO

Article history:

Received 9 December 2013

Received in revised form 26 May 2014

Accepted 12 July 2014

Available online 3 August 2014

Keywords:

Drought stress

Density

Polygonum hydropiper

Electrolyte leakage

MDA content

ABSTRACT

Wetland plants frequently experience drought events due to large water level fluctuations; drought might present a greater limitation to plant survival than flooding in many wetlands. High plant density can increase the relative humidity, which might facilitate the acclimation of plants to drought stress. However, studies on the effects of plant density on acclimation to drought are scarce. This study aimed to elucidate the influence of density on plant response to drought stress by investigating morphological and physiological characteristics of *Polygonum hydropiper* L. var. *flaccidum*, one of the dominant species in the Dongting Lake wetland, China. Experimental greenhouse treatments, including three density (16, 144, and 400 plants m⁻²) and three water level (0, –20, and –40 cm relative to the soil surface) treatments, were performed in a factorial design. Soil water content increased with increasing water level and plant density. In the 0- and –20 cm water level treatments, biomass accumulation per plant decreased significantly with increasing density. However, in the –40 cm treatment, biomass accumulation did not change among the three plant densities. Higher density and water level led to lower leaf mass fraction. Both electrolyte leakage and malondialdehyde (MDA) content increased along with decrease in water level. Moreover, they showed no change with plant density in the 0- and –20 cm water level treatments, but decreased with increased density in the –40 cm treatment. Soluble sugar content increased with decreasing water level and increasing density, whereas starch content decreased significantly with decreasing water level. These data indicate that a higher density facilitates the acclimation of *P. hydropiper* to drought stress. Moreover, our results might reveal the important role of density in the restoration of *P. hydropiper* communities.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Many natural wetlands are maintained primarily by large seasonal fluctuations in water level, such as complete inundation during wet parts of the year interspersed with prolonged dry periods (Parolin et al., 2010; Li et al., 2013). The tolerance or sensitivity of wetland plants to these hydrologic extremes is important to determine their survival and distribution (Pagter et al., 2005). The influence of flooding on the growth, distribution, and

composition of wetland macrophytes has been studied in detail (Luo et al., 2010; Li et al., 2011). However, the role of dry periods has been investigated less thoroughly, although this factor might have a greater impact on plant survival than flooding in many wetlands (Lopez and Kursar, 2007).

Drought adversely influences the growth and metabolism of wetland plants by stimulating increased production of superoxides such as malondialdehyde (MDA) or hydrogen peroxide (Luo et al., 2008; Zhao et al., 2008), and by leading to a decline in photosynthetic activity (Pagter et al., 2005). Generally, plants can tolerate or avoid drought stress through morphological, physiological, and phenological mechanisms such as adjustment of biomass allocation; modification of root length; adjustment of osmotic potential; and enhancement of MDA, proline, soluble sugar, and antioxidative enzyme contents (Bartels and Sunkar, 2005; Xie et al., 2008; Li et al.,

* Corresponding author at: Dongting Lake Station for Wetland Ecosystem Research, Institute of Subtropical Agriculture, Changsha 410125, China.
Tel.: +86 731 84615203.

E-mail address: yonghongxie@163.com (Y. Xie).

2013). These adjustments are beneficial for plants to improve their water-absorbing capacity and to resist desiccation or alleviate lipid peroxidation (Bartels and Sunkar, 2005). Moreover, these adjustments depend on the drought severity and duration, plant genotype and developmental stage, and environmental factors (Pagter et al., 2005).

The role of density in maintaining vegetation structure and population dynamics has been extensively investigated (Jiang et al., 2008; Li et al., 2009; Ala et al., 2014). Generally, plant growth can be inhibited in densely occupied habitats due to the competition for resources such as light and nutrients (Shilo-Volin et al., 2005). *Lemna minor* and *Spartina anglica* were shown to exhibit growth limitation or death at densities above 400–700 plants m^{-2} and 81 plants m^{-2} , respectively (Driever et al., 2005; Li et al., 2009). However, a dense canopy also has positive effects on plant growth, particularly under severe environmental conditions (Franks and Peterson, 2003; Luo et al., 2010). When treated with 40 cm water level, performances of three marsh plants were considerably better in high density treatments, since high plant density produced a relatively oxygen-rich environment (Luo et al., 2010). Physical and physiological effects of wind were ameliorated by higher density in *Sinapis alba* (Retuerto et al., 1996). High density might also facilitate plant acclimation to drought stress, since relative humidity and leaf wetness duration could be increased by reduced air movement and light penetration (Tu, 1997; Asghari et al., 2009). However, this positive effect requires further study.

Dongting Lake (28°30′–30°20′N, 111°40′–113°10′E) is the second-largest and the most typical river-connected lake in China. The wetlands in this lake are characterized by large seasonal fluctuations in water level; they are completely flooded during May to October and remain dry during November to April. In recent years, drought episodes have occurred frequently in the Dongting Lake ecosystem, leading to significant changes in the structure of plant communities (Li et al., 2013). In particular, during our annual field investigation, we (the staff of Dongting Lake Station for Wetland Ecosystem Research, the Chinese Academy of Sciences) found that *Polygonum hydropiper* community had degraded seriously in areas such as the lake shores of Chanpanzhou and Beizhouzi.

This study aimed to determine whether increased plant density could facilitate plant acclimation to drought stress. For this, a dominant wetland plant of Dongting Lake, *P. hydropiper*, was grown at three densities under three water levels, and morphological (biomass accumulation and allocation) and physiological (electrolyte leakage and contents of MDA, soluble sugar, and starch) characteristics were investigated to test the following hypotheses: (1) both drought and high-density treatment will lead to lower biomass accumulation of *P. hydropiper*, and the influence of drought will be stronger in the low-density than in the high-density treatments; (2) drought will lead to a higher belowground mass fraction but lower leaf and stem mass fraction for *P. hydropiper*, and these changes will be stronger in the low-density than in the high-density treatments; (3) electrolyte leakage and MDA and soluble sugar contents of *P. hydropiper* will increase, but starch content will decrease in the drought treatment, and there will be less evidence of these changes in the high-density than in the low-density treatments.

2. Materials and methods

2.1. Plant materials

P. hydropiper can form large patches of monodominant communities or occur as a codominant species with *Carex* species in Dongting Lake (Peng et al., 1984). This species normally flowers and fruits from April to May before the period of flooding, and the

recruitment of this species mainly occurs by producing vegetative ramets from rhizomes (Chen et al., 2011).

Ramets of *P. hydropiper* were collected in March 2012 from a monodominant stand in Chunfeng Village (29°13′49.72″N, 113°02′32.79″E), East Dongting Lake wetlands. The vegetation was cut into small blocks (20 × 20 cm) and transported to a greenhouse at the Institute of Subtropical Agriculture, the Chinese Academy of Sciences. New ramets were germinated by placing plant fragments with roots into plastic buckets containing 15 cm soil (1.87% organic matter, 28.5 $\mu g\ g^{-1}$ total N, and 7.83 $\mu g\ g^{-1}$ total P) collected from the *P. hydropiper* community. The temperature was controlled at $25 \pm 2^\circ C$ during the day and $17 \pm 2^\circ C$ at night. Light was provided using 400 W SON-T ARGO sodium lamps (Phillips, Guildford, UK) at a photon flux density of 600 $\mu mol\ m^{-2}\ s^{-1}$ (physiologically active radiation) for a 14-h photoperiod. Tap water (containing 0.511 $\mu g\ L^{-1}$ NH_4^+-N , 1.760 $\mu g\ L^{-1}$ $NO_3^- -N$, and 0.527 $\mu g\ L^{-1}$ $PO_4^{3+} -P$, pH = 7.2) was supplied daily.

2.2. Experimental set-up

The experiment included three water levels (0, –20, and –40 cm relative to the soil surface) with three densities (low, 16 plants m^{-2} ; medium, 144 plants m^{-2} ; and high, 400 plants m^{-2} ; field density ranged from 25 to 390 plants m^{-2}) in a factorial design with three replicates. In all, the experiment included nine treatments: three single-individual treatments (one plant per water level) and six multi-individual treatments (9 or 25 plants per water level, with the individual in the center designated as the target plant). A total of 630 similar-sized ramets (6–7 leaves, approximately 23 cm high) were cut from plant cultures on April 15, 2012. These ramets were planted in polyvinyl chloride tubes (45 cm high × 30 cm diameter) filled with 45 cm of the same soil used for plant incubation. Six tubes (two tubes per density treatment) were placed in each of nine larger plastic basins (98 × 76 × 68 cm) to control the water level.

After planting, the water level in all the basins was maintained at 45 cm depth for 1 week, after which it was adjusted to the required level. Water depth was adjusted weekly during the experimental period, and tap water was supplied as necessary. The entire volume of water was replaced every 2 weeks to prevent algal growth, and the water content of each tube was measured monthly by using a soil water analyzer (TZS-2X; Zhengjiang).

2.2.1 Harvest and measurement of physiological indexes
The plants were harvested after 6 months. First, fresh leaf samples were collected for the analysis of physiological traits, and roots were carefully removed from the soil and cleaned using tap water. Plants were then divided into leaves, shoots, and belowground parts (rhizomes and roots), oven dried at 80 °C for 48 h, and weighed. Leaf mass removed for analysis of physiological traits was added to the corresponding plant biomass by converting fresh weight to dry weight. Leaf, stem and belowground mass fractions were determined as the ratios of leaf mass, stem mass, and belowground mass (rhizome and root) to total biomass, respectively.

Electrolyte leakage was determined as described by Dionisio-Sese and Tobita (1998). The content of MDA was measured using the method described by Duan et al. (2005). Dry leaf samples were used for soluble sugar and starch content analysis according to the methodology described by Yemm and Willis (1954).

2.3. Data analysis

For each basin, the mean values of two tubes per density treatment were used for the data analysis. Repeated-measures analysis of variance (ANOVA), with water level and density as dependent factors and time as a repeated-measures factor, was used to evaluate treatment effects on soil water content. Two-way ANOVA,

Download English Version:

<https://daneshyari.com/en/article/4389138>

Download Persian Version:

<https://daneshyari.com/article/4389138>

[Daneshyari.com](https://daneshyari.com)