



Short communication

Amount vs. temporal pattern: On the importance of intra-annual climatic conditions on tree growth in a dry environment

Michael Dorman^{a,*}, Avi Perevolotsky^b, Dimitrios Sarris^{c,d,e}, Tal Svoray^a^a Department of Geography and Environmental Development, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel^b Department of Natural Resources, Agricultural Research Organization, Volcani Ctr, Bet Dagan 50250, Israel^c Faculty of Pure & Applied Sciences, Open University of Cyprus, 2252 Latsia, Nicosia, Cyprus^d Department of Biological Sciences, University of Cyprus, 1678 Nicosia, Cyprus^e Division of Plant Biology, Department of Biology, University of Patras, 265 00 Patras, Greece

ARTICLE INFO

Article history:

Received 23 December 2014

Received in revised form

21 February 2015

Accepted 2 March 2015

Available online 10 March 2015

Keywords:

Aleppo pine

Climate change

Dendrochronology

Pinus halepensis

Semi-arid

Tree rings

ABSTRACT

Forests in semi-arid regions are at particular risk for climate change impacts. Although it has been recently acknowledged that vulnerability to climate change depends on changes in climatic variability and the occurrence of extreme events, and not just the mean climatic conditions, the relative importance of such effects remains largely unexamined. In the present study we investigated the effects of intra-annual rainfall distribution characteristics, as opposed to total rainfall amount, on tree growth. More specifically, proportion of large rain events and dry season length – two climatic characteristics considered key to the survival of planted *Pinus halepensis* forests in a semi-arid region – were evaluated based on a tree-rings dataset. Dry season length did not have a significant effect on growth, highlighting the high resilience of this species when facing harsh climatic conditions. Proportion of large rain events had a positive effect on growth under dry conditions, as expected. The magnitude of this effect was relatively small, compared to that of total rainfall amount. Nevertheless, an increase in the proportion of large rain events as a result of climate change may potentially balance the decline in its total amount, in terms of trees growth rate, to an extent quantifiable using our statistical model predictions.

© 2015 Elsevier Ltd. All rights reserved.

Forests in semi-arid regions of mid-latitudes are an example of an ecosystem where negative effects of climate change are already underway (Allen et al. 2010; Steinkamp and Hickler, 2015), with widespread mortality events reducing forest cover and increasing the potential for soil erosion. It has been recently acknowledged that vulnerability of plants to climate change will largely depend on changes in climatic variability and the occurrence of extreme events, and not just the mean climatic conditions (Smith, 2011; Reyner et al. 2013). For example, precipitation events, as well as drought, are predicted to become more extreme, potentially changing soil water distribution across different depths in a way which benefits deeper-rooted woody plants (Kulmatiski and Beard, 2013). Increased climatic variability, however, may also translate to a higher probability of long periods of drought, which may negatively affect perennial species even if the total amount of precipitation is unchanged. For instance, in highly seasonal semi-arid

climates tree mortality can occur when dry season length (DSL) surpasses the physiological tolerance threshold of the given tree species (Breshears et al. 2009; Klein et al. 2014). Nevertheless, the consequences of changes in the hydrological regime to vegetation productivity have mainly been limited to theoretical examination, at least for woody plants (Anderegg et al. 2013).

Dendrochronology can provide a retrospective characterization of year-to-year growth variation of trees, making it the most suitable tool for studying climate-growth relationships. Taking into account concurrent climatic conditions, dendrochronological studies have long ago established the link between tree growth processes and limiting climatic factors, one of them being water availability, with varying degree of sensitivity according to the timing of the climatic stress. A recently adopted practice in tree rings research is also taking into account the duration, or time-scale, over which climatic state is observed (Vicente-Serrano et al. 2013). Nevertheless, only a monthly resolution is generally considered (e.g. annual rainfall sum, last 3-months mean temperature in January, etc.). An alternative way of characterizing water availability conditions in a more comprehensive manner involves

* Corresponding author.

E-mail address: michael.dorman@mail.huji.ac.il (M. Dorman).

taking one step further and constructing soil water content (SWC) models, fed with detailed climatic data regarding precipitation and potential evapotranspiration (PET). SWC prediction, however, is frequently also based on monthly data (e.g. Lebourgeois et al. 2013) or aggregated to such prior to the contrasting with tree growth (e.g. Michelot et al. 2012). Moreover, SWC models inevitably rely on numerous assumptions, such as soil water holding capacity and maximal transpiration rate, which are difficult to verify, especially for trees (Granier et al. 1999). Thus, climatic characteristics that address higher-than-monthly resolution or characterize timing rather than amount, such as the proportion of large rain events (LRE) and DSL (respectively) largely remain unexamined.

Pinus halepensis is a relatively drought-resistant pine species naturally found in the western Mediterranean region, with a few isolated populations in the eastern Mediterranean. It was also extensively planted, including in semi-arid environments where climate is drier than encountered within the natural distribution range (de Luis et al. 2013). One such region is southern Israel, where planted forests of the species grow under average annual rainfall amounts (P) of 250–350 mm (Schiller and Atzmon, 2009). Recent dendrochronological studies conducted in the region implied that P is the principal climatic variable affecting performance of *P. halepensis*, rather than rainfall amounts over shorter (monthly) or longer (multi-annual) time scales, or other factors such as temperature (Maseyk et al. 2011; Dorman et al. in press-a). Because P is much lower than PET, lack of deep ground infiltration and formation of deep roots may limit the influence of multiple years' rainfall on tree growth. On the other hand, due to the relative insignificance of other limiting factors, all rainfall is potentially utilized for growth, and thus rainfall in no particular month or season is more important than the whole annual sum (Sarris et al. 2007; Dorman et al. in press-a).

A detailed hydrological investigation of the planted forests' ecosystem conducted in the region (Raz-Yaseef et al. 2010, 2012) suggested that forest productivity was more strongly dependent on the proportion of rainfall occurring as part of intensive (>30 mm) storms (P_{30}/P ; see below) leading to relatively deep infiltration, than on P. It was also proposed that DSL plays an important role in tree survival – since it determines the amount of time trees are exposed to a zero transpirable SWC regime – given the initial water availability attained by the end of the wet season and the time span until the following wet season begins (Klein et al. 2014). However, both above-mentioned hypotheses are based on a limited time frame (4–5 years), thus the significance and relative importance of the suggested effects could not be examined.

The purpose of the present study is to investigate the relative importance of P_{30}/P and DSL, as opposed to P, for *P. halepensis* tree growth in semi-arid regions. This is achieved by combining (1) simple climatic indices (addressing specific timing- and intensity-related climatic characteristics), the biological importance of which has been previously recognized, with (2) a comprehensive tree-rings dataset spanning a time-frame of 30 years. The results are also ultimately intended to aid in the prediction of forest response to future climatic changes, by highlighting the relative importance of intra-annual rainfall distribution as opposed to its total amount.

The studied area (see Fig. 1 in Dorman et al. in press-b) included planted monoculture *P. halepensis* stands in Lahav and Dvira forests in southern Israel. The climate in this area is semi-arid (average annual rainfall of ~300 mm), and highly seasonal (5 consecutive months with <2.5 mm rainfall depth). Three sites were randomly selected within the area of Lahav and Dvira forests for dendrochronological sampling. All sites were located on south-facing slopes (to test responses of trees to elevated drought stress as such slopes are drier due to higher solar radiation load) of similar

lithology, and were of similar age (43–45 years old in 2012).

Dendrochronological sampling was conducted during autumn 2011–spring 2012. In each of the three sites, 30 living unsupported, i.e., not overshadowed by their neighbors, trees were randomly selected (Fig. A.1). Two wood cores were extracted from opposite sides of each tree at breast height using an increment borer. Cores were sanded using increasingly fine sanding paper until tree rings were clearly visible under a binocular microscope. Tree Ring Width (TRW) was measured to an accuracy of 0.01 mm using a LINTAB 6 measuring device (Rinntech, Heidelberg, Germany). Five trees with damaged cores or un-datable missing rings were removed from the analysis. Finally, TRW values were converted to Basal Area Increment (BAI) values (Biondi and Qeadan, 2008), and average BAI chronologies were calculated per tree. Structural characteristics of the sites (tree dimensions, density and mortality) are provided in Dorman et al. (in press-b). Records from the last 30 years of tree growth (1983–2012), when growth patterns had already stabilized after the establishment phase (Fig. A.2), were used for analysis of climatic effects on forest growth in each region. The final dataset included 2549 BAI records, from 85 trees over 30 years.

To characterize climatic conditions in the studied area, daily precipitation records for the period 1953–2012 were obtained from the standard meteorological station “Lahav” bordering Lahav forest (31.38N/34.87E). Climatic conditions were characterized using three variables: P, P_{30}/P and DSL. P for the year t was defined as the rainfall sum over the respective hydrological year (1st of August in year $t-1$ until 31st of July in year t). LRE were defined as consecutive runs of rainy days with the total amount of rainfall >30 mm. Annual sum of rainfall during LRE (P_{30}) was divided by P to obtain P_{30}/P (Fig. 1). The onset and the end of the wet season were considered the days when 10% and 90% of the annual rainfall amount have been accumulated, respectively. Accordingly, the DSL for year t has been defined as the period of time between the end of the wet season in year $t-1$ and the onset of the wet season of year $t-1$ (Fig. 1).

A Linear Mixed-Effects Model has been constructed to consider the effects of fixed (P, P_{30}/P and DSL) and random (site and tree-within-site) effects on BAI. The most parsimonious model has been selected according to the Akaike Information Criterion (AIC). For easier interpretation of the model (Table 1), predicted BAI was calculated for an array of climatic settings encompassing the range of observed condition in 1983–2012 (Fig. 2). Model fitting was done in R (R Core Team, 2014), using package nlme (Pinheiro et al. 2014).

During the whole period of climatic data availability (1953–2012), average P in the studied region was 292 mm, about half of which fell during LRE (average $P_{30}/P = 0.53$). The average wet season started on November 29th and ended on March 17th. DSL varied between 207 and 357 days, with the average being 258 days. According to linear regression, there was no significant change over time ($p > 0.1$) in any of the three variables (P, P_{30}/P , DSL).

The most parsimonious model of tree growth included only P, P_{30}/P and their interaction (Table 1). This means that DSL had no substantial effect on growth. BAI was strongly and positively affected by P (Fig. 2). The effect of P_{30}/P was more complicated, due to the $P \times P_{30}/P$ interaction. P_{30}/P had a positive effect on growth under relatively dry conditions, and a negative effect under relatively wet conditions (Fig. 2). For example, under average conditions (P = 300 mm), predicted BAI was 6.7% higher under an intensive rainfall regime compared to a less intensive one (P_{30}/P increase from 0.44 to 0.65; the 25% and 75% quantiles of the variation observed during 1983–2012). Under drier conditions (P = 200 mm) the beneficial effect of LRE was much more substantial (BAI increase by 22.7%). Under more humid conditions (P = 400 mm), however, the same increase in LRE led to –1.4%

Download English Version:

<https://daneshyari.com/en/article/4392850>

Download Persian Version:

<https://daneshyari.com/article/4392850>

[Daneshyari.com](https://daneshyari.com)