

Short communication

Seed harvester ants (*Pogonomyrmex rugosus*) as “pulse” predators

W.G. Whitford*, E. Jackson

USDA-ARS Jornada Experimental Range, MSC 3JER, New Mexico State University,
Las Cruces, NM 88003, USA

Received 14 August 2006; received in revised form 2 January 2007; accepted 10 January 2007
Available online 6 March 2007

Abstract

Seed harvesting ants, *Pogonomyrmex rugosus*, collected grass cicadas at a high rate ($>5 \text{ min}^{-1}$ taken into the nest) at one location where cicada emergence exceeded 3 m^{-2} . Dry conditions in the winter-spring resulted in no annual plants in the northern Chihuahuan Desert. *P. rugosus* colonies were inactive in areas where grass cicada emergence was less than 1 m^{-2} . *P. rugosus* initiate intense predatory activity in response to pulse of large numbers of prey. This study demonstrates that predatory behavior of seed harvesting ants is not limited to incidental encounters with prey during seed harvesting activities. Pulse predation demonstrates the importance of protein to seed harvester ant colonies.

© 2007 Elsevier Ltd. All rights reserved.

Keywords: *Beameria vanosa*; Foraging ecology; Grass cicadas

Patterns and processes of arid and semi-arid ecosystems are characterized by temporal and spatial heterogeneity. The temporal dynamics of desert ecosystem processes have been described by the pulse reserve paradigm (Noy-Meir, 1973). In the pulse-reserve conceptual model, rainfall stimulates a pulse of activity (new growth, reproduction etc.) drawing on reserve resources. At the end of the pulse, reserves are replenished (production of seeds, energy storage, etc.). Thus arid and semi-arid ecosystems are typically viewed as “pulsed” systems in which all ecosystem processes proceed in short burst of intense biological activity initiated by water availability (Noy-Meir, 1979/80).

*Corresponding author. Tel.: +1 505 521 1358.

E-mail address: wawhitfo@nmsu.edu (W.G. Whitford).

Seed harvester ants, *Pogonomyrmex rugosus*, forage intensively on the seeds of herbaceous annuals and annual grasses. Intense foraging by *P. rugosus* is limited to time periods when seeds are available on the soil surface (Whitford, 1976, 1978). Although many studies document the dependence of species of *Pogonomyrmex* on seeds (MacMahon et al., 2000) some of the data on foraging documents predatory behavior of these ants with termites as the most frequent prey returned to the colonies of harvester ants (Whitford, 1978). These studies suggest that predatory behavior of harvester ants is an opportunistic activity that occurs when ants encounter suitable insect prey while hunting seeds. Here we report a brief study that demonstrates that seed harvester ants exhibit a “pulse” of predation in response to a short duration episodic event (the emergence of a large number of prey). Our study was limited to one area where the numbers of emerging grass cicada were apparently sufficiently abundant to stimulate intensive foraging on emergent cicadas by harvester ants (*Pogonomyrmex rugosus*). Additional sites were not included because harvester ants were not actively foraging in other areas where there were low numbers of emerging cicadas.

Not all desert organisms exhibit a rainfall “pulse” phenology. Some desert species follow seasonal patterns of development that are independent from the timing of rain events. Cicadas are an example of rain event independent seasonal species. In the Chihuahuan Desert, the emergence of cicadas is synchronous and seasonally predictable (mid-May to mid-June). The synchronized, short period of emergence of grass cicadas is important for some fraction of the cicadas to avoid predation. Chihuahuan Desert grass cicadas (*Beameria vanosa* (Uhler)) inhabit fine textured soils in grasslands dominated by burrograss (*Scleropogon brevifolia*), alkali sacaton (*Sporobolus airoides*) tobosa grass (*Pleuraphis mutica*) and grama grasses (*Bouteloua* spp.). In May 2000, surveys of 10 m radius circular plots in these habitats on the USDA-ARS Jornada Experimental Range (JER; 50 km NNE of Las Cruces, Dona Ana Co., NM) yielded less than 1 grass cicada emergence hole m^{-2} . The harvester ant (*P. rugosus*) colonies in these habitats on the Jornada were not active during the period that grass cicadas were emerging. The absence of foraging activity by harvester ants was attributed to the dry conditions of the preceding 6 months (total rainfall—8.38 mm) and the absence of seed producing spring annual plants.

In late May, while visiting potential study sites at the Sevilleta Long-Term Ecological Research Site (approximately 25 km N of Socorro, NM), we found an area with numerous active colonies of harvester ants (*P. rugosus*). *P. rugosus* foragers were returning to the nests with grass cicadas. The active harvester ant colonies were on an ecotone that was a mixture of blue grama (*Bouteloua gracilis*)—black grama (*Bouteloua eriopoda*) grassland on fine textured soil on a low slope bajada of the Pino Altos mountains (N 34°0.209', W 106°37.656'). This area had burned in June 1998 by a lightning set fire. Grass cover in May 2000 was close to pre-burn values.

We recorded the rate of grass cicada harvest by *P. rugosus* by counting the number of foragers entering a nest with cicadas in their mandibles within a 1-min period. Counts were made at 12 nests within a 5-ha area beginning at 1630 h. Repeat counts were made over a 1.5-h period. The abundance of grass cicadas was estimated by counting emergence holes in seven, 4 m radius circular plots centered on points located by a surveyors pin tossed at random behind the observer. The following day, counts of active foragers with and without cicadas were made at 12 nests beginning at 0900 h; ending at 1130 h.

The average density of grass cicada emergence holes was $3.1 \pm 2.1 m^{-2}$ (range 0.9–6.7 m^{-2}). Emergence holes were contagiously distributed with highest concentrations

Download English Version:

<https://daneshyari.com/en/article/4394534>

Download Persian Version:

<https://daneshyari.com/article/4394534>

[Daneshyari.com](https://daneshyari.com)