

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Journal of Asia-Pacific Biodiversity

journal homepage: <http://www.elsevier.com/locate/japb>Journal of
Asia-Pacific
Biodiversity

Original article

Variation of avifaunal diversity in relation to land-use modifications around a tropical estuary, the Negombo estuary in Sri Lanka

Manjari Bhagya Jayathilake, Withanage Upali Chandrasekara*

Department of Zoology and Environmental Management, Faculty of Science, University of Kelaniya, Kelaniya 11600, Sri Lanka

ARTICLE INFO

Article history:

Received 21 November 2014

Received in revised form

30 January 2015

Accepted 3 February 2015

Available online 11 February 2015

Keywords:

Anthropogenic disturbances

Avifauna

Estuary

House crow

Land-use modifications

ABSTRACT

We assessed variation in avifaunal diversity at some selected habitats around the Negombo estuary in Sri Lanka in relation to land-use modifications. During the study period, we observed 48 bird species of which 47 species are residents to Sri Lanka. The avian species richness, evenness, and heterogeneity were found to be the highest at undisturbed habitats. Further, these diversity measures were negatively correlated with the intensity of anthropogenic land-use activities. Total abundance of birds increased at highly disturbed habitats due to the presence of the house crow, as it is the most abundant of all birds observed at these habitats. This study highlights the need for habitat management around estuaries, giving due consideration to existing ecological theories to conserve avifaunal diversity. It also highlights the negative impacts of the house crow on diversity of other resident avian fauna in these habitats.

Copyright © 2015, National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA). Production and hosting by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Coastal ecosystems, including estuaries and associated habitats, have been increasingly altered and developed for human settlement (DeLuca et al 2008) and for commercial and recreational use. These anthropogenic disturbances have severely affected the long-term viability, health (Kennish 2002), and the biota including birds inhabiting these sensitive habitats.

Due to the high productivity, estuaries and coastal regions around the world have been the focal points of human settlement and marine resource use, and such use has strong negative impacts on plant and animal communities (Hilbert 2006; Lotze et al 2006). In many parts of the world human activities around estuaries significantly affect bird communities, their behavior, and existence. For example, DeLuca et al (2008) found that coastal urbanization, even at low levels, significantly affects the integrity of aquatic bird communities in the Chesapeake Bay, USA. Similarly, the abundance and richness of shorebirds and other aquatic birds were low in Southern California, where human activities were high (Lafferty et al 2013). It has also been found that human influences at intertidal mud flats of Queule River estuary in Chile have affected the

distribution of both migratory and resident birds there (Suazo et al 2012). However, prominent features or characteristics of urbanization and human habitations around estuarine land-use patterns that govern the avifaunal distributions are yet to be assessed.

Sri Lanka is a small island in the Indian Ocean and together with Western Ghats of India, is recognized as one of the 34 biodiversity hotspots of the world (Gunatilleke et al 2008). In spite of the small size, Sri Lanka is known to harbor 426 avian species including residents and winter migrants (Harrison 2011). The island has its unique specialties as well, with more than 20 species and over 70 subspecies being endemic to Sri Lanka (Harrison 2011). A few studies that assess estuarine avian communities have already been carried out in Sri Lanka. For example, Bellio and Kingsford (2013) studied the alteration of wetland hydrology and its implications on shorebird conservation in Bundala National Park (a Ramsar site), and Embilikele lagoons and found that human activities such as pollution had detrimental effects on bird communities. Kaluthota et al (2008) also conducted studies in Bundala on migratory wading bird communities. Chandana et al (2008) studied the factors affecting avifaunal distribution in three lagoons, namely, Malala, Embilikele, and Bundala and showed that salinity, water depth, and abundance of aquatic macrophytes were the key determinants of bird diversity there. However, the direct impacts of urbanization on estuarine avian diversity in Sri Lanka have not yet been addressed. This gap has been identified throughout the world too, as the conservation biologists focus predominantly on the protection of natural ecosystems and have placed little importance on

* Corresponding author. Tel.: +94 (0)11 2914479; fax: +94 (0) 11 2914479.

E-mail address: upali@kln.ac.lk (W.U. Chandrasekara).

Peer review under responsibility of National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA).




Figure 1. Satellite images of the Negombo estuary showing the eight study sites: Site 1 - Munnakkara (a highly populated village), Site 2 - Kadolkele (a pristine mangrove forest), Site 3 - Seeduwa (marsh lands adjacent to the Colombo-Katunayake expressway), Site 4 - Kindigoda (an undisturbed marsh and mangrove forest area near the head of the estuary), Site 5 - Kepungoda (a village with coconut plantations), Site 6 - Dungalpitiya (a village with a low human population density), Site 7 - Pitipana (an area with aquaculture grow-out ponds), Site 8 - Duwa (a fishing harbor/boating area). Image source: Google Earth Pro Software (Google Inc., 2013). Location of Negombo estuary in Sri Lanka is also shown.

urban biodiversity overall (Melles et al 2003). Therefore, it is important to focus on this particular aspect with regard to estuaries with varying degrees of urbanization among different land-use patterns associated with them.

The Muthurajawela marsh and Negombo estuary have received widespread interest and attention and, when considered together, are designated as a protected area for biodiversity conservation (Devendra 2002). As these two wetland ecosystems occur in one of the most populated regions of the west coast of the country, they are threatened by a variety of anthropogenic user patterns including fishery, agriculture, shipping, and habitation. This region has undergone a considerable land-use change over the past few decades (Sellamuttu et al 2011), particularly towards the northern side in the Negombo estuary. Although only a few studies have

been carried out to investigate the effects of these land-use changes on the floral and faunal communities around the Negombo estuary in general, their impacts on the avian diversity have never been addressed. Therefore, the present study was carried out to assess the changes of the avifaunal diversity in relation to land-use patterns around this estuary in Sri Lanka.

Materials and methods

Study area and sampling sites

The Negombo estuary (7° 6'–7° 12' N; 79° 49'–79° 53' E) is situated on the west coast of Sri Lanka (Figure 1). It is a shallow basin type estuary with a surface area of approximately 35 km². The

Download English Version:

<https://daneshyari.com/en/article/4395109>

Download Persian Version:

<https://daneshyari.com/article/4395109>

[Daneshyari.com](https://daneshyari.com)