

Heavy Metal Accumulations of 4 Species of Anseriformes in Korea

Seon-Deok Jin¹, Seul-Gi Seo², Young-Un Shin³, Ki-Chang Bing², Tae-Han Kang³,
Woon-Kee Paek¹ and Doo-Pyo Lee^{4*}

¹Research and Planning Division, National Science Museum, Daejeon 305-705, Korea

²Migratory Birds Center, Korea National Park Service, Sinan 535-916, Korea

³Korea Institute of Environmental Ecology, Daejeon 305-509, Korea

⁴Department of Biological Science, Honam University, Gwangju 506-714, Korea

Abstract: The study figured out the tendency of distribution of lead (Pb) and cadmium (Cd), and compared and examined the heavy metal accumulations among species by measuring the heavy metal accumulations in major organs (liver, muscle, kidney and bones) targeting 31 birds in 4 species (*Cygnus cygnus*, *Anser albifrons*, *Aix galericulata* and *Anas formosa*) of Anseriformes. Concentration of lead (Pb) in tissues was generally high in kidney (1.54 µg/wet g) or bones (5.94 µg/wet g), and concentration of cadmium (Cd) was generally high in kidney (0.474 µg/wet g) and low in muscles (0.019 µg/wet g). As shown above, there was no bird containing more than the standard amount of cadmium (Cd) among Anseriformes, but the study confirmed from 3 species (*Cygnus cygnus*, *Anser albifrons*, *Aix galericulata* and *Anas formosa*) containing lead at exposed levels and at poisoned levels. It is judged that the study result will be used as a comparing material for studies monitoring heavy metal of Anseriformes, a winter bird and a bio-indicator in an aquatic ecosystem and a basic material for reasonable environmental policy of lead shots and lead sinker in the future.

Keywords: Anseriformes, Lead (Pb), Cadmium (Cd), bio-indicator, Lead poisoning

Introduction

Heavy metals are not purified by the sanitation facilities or the nature's self-purification capacity unlike other pollutants, and their amount increases with the course of time in the natural world, causing biological magnification which is accumulated in a specific organism (Phillip, 1980). Heavy metals mean a metallic element with 4.0 or more specific gravity, and cadmium (Cd), mercury (Hg), lead (Pb), and antimony (Sb) are metals unnecessary in the living body, and they are not decomposed but accumulated in the body, causing harmful effects. Also, when they are ionized, they are easy to react with biogenic substances such as protein or nucleic acid, and have significant influences on 3-dimensional structure of protein or active revelation of enzyme. In addition, heavy metals are sometimes displaced with the necessary metals located at the center of vitality of metalloenzymes. In order to directly or indirectly evaluate the degree of exposure to heavy-metal contamination and the level of hazard, it is necessary to conduct biological monitoring using biological indicator species (Furness and

Greenwood, 1993). Birds which are located in higher trophic levels in the ecosystem and can provide information on the range of contamination in the whole food chains are widely used for monitoring heavy-metal contamination or as an indicator species (Blus *et al.*, 1993). Birds are known for being influenced by the distribution and accumulation of heavy metals in the body due to complex physiological functions such as breeding, growth, molt and spawning and various factors such as seasonal migration, types of food and the level of contamination in their food field (Hutton, 1981; Honda *et al.*, 1985; Honda *et al.*, 1986a, b, c). They show a close correlation between each organ and between elements in the process to absorb and redistribute heavy metals in the body. In such a process, they show difference in the level of distribution and accumulation of heavy metals in the body (Burger and Gochfeld, 2000). Also, as birds live in various kinds of environment including deserts, forests, grasslands, rivers, lakes and oceans, have various kinds of food habits including herbivorous, insectivorous, carnivorous, ichthyophages and omnivorous, and have different lifespans from several years to several decades, they have a feature to accumulate high-concentration of various pollutants emitted in the environment in their body (Lee, 1995).

A study on heavy-metal accumulation in Anseriformes is

*To whom correspondence should be addressed.

Tel: +82-62-940-5434

E-mail: dplee@honam.ac.kr

Table 1. List of bird analyzed

Species	*N	Sampling date	Sampling site	Cause of death (n)
<i>Cygnus cygnus</i>	5	00.12.~02.12.	Jeollanam-do Yeosu-si Chungcheongnam-do Dangjin-gun Chungcheongnam-do Taean-gun Chungcheongnam-do Seosan-si	Unknown (3) Exhaustion (1) Shot (1)
<i>Anser albifrons</i>	10	00.10.~04.03	Chungcheongnam-do Seosan-si Gyeonggi-do Yeuju-gun, Paju-si Gangwon-do Cheorwon-gun Incheon Ganghwa-gun Jeollabuk-do Gunsan-si	Cholera (1) Exhaustion (1) Poison (8)
<i>Aix galericulata</i>	12	01.03.	Gyeongsangnam-do Sacheon-si	Poison
<i>Anas formosa</i>	4	00.10	Chungcheongnam-do Seosan-si	Cholera
Total	31			

*N: Number of individual

considered as an appropriate bio-indicator toward heavy-metal pollutants in water environment (Scheuhammer, 1989; Congiu *et al.*, 2000). As for the study on heavy-metal accumulation in Anseriformes in Korea, there is a study on the difference among species in the accumulating level of heavy metals in birds' organs (Lee, 1995), examining several species belonging to Anseriformes, but it is not sufficient domestically.

Therefore, this study was conducted in order to provide the basic materials for monitoring environmental pollution by figuring out the tendency of distribution of elements through the measurement of the concentration of heavy metals in birds' body targeting 4 species of Anseriformes including *Cygnus cygnus*, *Anser albifrons*, *Aix galericulata* and *Anas Formosa* and by comparing and examining the concentration of heavy metals between species.

Materials and Methods

The study subjects are 31 individuals of 4 species collected by being supported by the Korean Association for Bird Protection from 2000 to 2004 (Table 1). In order to analyze heavy metals in the body, collected birds were dissected for measuring the weight of each part, 1-5 g of specimen was completely disassembled in the Kjeldahl digestion apparatus using sulfuric acid, nitric acid and perchloric acid, and the dissolved solution was diluted into 100ml solution. The dissolved solution was enriched and extracted in DDTC-

MIBK method, and lead (Pb) and cadmium (Cd) were analyzed using an atomic absorption spectrometer (Shimadzu AA-6400, Co. Ltd, Japan) (Lee *et al.*, 1989). Detection limit was 0.1 µg/wet g or less for lead (Pb) and 0.01 µg/wet g or less for cadmium (Cd). Statistical analysis of data was conducted using SPSS program (version 10.0, SPSS), and as the concentration of heavy metals was abnormal distribution, Kruskal-Wallis test was used for comparing concentration of each element between species through a non-parametric test for data analysis (Alessandra *et al.*, 2005).

Results

Lead (Pb)

The concentration of lead (Pb) was generally high in kidney or bones. Lead (Pb) in liver was the highest in *Aix galericulata* (3.18 µg/wet g), and the lowest in *Cygnus cygnus* (0.39 µg/wet g) ($\chi^2=46.4$, $p<0.05$). As for muscles, it was the highest in *Anser albifrons* (2.60 µg/wet g) and the lowest in *Aix galericulata* (0.48 µg/wet g) ($\chi^2=46.0$, $p<0.05$). In kidney, it was the highest in *Aix galericulata* (2.40 µg/wet g) and the lowest in *Cygnus cygnus* (0.69 µg/wet g) ($\chi^2=46.8$, $p<0.05$). In bones, *Anser albifrons* showed the highest concentration by about 20 times (20.7 µg/wet g) compared with other species, and *Cygnus cygnus* showed the lowest concentration (0.59 µg/wet g) ($\chi^2=55.3$, $p<0.05$) (Table 2).

Table 2. Lead (Pb) concentration (µg/wet g) in tissues of 4 bird species from Korea

Family/species	N		Liver	Muscle	Kidney	Bone
<i>Cygnus cygnus</i>	5	Median Range	0.56 0.03~28.9	0.55 0.11~2.81	0.69 0.27~17.7	0.59 0.04~23.2
<i>Anser albifrons</i>	10	Median Range	0.61 0.08~34.4	0.63 0.03~7.85	0.93 0.01~95.2	1.17 0.16~24.7
<i>Aix galericulata</i>	12	Median Range	3.18 2.11~303	0.48 0.22~0.94	2.40 1.50~4.41	1.30 0.90~1.95
<i>Anas formosa</i>	4	Median Range	0.88 0.78~4.02	2.60 0.60~5.61	2.12 0.60~10.9	20.7 8.79~147

Download English Version:

<https://daneshyari.com/en/article/4399522>

Download Persian Version:

<https://daneshyari.com/article/4399522>

[Daneshyari.com](https://daneshyari.com)