

Update of the 2014 Drought on California Rangelands

By Royce E. Larsen, Marc R. Horney, and Daniel Macon

On the Ground

- Droughts are common on California rangelands.
- The current drought in California is listed as severe or exceptional for most of the state.
- The drought has affected rangeland, and the live-stock industry, more than other commodities.
- The actual costs associated with this drought are just beginning to be realized.

Keywords: drought, California, rangelands.

Rangelands 36(5):52–58

doi: 10.2111/Rangelands-D-14-00032.1

© 2014 The Society for Range Management

Drought is a recurring phenomenon characteristic of arid and semiarid landscapes that challenges efforts to make reliable uses of them. The United States has experienced mounting costs from droughts over the last 30 years. The costs of the 2012 drought alone surpassed \$50 billion.¹ In California, as in other western states, costs of the 2014 drought are already significant and growing rapidly. By late spring 2014, preliminary estimates of agricultural economic losses exceeded \$2.2 billion, and more than 17,000 jobs had been lost.² Though drought impacts many agricultural segments, our main interest here is how it has impacted rangelands in California, particularly rangelands in the Central Coast region.

California Climate and Common Rangeland Types

Much of California enjoys a Mediterranean climate that is largely unique in North America. It is characterized by cool wet winters and hot dry summers. While occasional weather systems may bring some monsoonal moisture in the summertime, those events are presently rare and short-lived. California's large geographic extent and considerable topographic variation have yielded a great many ecological sites and associated plant communities. One broad division is between the annual grassland type, which commonly exists

below 1,200 feet in elevation³ and is made up of naturalized species, many introduced from the Mediterranean region of southern Europe, and the perennial-grass-dominated Great Basin sagebrush-steppe type, which is common on the east slope of the Sierra Nevada and Cascade mountains and the Modoc plateau in the NE portion of the state. The annual grasslands of the Central Valley and Central Coast regions are often codominant with oak woodlands or savannas, which in many areas are vestiges of their former extent.⁴ Weather and growing conditions for California rangelands are driven by complex interactions among coastal influences, elevation and other orographic effects, and distribution of summertime heat from the central valleys and southern deserts. Precipitation amounts and patterns can vary substantially over relatively short distances in response to these interactions.

History of Droughts in California

According to tree ring records there have been several multi-decadal dry periods in the past 500 years in California. These records reveal a major dry period from 1760 to 1820⁵ and another drought from 1860 to 1885. Severe droughts in 1850–1851 and 1862–1864, together with other factors, including grazing practices, have been implicated in the conversion of the former native perennial grassland to a vegetation type that is dominated by annual grasses and forbs.⁶ The severe drought from 1862 to 1864 was also implicated in a major shift from livestock to cropland production in California.⁷

There have been at least eight multiyear periods of low precipitation in California since 1900. Droughts that exceed three years are uncommon, though occurrences in the past century include 1929–1934, 1947–1950, and 1987–1992. One remarkable example of drought in California was the two-year dry period in 1976 and 1977. Precipitation during each of these calendar years ranked among five lowest amounts ever recorded in California.⁴

In some years low precipitation results in forage production that is well below average. Because the amount and dependability of precipitation increases from south to north, and with elevation, the frequency of years with forage production significantly below average varies greatly across the

Lessons From the Dust Bowl Drought and Rangeland Conversion

By Dan Macon, Flying Mule Farm: A Producer's Viewpoint

In late January 2014, I attended the 9th annual California Rangeland Conservation Coalition summit in Oakdale, California. While past summits have focused on the threats posed by real estate development, this year's event focused on a different and, for me, unexpected threat—the conversion of our rangelands to orchards and vineyards. After returning home from this year's summit, I happened to start watching Ken Burns' latest PBS documentary, *The Dust Bowl*. As I watched, I was struck by the similarity between the attitudes and actions of farmers in the Southern Plains in the years leading up to the Dust Bowl and our own efforts to convert “unproductive” grasslands into intensively farmed orchards. At the risk of being melodramatic, I think we should consider the lessons of the Dust Bowl and the parallels with today's rangeland conversion in California.

My own “shepherd” definition of rangeland is any land that is too steep, too dry, too wet, too “something” for cultivated agriculture. Prehistorically, our rangelands were grazed by wild ungulates. Today, California's rangelands are the foundation of our livestock industry: cattle and sheep have grazed on rangelands in California for more than two and a half centuries. Rangelands continue to provide habitat for an incredible array of wildlife and native plants—everything from raptors to reptiles and the majestic oak to the smallest wildflower. Scientists are increasingly convinced that well-managed livestock grazing is crucial to the health of the native flora and fauna on our rangelands. And ranching, as an economic endeavor, is crucial to keeping these rangelands intact: it seems that ranching and wildlife both require large, contiguous tracts of open land.

The wheat boom of the 1920s resulted in the plowing of native prairie and planting of wheat on millions of acres in the Southern Plains. The price of wheat, driven up by World War I, seemed to be on an endless upward path. Even though the Plains were prone to periodic drought, historian Donald Worster says, “Promoters promised that the very act of farming would increase the precipitation—the rain follows the plow.” Wayne Lewis, who grew up on a farm in the Oklahoma panhandle, adds, “In the late 20s, the crops were good, the prices were good and so everybody ... the thing to do was to break out everything and get it in wheat.”¹⁸

Not everyone was convinced that the wheat boom was beneficial or sustainable. According to Duncan,¹⁸ “A handful of old timers, especially the cattlemen who had been there through those droughts, weren't so sure [about sod-busting the Plains]. To them, the Southern Plains were a grassland and the sod should never be turned.” Following the stock market crash in 1929 and several years of record wheat crops, the price of wheat crashed. As prices fell, the federal government encouraged farmers to plant

state's annual-type rangelands. Analysis of annual forage production data from 26 locations in California's annual rangelands reveal that a 50% reduction in range forage production rarely occurs north of Sacramento.⁸ The 50% reduction value is important because that is the value used by the USDA Farm Service Agency to begin offering assistance programs like the Noninsured Crop Disaster Assistance Program. Forage losses greater than 50% are more common in the rain shadow of the Coast Range adjacent to the west edge of the San Joaquin Valley.⁸

California's annual rangeland forage production also varies greatly over short distances due to variations in precipitation, soil characteristics, and topography. For example, data from San Luis Obispo County reveal that forage reductions are less frequent at coastal sites than inland sites.⁹ This high variability from year to year and from site to site makes it difficult for producers to manage livestock enterprises.

Current Drought in California

State of the Drought

The current drought in California has placed the majority of the state in the severe or exceptional drought category according to the US Drought Monitor.¹⁰ Even though this year has received much attention from the media, California has actually been experiencing drought conditions for at least six out of the last eight years (Fig. 1). The 2013–2014 water year was exceptionally low in rainfall and snowpack, and most reservoirs did not fill to capacity, many communities are demanding water cuts, crops are not being watered, and forage production for livestock was significantly below average this year. A recent report shows that the 2014 drought is responsible for the greatest reduction in surface water availability ever measured.² Groundwater is expected to make up 75% (5 million acre-feet) of the water shortage (6.5 million acre-feet) for agriculture during 2014. In addition to higher pumping costs, this could lead to depletion of groundwater sources, wells may dry up, and land may subside. The net water shortage will affect the Tulare Basin, in the southern Central Valley the most,² which is in the “exceptional drought” zone. Although most crops may be irrigated by pumping groundwater, where available, rangeland productivity depends on rainfall. Rangeland use is not only dependent on forage availability but water availability as well. It has not been uncommon for pastures to be left with forage still available simply because drinking water sources for livestock, whether stock-water ponds, streams, springs, or wells, have run dry.

It is common that droughts affect the areas south of Sacramento more often than those north of Sacramento.⁸ The Central Coast of California, which is more than 200 miles south of Sacramento, is in the heart of the D4 (Exceptional) drought rating area of the state (Fig. 1). Within this area, the city of Paso Robles, California, has kept rainfall records since 1887 (Fig. 2). The water years 2006–2007, 2008–2009, 2011–2012, 2012–2013, and 2013–2014 were all well below average. Annual forage production has been monitored in San Luis

Continued

Download English Version:

<https://daneshyari.com/en/article/4405264>

Download Persian Version:

<https://daneshyari.com/article/4405264>

[Daneshyari.com](https://daneshyari.com)