



ORIGINAL ARTICLE

A new record of *Myxobolus brachysporus* and *M. israelensis* in the tilapia (*Oreochromis niloticus*) collected from the Nile River, Egypt



Abdel-Azeem S. Abdel-Baki ^{a,b,*}, Eman Zayed ^b, Thabet Sakran ^b,
Saleh Al-Quraishy ^a

^a Zoology Department, College of Science, King Saud University, Saudi Arabia, P.O. Box 2455, Riyadh 11451, Saudi Arabia

^b Zoology Department, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt

Received 11 November 2014; revised 28 December 2014; accepted 10 January 2015

Available online 19 January 2015

KEYWORDS

Myxozoa;
Fish;
Nile;
Tilapia

Abstract The present study was carried out as part of an ongoing general survey for myxosporean parasites infecting tilapias in the River Nile, Egypt. In the present study, 77 Nile tilapia (*Oreochromis niloticus*) were collected from boat landing sites at Beni-Suef governorate, Egypt and examined for the myxosporean infection. The infection was encountered as a huge number of free spores in the kidney and the spleen. The infection showed a prevalence of 51.9% (40/77) for *Myxobolus brachysporus* while it was 25.9% (20/77) for *Myxobolus israelensis*. Mature spores of *M. brachysporus* were ellipsoidal and measured $8.6 \times 13.2 \mu\text{m}$. The polar capsules were subcircular with 5–6 filament turns and measured $4.7 \times 3.6 \mu\text{m}$. Spores of *M. israelensis* were ellipsoidal in the frontal view and fusiform in the lateral view. Spore measurements were $13.4 \mu\text{m}$ long and $8.7 \mu\text{m}$ wide. The polar capsules were elongated with 6–7 filament coils and measured $8.6 \times 3.1 \mu\text{m}$. The findings presented here proved that tilapia fishes in the Nile River are still suffering from infections with *Myxobolus* species. Therefore, further studies should be carried out to survey the *Myxobolus* infection among tilapias under culture conditions to clarify the pathological impacts of this parasite in tilapias aquaculture.

© 2015 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author at: Zoology Department, College of Science, King Saud University, Saudi Arabia, P.O. Box 2455, Riyadh 11451, Saudi Arabia. Tel.: +966 (1)4675754.

E-mail address: azemal@yahoo.com (A.-A.S. Abdel-Baki).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

1. Introduction

It is quite certain that fishes represent the foremost necessary sources of animal protein all over the world. Since the beginning, fishing is considered one of the most important activities that is practiced by ancient Egyptians (Abdel-Ghaffar et al., 2008). Tilapia are essentially freshwater fish commonly found in rivers, streams, lakes, ponds and less frequently inhabiting brackish and even marine water. The overall production of

tilapia culture in Egypt throughout 2007 was calculable by 265,862 Ton that represents about 41.8% of the whole national fish culture production (GAFRD, 2010). Myxosporeans are an abundant and diverse group of economically important microscopic parasites, which cause disease in a large variety of commercially important fishes including tilapias (Lom and Dykova, 2006). Myxosporean parasites are known to be responsible for several forms of damage, including post-mortem myoliquefaction of the host (Pampoulie et al., 1999), damage of ovaries (Gbankoto et al., 2001; Mansour et al., 2013) and reduction of the capacity of respiration (Molnár and Székely, 1999). Among the myxosporeans, species of the genus *Myxobolus* are, so far, the foremost unremarkably found in fish, with about 856 known species throughout the world (Eiras et al., 2014). Of them, about 12 species were described from tilapias spp. in natural and cultural habitats. In the present study, we described two *Myxobolus* species from the Nile tilapia *Oreochromis niloticus* collected from the Nile River at Beni-Suef governorate which were delineated for the first time from Egypt.

2. Materials and methods

During the present investigation, 77 tilapia fish (*Oreochromis niloticus*) live or freshly caught were collected from boat landing sites at the Beni-Suef governorate (29°3'50"N, 31°5'20"E), throughout a period from October 2012 to October 2013. Fish were collected from both sexes and with lengths ranging from 15 to 20 cm and weighs about 150–200 g. Skins and gills scraps were performed from all fishes and examined for myxosporean infection. Also, eyes of the fish were examined for the presence of any myxosporean spores within both the cornea and lens. Further, the fish were dissected then impression smears from different organs including liver, spleen, kidneys, gall bladder were made. Stomach and intestinal scraps were also made.

All samples were freshly examined using a regular light microscope as well as a dissecting microscope. Fresh spores were examined and photographed using differential interference contrast Zeiss Axiovert 135 microscope equipped with a camera. The morphometric measurements of spores followed the guidelines devised by Lom and Arthur (1989) for species descriptions of Myxosporea. Data are represented as: mean \pm SD followed by (Range).

3. Results

The infection was encountered as a huge number of free spores in the kidney and the spleen. The infection showed a prevalence of 51.9% (40/77) for *Myxobolus brachysporus* whereas it was 25.9 (20/77) for *Myxobolus israelensis*. The recorded species are described as follows:

3.1. *Myxobolus brachysporus* Baker, 1963

The spores were ellipsoidal in shape and characterized by their width that greatly exceeded the length (Figs. 1, panels 1–3 and 2). They measured 8.6 ± 0.4 (7.8–9.2) long \times 13.2 ± 0.6 (12.1–14.2) wide. The polar capsules were subcircular, mostly equal in size and measured 4.7 ± 0.3 (4.2–5.1) long \times 3.6 ± 0.3 (3.2–4.2) wide. The polar filament spired in 5–6 turns perpendicular or slightly oblique to the longitudinal axis of the capsule. The sporoplasm filled up the rest of the spore cavity (Figs. 1, panels 1–3 and 2).

3.2. *Myxobolus israelensis* Landsberg, 1985

The spores were ellipsoidal with rounded anterior and posterior ends in the frontal view and fusiform in the lateral view (Figs. 1, panels 4–6, 3). The spores measured 13.4 ± 0.9




Figure 1 Light micrographs of fresh mature spores of *Myxobolus brachysporus* (panels 1–3) and *Myxobolus israelensis* (panels 4–6) infecting the kidney and the spleen of Nile tilapia *Oreochromis niloticus* from the River Nile.

Download English Version:

<https://daneshyari.com/en/article/4406326>

Download Persian Version:

<https://daneshyari.com/article/4406326>

[Daneshyari.com](https://daneshyari.com)