



ORIGINAL ARTICLE

# Assessment of the role of general, biochemical and family history characteristics in kidney stone formation



Faiza Jabbar <sup>a</sup>, Muhammad Asif <sup>a,b</sup>, Hajirah Dutani <sup>a</sup>, Abrar Hussain <sup>a</sup>, Arif Malik <sup>c</sup>, Mohammad Amjad Kamal <sup>d</sup>, Mahmood Rasool <sup>e,\*</sup>

<sup>a</sup> Department of Biotechnology and Informatics, BUIITEMS, Quetta, Pakistan

<sup>b</sup> Office of Research Innovation and Commercialization, BUIITEMS, Quetta, Pakistan

<sup>c</sup> Institute of Molecular Biology and Biotechnology (IMBB), The University of Lahore, Lahore, Pakistan

<sup>d</sup> King Fahd Medical Research Center, King Abdulaziz University, Jeddah, Saudi Arabia

<sup>e</sup> Center of Excellence in Genomic Medicine Research (CEGMR), King Abdulaziz University, Jeddah, Saudi Arabia

Received 27 May 2014; revised 3 June 2014; accepted 3 June 2014

Available online 12 June 2014

## KEYWORDS

Risk factors;  
Kidney stone formation;  
Nephrolithiasis;  
Water intake

**Abstract** *Aim:* The main objective of the study was to determine the urinary risk factors involved in kidney stone formation.

*Method:* In this study a total number of 101 patients (64 males and 37 females) between the age group 2 and 70 years were selected. Personal characteristics like age, family history, clinical sign and symptoms, education, monthly income, living style, smoking or tobacco chewing habit, dietary intake and daily amount of drinking water were recorded.

*Results:* The study showed that the risk of kidney stone formation was high in the median age group (16–25 years) both in male and female population. The most important factors associated with this were lack of drinking clean water, over weight and obesity as well as family history (37.5% and 27.02% in men and women, respectively).

*Conclusion:* Our study has confirmed that lack of drinking sufficient amount of water, increasing weight and obesity and family history are some major factors contributing to the increased risk of kidney stone formation. Therefore it is very important to live a healthy life, drink clean water and control weight to prevent such diseases.

© 2014 Production and hosting by Elsevier B.V. on behalf of King Saud University.

\* Corresponding author. Address: Center of Excellence in Genomic Medicine Research (CEGMR), Post Box No. 80216, King Abdulaziz University, Jeddah 21589 Saudi Arabia. Tel.: +966 6401000x25479; fax: +966 6952521.

E-mail address: [mahmoodrasool@yahoo.com](mailto:mahmoodrasool@yahoo.com) (M. Rasool).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

## 1. Introduction

The prevalence and incidence of kidney stone is a major cause of death all over the world. Life time prevalence of symptomatic nephrolithiasis is around 10% in men and 5% in women and about more than \$2 billion spent each year for treatment purpose (Taylor and Curhan, 2008; Taylor et al., 2005). A number of different kinds of factors are involved in

increasing the risk of kidney stone formation like; excess calcium, phosphate, oxalate and uric acid in the urine, inadequate hydration, lack of stone inhibitors in the urine, family history of stone (Curhan et al., 1997), daily urine volume, high, large body size (Curhan et al., 1998), some medications and ongoing urine infection (Sowers et al., 1998; Stamatelou et al., 2003; Pandeya et al., 2010; Leonetti et al., 1998). Dietary risk factors play a very vital role in stone formation. There is a proof that diminished fluid and calcium consumption is a strong risk factor (Stamatiou et al., 2006; Hirvonen et al., 1999), increased consumption of oxalate is also a major contributor to enhance the stone formation (Taylor and Curhan, 2008). It is verified by epidemiological studies that increased sodium, salt and animal proteins intake have an equivocal impact on stone formation risk (Curhan et al., 1997; Stamatelou et al., 2003). The global climate changes, which is an environmental factor also promote the rates of kidney stone disease. The broad consensus is that average global temperatures have increased Curhan et al. (1997). Common clinical conditions involving the kidney stone formation have been linked to a number of medical co morbidities including obesity (Taylor and Curhan, 2008), diabetes mellitus, hypertension (Cappuccio et al., 1990), chronic kidney disease, and cardiovascular problems (Rule et al., 2009). Stones of kidneys can be easily diagnosed with sudden onset of pain, blood excretion from urine and stones that appear on X-ray. Analyzing the stone prior to treatment is important because it helps to decide on the different options for treatment. Majority of the stones can be treated without undergoing surgery and about 90% of the stones will pass by themselves within 3–6 weeks. In these cases the only medication required is pain relief. In cases where the pain onset is severe and unbearable then hospital admission and analgesia may be required (Stamatiou et al., 2006).

The aim of the present study is to assess prevalence of kidney stone diseases in different age groups and to evaluate the association of self history of kidney stones with age, sex, history and geographical residence.

## 2. Materials and methods

### 2.1. Setting and study area

The study was conducted in the Quetta city and samples were collected from Bolan Medical Complex Hospital and Sandeman Civil Hospital, Quetta, Pakistan. Informed consent was taken from all participants. The study was approved by the local ethics committee of Balochistan University of Information Technology and Management Sciences (BUIITEMS), Quetta Pakistan. A total of 101 patients (64 males and 37 females) were selected and information was taken in a detailed questionnaire. The study was performed from June 2011 to February 2012.

### 2.2. Sample and data collection

All the required information was collected by professional doctors and nursing staff. Questionnaire used for information which included, name, age, gender, location of patients, sign and symptoms, family history, daily amount of drinking water, source and treatment of water, and ultrasound reports. On the

basis of age, patients were divided into 5 groups, 0–15, 16–25, 26–35, 36–45, 46–55 and above 55, respectively.

### 2.3. Blood collection and laboratory investigation

The skin was cleaned thoroughly and sterilized with 70% isopropyl alcohol swab (Kandall HealthCare, USA) and dried before withdrawing 2 ml peripheral blood by a 5 cc disposable syringe (Becton Dickinson Pak) from enrolled subjects. The blood was transferred to an ethylenediamine tetra acetic acid (EDTA) coated purple-top test tube. The blood was mixed in the test tubes with 5 complete inversions and tubes were marked with codes and immediately taken to the laboratory for investigation.

Complete blood cell (CBC) count, white blood cells (WBC), red blood cells (RBC), hemoglobin (Hb), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), (RDW) and platelets, were measured in this study.

The glucose level of each individual was carried out by using a glucometer (Accu-Check Active, Roche Company). Concentrations of urea and creatinine were estimated by the direct kit method using precipitant of Spinreact Co, Spain estimated in the sera of the samples.

### 2.4. Data analysis

SPSS 16 version was used for data analysis. All the results were checked by descriptive statistics. Results were expressed by mean standard and *P* value was used for checking significance level at <0.05. Graphs were also drawn for comparison between two groups.

## 3. Results

A total number of 101 kidney stone subjects were indentified in which male were 64 (63.36%) and female were 37 (36.63%).

**Table 1** General characteristics of kidney stone patients.

Characteristics	Male <i>N</i> = 64	Female <i>N</i> = 37
<i>Age (years)</i>		
1–15	16 (25%)	10 (27.02%)
16–25	24 (37.5%)	11 (29.72%)
26–35	8 (12.5%)	9 (24.32%)
36–45	5 (7.81%)	6 (16.21%)
46–55	5 (7.81%)	3 (8.10%)
56 and above	3 (4.68%)	1 (2.70%)
<i>Family history of kidney stone</i>		
Yes	24 (37.5)	10 (27.02)
No	40 (62.5)	27 (72.97)
<i>Location</i>		
Urban	32 (50%)	29 (78.37%)
Rural	29 (45.31%)	11 (29.72%)
<i>History about disease</i>		
Hypertension	12 (18.75%)	05 (13.51%)
CHD	08 (12.5%)	04 (10.81%)
Obesity	10 (15.62%)	7 (18.91%)

Download English Version:

<https://daneshyari.com/en/article/4406362>

Download Persian Version:

<https://daneshyari.com/article/4406362>

[Daneshyari.com](https://daneshyari.com)