



King Saud University

Saudi Journal of Biological Sciences

www.ksu.edu.sa
www.sciencedirect.com



الجمعية السعودية لعلم الحياتة
SAUDI BIOLOGICAL SOCIETY

ORIGINAL ARTICLE

Studies on the role of goat heart galectin-1 as an erythrocyte membrane perturbing agent



Ghulam Md Ashraf ^{a,*}, Asma Perveen ^b, Syed Kashif Zaidi ^c, Shams Tabrez ^a,
Mohammad A. Kamal ^a, Naheed Banu ^{b,d}

^a King Fahd Medical Research Center, King Abdulaziz University, P.O. Box 80216, Jeddah 21589, Saudi Arabia

^b Department of Biochemistry, Faculty of Life Sciences, Aligarh Muslim University, Aligarh 202002, U.P., India

^c Center of Excellence in Genomic Medicine Research, King Abdulaziz University, P.O. Box 80216, Jeddah 21589, Saudi Arabia

^d College of Medical Rehabilitation, Qassim University, Buraydah, Saudi Arabia

Received 22 May 2014; revised 25 September 2014; accepted 26 September 2014

Available online 5 October 2014

KEYWORDS

Goat heart galectin-1;
Agglutination;
Erythrocyte;
Membrane;
Hemolysis

Abstract Galectins are β -galactoside binding lectins with a potential hemolytic role on erythrocyte membrane integrity and permeability. In the present study, goat heart galectin-1 (GHG-1) was purified and investigated for its hemolytic actions on erythrocyte membrane. When exposed to various saccharides, lactose and sucrose provided maximum protection against hemolysis, while glucose and galactose provided lesser protection against hemolysis. GHG-1 agglutinated erythrocytes were found to be significantly hemolyzed in comparison with unagglutinated erythrocytes. A concentration dependent rise in the hemolysis of trypsinized rabbit erythrocytes was observed in the presence of GHG-1. Similarly, a temperature dependent gradual increase in percent hemolysis was observed in GHG-1 agglutinated erythrocytes as compared to negligible hemolysis in unagglutinated cells. The hemolysis of GHG-1 treated erythrocytes showed a sharp rise with the increasing pH up to 7.5 which became constant till pH 9.5. The extent of erythrocyte hemolysis increased with the increase in the incubation period, with maximum hemolysis after 5 h of incubation. The results of this study establish the ability of galectins as a potential hemolytic agent of erythrocyte membrane, which in turn opens an interesting avenue in the field of proteomics and glycobiology.

© 2014 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Abbreviations: GHG-1, goat heart galectin-1; PBS, phosphate buffered saline.

* Corresponding author.

E-mail addresses: ashraf.gm@gmail.com, gashraf@kau.edu.sa (G.M. Ashraf).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

1. Introduction

Galectins are β -galactoside specific binding lectins reported to be expressed ubiquitously in all taxa of living world and is involved in vital biological functions (Hasan et al., 2007). The ability of galectins to act as humoral factors in the defense mechanism against various pathogenic agents suggests important applications for these proteins (Hussain et al., 2013; Liu et al., 2014; Shi et al., 2014). Over the years, galectins have

been reported to have potential hemolytic actions on membranes and other cell organelles (Armstrong et al., 1996; Unno et al., 2014). Lytic action of some galectins has been ascribed to enzymatic activity (Hittelet et al., 2003), perturbation of the activities of membrane associated enzymes (Lowe and Marth, 2003), and pore formation in the membranes (Yu et al., 2002). In the present study, the hemolytic actions of purified goat heart galectin-1 (GHG-1) (Ashraf et al., 2011) have been studied by monitoring their effect on erythrocyte membrane integrity and permeability to understand the possible role of galectins in hemolytic destruction of foreign cells in the mammalian nervous system. The effect of various physiological parameters like concentration, temperature, pH and incubation time on GHG-1 mediated erythrocyte membrane lysis was also investigated.

The galectin dependent association of glycans has been suggested to potentially affect the key membrane features such as permeability, fluidity and osmofragility, and enhances its impact on membrane characteristics by interacting with hydrophobic membrane patches (Gupta et al., 2006). Moreover, glycan residues attached to cell membranes play a crucial role in membrane lysis (Semrau et al., 2010; St-Pierre et al., 2011). Galectins are the lectins with specific attraction for β -galactoside moieties, and the presence of this glycan residue makes them a potent membrane lysis molecule (Yu et al., 2002). Hence, the effect of various saccharides on erythrocyte hemolysis was also investigated to assess the ability of non-electrolytes to protect GHG-1 induced lysis of erythrocytes, and also to identify the sugar molecule providing maximum protection against membrane lysis.

2. Materials and methods

2.1. Reagents

Sephadex G100 and G50, molecular weight markers (14.4–97.4 kDa), coomassie brilliant blue (CBB) G-250 and R-250, and sugars were purchased from Sigma Aldrich (St Louis, MO, USA). All other chemicals used were of analytical grade and purchased from Qualigens Fine Chemicals and Merck India Ltd, India. All the experiments were carried out in triplicates and mean value has been reported.

2.2. Isolation and purification of GHG-1

The GHG-1 was isolated and purified in essence according to our previously published studies on heart lectins (Ashraf et al., 2010a,b, 2011). Protein concentration was estimated by Lowry et al. (1951) and its activity was determined by hemagglutination activity using trypsinized rabbit erythrocytes by two fold serial dilutions on a microtiter V-shaped plate (Laxbro, India) (Raz and Lotan, 1981).

2.3. Isolation of erythrocytes and trypsinization

A fresh blood sample from the vein of a healthy rabbit was collected in a tube containing heparin. The erythrocytes were isolated by centrifuging the blood sample at 2000 rpm for 5 min at room temperature and washed three to four times with Phosphate buffered saline (PBS) 'B' (5 ml per ml packed cells). Washed erythrocytes were then made 4% (v/v) by adding PBS

'B'. An 8% (v/v) suspension of washed erythrocytes was prepared in PBS 'B'. A trypsin solution (100 mg%) was then added to the erythrocytes (0.1 ml of trypsin solution per ml of erythrocyte suspension) and incubated for 1 h at 37 °C. The trypsinized erythrocytes were washed four to five times with PBS 'B' and 4% (v/v) suspension was prepared for further use.

2.4. Effect of GHG-1 mediated agglutination on the osmofragility of trypsinized rabbit erythrocytes

The osmofragility of trypsinized rabbit erythrocytes in the presence of GHG-1 was determined according to a previously described method (Pande et al., 1998) with some modifications. An 8% trypsinized rabbit erythrocyte suspension (200 μ l) was mixed with 100 μ l GHG-1 (100 μ g/ml in 50 μ l of normal saline containing 5 mM β -ME) and incubated for one hour at room temperature to allow agglutination, followed by the addition of 1.5 ml hypotonic NaCl solution with concentrations ranging from 0.3% to 0.8%. This was then followed by incubation of cells for 4 h at room temperature with intermittent gentle shaking and centrifuged at 2000 rpm for 5 min. The supernatant was collected after centrifugation, the extent of hemolysis was measured at 540 nm and the results were reported as percent lysis. The lysis of erythrocytes with an equal volume of distilled water was taken as 100%.

2.5. Effect of varying concentrations of GHG-1 on erythrocyte hemolysis

An 8% trypsinized rabbit erythrocyte suspension (200 μ l) treated with varying concentrations of 100 μ l GHG-1 (10–100 μ g/ml in 50 μ l of normal saline containing 5 mM β -ME) was incubated for 4 h at 37 °C. Samples were centrifuged and RBC lysates were analyzed at 540 nm.

2.6. Hemolysis of GHG-1 agglutinated erythrocytes as a function of increasing temperature

An 8% trypsinized rabbit erythrocyte suspension (200 μ l) in the presence and absence of 100 μ l GHG-1 (100 μ g/ml in 50 μ l of normal saline containing 5 mM β -ME) was incubated at increasing temperatures (10–50 °C) for 4 h. Samples were centrifuged and RBC lysates were analyzed at 540 nm.

2.7. pH induced hemolysis of GHG-1 agglutinated erythrocytes

An 8% trypsinized rabbit erythrocyte suspension (200 μ l) in the presence and absence of 100 μ l GHG-1 (100 μ g/ml in 50 μ l of normal saline containing 5 mM β -ME) was treated at varying pH values (3.5–11.5) at 37 °C for 4 h. The buffers used were, 0.1 M sodium acetate buffer (pH 3.5–5.5), 0.1 M sodium phosphate buffer (pH 6.5–7.5), 0.1 M Tris-HCl buffer (pH 8.5–9.5) and 0.1 M glycine-NaOH buffer (pH 10.5–11.5) for 24 h at 4 °C. Samples were centrifuged and RBC lysates were analyzed at 540 nm.

2.8. Hemolysis of GHG-1 agglutinated erythrocytes as a function of incubation period

An 8% trypsinized rabbit erythrocyte suspension (200 μ l) in the presence and absence of 100 μ l GHG-1 (100 μ g/ml in

Download English Version:

<https://daneshyari.com/en/article/4406369>

Download Persian Version:

<https://daneshyari.com/article/4406369>

[Daneshyari.com](https://daneshyari.com)