



Enhanced stability of a naringenin/2,6-dimethyl β -cyclodextrin inclusion complex: Molecular dynamics and free energy calculations based on MM- and QM-PBSA/GBSA

Waratchada Sangpheak^a, Wasinee Khuntawee^b, Peter Wolschann^{c,d}, Piamsook Pongsawasdi^a, Thanyada Rungrotmongkol^{a,*}

^a Department of Biochemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

^b Nanoscience and Technology Program, Graduate School, Chulalongkorn University, Bangkok 10330, Thailand

^c Department of Pharmaceutical Technology and Biopharmaceutics, University of Vienna, Vienna 1090, Austria

^d Institute of Theoretical Chemistry, University of Vienna, Vienna 1090, Austria

ARTICLE INFO

Article history:

Received 15 January 2014

Received in revised form 27 February 2014

Accepted 1 March 2014

Available online 12 March 2014

Keywords:

Naringenin

Cyclodextrin

Inclusion complex

Binding free energy

MM- and QM-PBSA/GBSA

ABSTRACT

The structure, dynamic behavior and binding affinity of the inclusion complexes between naringenin and the two cyclodextrins (CDs), β -CD and its 2,6-dimethyl derivative (DM- β -CD), were theoretically studied by multiple molecular dynamics simulations and free energy calculations. Naringenin most likely prefers to bind with CDs through the phenyl ring. Although a lower hydrogen bond formation of naringenin with the 3-hydroxyl group of DM- β -CD (relative to β -CD) was observed, the higher cavity could encapsulate almost the whole naringenin molecule. In contrast for the naringenin/ β -CD complex, the phenyl ring feasibly passed through the primary rim resulting in the chromone ring binding inside instead. MM-PBSA/GBSA and QM-PBSA/GBSA binding free energies strongly suggested a greater stability of the naringenin/DM- β -CD inclusion complex. Van der Waals force played an important role as the key guest–host interaction for the complexation between naringenin and each cyclodextrin.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

The polyphenols, and in particular flavonoids, are natural products found in many vegetables and fruits. Based on the variations in their polyphenolic structure, flavonoids can be divided into the different classes of flavonols, flavones, flavanones, catechins, anthocyanidins, isoflavones, dihydroflavonols and chalcones. Their broad pharmacological profiles include anti-lipoperoxidant and anti-inflammatory properties [1] and the ability to exert anticancer and chemopreventive activities [2]. Naringenin (5,7,4-trihydroxyflavanone or 5,7-dihydroxy-2-(4-hydroxyphenyl) chroman-4-one in Fig. 1A) belongs to the flavanones class of flavonoids. It is abundant in citrus fruits, such as grapefruit (*Citrus paradise*) and orange (*Citrus sinensis*). It exerts a variety of pharmacological effects [3] including antioxidant effects, blood lipid-lowering effects (Borradaile et al., 1999; Santos et al., 1999), anti-inflammatory activity through inhibition of the enzymes

involved in arachidonate metabolism [4–6], drug inhibition against cytochrome P-450 [7] and anti-carcinogenic effects [8,9]. In the cancer case, naringenin was reported to induce apoptosis in various cancer cell lines with no effect on normal cells [10]. Additionally, it has an insulin-like effect to decrease apolipoprotein B (Apo B) secretion in hepatocytes and to lower blood glucose levels in healthy male wistar rats [11]. From this point of view, there is a growing interest in applying naringenin in food products to enhance their health functions as well as its use in developing nutraceutical and pharmaceutical products. Unfortunately, a common limiting factor in the use of flavonoids in many applications is their low water solubility and stability. The formation of an inclusion complex with cyclodextrin (CD) is one effective way to solve this problem.

The cyclodextrins (CDs) are a group of cyclic oligosaccharides that consist of six (α -cyclodextrin or α -CD), seven (β -cyclodextrin or β -CD) or eight (γ -cyclodextrin or γ -CD) glucopyranose units linked by α -(1,4) glycosidic bonds. From their geometry, the internal cavity is highly hydrophobic, whilst the external hydroxyl face establishes a hydrophilic character. In the recent years, CD complexation has been successfully used to improve the solubility, stability toward light and oxidative degradation, dissolution rate, and bioavailability of poorly water soluble drugs [12–15]

* Corresponding author. Tel.: +66 2 2185426; fax: +66 22 185418.

E-mail addresses: t.rungrotmongkol@gmail.com, thanyada.r@chula.ac.th (T. Rungrotmongkol).




Fig. 1. Chemical structures of (A) naringenin and (B) the two CDs, β -CD and DM- β -CD, where -R is H- and -CH₃, respectively.

such as nimbin [16], crassicauline A [17] and alpinetin [18]. Since the application of β -CD in pharmaceutical fields is limited by its rather low aqueous solubility, more soluble derivatives of β -CD are presently of interest [12,13]. The amorphous CD, 2,6-dimethyl- β -cyclodextrin (DM- β -CD, Fig. 1B), is a chemically modified cyclomaltoheptaose (β -CD) derivative obtained by methylation of the 2- and 6-hydroxyl groups. Compared to the native β -CD, DM- β -CD exhibits an increased solubility in water and organic solvents [15] which enable its use as a carrier for poorly soluble molecules in water or hydrophobic solvents [19,20]. In addition, DM- β -CD was found to be an effective absorption promoter for nasal glucagon in rabbit [19] and has been reported in a European patent application for oral or parenteral administration [21]. To date, the solubility enhancement of naringenin by natural CD has been reported on from experimentally and theoretically derived studies, but only a few studies on the DM- β -CD derivative are available.

In the present work, the stability of naringenin was theoretically achieved by complexation with natural β -CD and DM- β -CD using multiple molecular dynamic (MD) simulations and free energy calculations. We aimed to obtain a better understanding of the structure and dynamic properties of naringenin inside the CD cavity. These results constitute new and valuable information on the naringenin binding strength, the key guest–host interaction and the binding free energy of complex formation.

2. Methods of computation

The starting configuration of naringenin was created and fully optimized with the *ab initio* calculation using the HF/6-31G* basis set in the Gaussian09 program [22]. The geometry of β -CD was taken from a systematic investigation on the structure of β -CD [23], and the structure of DM- β -CD was obtained by adding methyl groups to β -CD and subsequent geometry optimization with B3LYP/6-31G**. Prediction of a naringenin inclusion complex with these two CDs was performed *via* a docking procedure with 500 independent runs using the CDocker module of Accelrys Discovery Studio 2.5 (Accelrys, Inc.). The three docked inclusion complexes with the best ranked interaction energy and hydrogen bond (H-bond) formation between naringenin and each CD were then chosen for the MD study. The following system preparation and MD simulations on all inclusion complexes were performed with the Amber 12 software package [24]. The Glycam-06 bimolecular force field [25] was applied for the β -CD and DM- β -CD. In case of naringenin, the atomic charges and parameters were developed according to standard procedures [26–28]. The electrostatic potential (ESP) charges around the optimized naringenin molecule were evaluated by the HF/6-31G* calculation using Gaussian09. The restrained electrostatic potential (RESP) charges of

naringenin were then obtained by a charge fitting procedure using the antechamber module implemented in the Amber 12, while its parameters were obtained using the parmchk module. The hydrogen atoms of each inclusion complex were minimized with the 1000 steps of steepest descents (SD) followed by 3000 steps of conjugated gradients (CG) to relax the structure and release bad contacts. The complex was solvated by SPC water molecules with a spacing distance of 12 Å around the system surface. The naringenin/ β -CD and naringenin/DM- β -CD complexes consisted of 1480 ± 10 and 1750 ± 3 water molecules in the $41 \text{ \AA} \times 41 \text{ \AA} \times 41 \text{ \AA}$ and $44.0 \text{ \AA} \times 44.0 \text{ \AA} \times 44.0 \text{ \AA}$ truncated octahedron periodic boxes, respectively. The water molecules alone were minimized with the SD (1000 steps) and CG (3000 steps) and then minimization with the same process was applied on the whole system. In the next step, the MD simulation with periodic boundary condition using the NPT ensemble and a time step of 2 fs. The electrostatic interactions were taken into account by the particle mesh Ewald approach [29] with a cutoff distance of 12 Å. The SHAKE algorithm was used to constrain all bonds with hydrogen atoms. Each system was heated up to 298 K with a relaxation time of 100 ps and then simulated at the same temperature for 80 ns. The coordinates were recorded every 500 steps for analysis. The root means square displacement (RMSD), distance between the centers of gravity of the naringenin ring and CD ($d(Cg_{\text{chromone/phenyl ring}} - Cg_{\text{CD}})$) along the simulation time were analyzed by the ptraj module. The 30 MD trajectories of inclusion complex extracted from simulation at equilibrium (discussed later) were used for the binding free energy prediction. MM-PBSA and MM-GBSA approaches, the acceptable methods for predicting the absolute binding free energy of molecules in solution [30] were adopted for calculating the free energy of naringenin binding to β -CD and DM- β -CD. The methodology details of both methods have been described elsewhere [27,31–33]. To correct the MM energy, single point M06-2X/6-31+G** calculations were applied to the structures of the inclusion complex, CD and naringenin. Note that the DFT M06-2X functional energy includes the empirical dispersion correction energy [34], which is an important factor in the guest–host interaction.

3. Results and discussion

3.1. Naringenin/CD inclusion complex

From the 500 independent docking runs, the conformation, interaction energy and H-bond formation of naringenin inside the hydrophobic cavity of the CDs are given in Table 1.

Note that the interaction energy of inclusion complex was calculated by the energy difference between inclusion complex, cyclodextrin and naringenin using the CHARMM-based docking algorithm [35]. It can be clearly seen that the phenyl ring of naringenin molecule (Fig. 1A) preferentially dipped to β -CD (99% docked conformation) and DM- β -CD (100%) with a location close to the primary rim. The obtained information was in good agreement with the recent ¹H and 2D NMR study of naringenin in complex with β -CD and DM- β -CD [36] and the theoretical studies by 6-ns MD simulation [37] as well as the semi-empirical AM1 method [38] on naringenin/ β -CD complex.

Based on docked interaction energy and H-bond between naringenin and cyclodextrin, the best formed inclusion complex with β -CD and DM- β -CD were depicted in Fig. 2. It is worth noting that the DM- β -CD encapsulates this flavanones molecule better than the native β -CD by 2 kcal/mol in the gas phase. For naringenin/ β -CD complex (Fig. 2A), the hydroxyl oxygen (O7) on its chromone ring formed a H-bond with the 3-OH group located in the interior of β -CD (1.9 Å). Differentially, a weaker H-bond was detected in the naringenin/DM- β -CD complex (2.5 Å) between the O1 atom and

Download English Version:

<https://daneshyari.com/en/article/443339>

Download Persian Version:

<https://daneshyari.com/article/443339>

[Daneshyari.com](https://daneshyari.com)