



Evaluation of airborne Actinomycetes at waste application facilities

Abdel Hameed Awad^{1,2}, Safa A. El Gendy¹

¹ Air Pollution Department, National Research Centre, Dokki, Giza, Egypt

² Department of Environmental and Health Research, the Custodian of the Two Holy Mosques Institute for Hajj and Umrah Research, Umm Al Qura University, Saudi Arabia

ABSTRACT

This study aimed to evaluate airborne meso- and thermophilic actinomycete concentrations and their types at a wastewater treatment plant and a biosolid landfill, in Egypt. Air samples were collected at 200 m upwind, and onsite and 300 m downwind by using liquid impinger sampler, calibrated to draw 12.5 L/min, for 20 minutes. The concentrations ranged between 0.0–7360 CFU/m³ for mesophilic, and 106–586 CFU/m³ for thermophilic actinomycetes. Airborne actinomycete concentrations exceeded the suggested occupational exposure limit value of 100 CFU/m³. No significant correlations were found between actinomycete concentrations onsite and 300 m downwind distance. At the biosolid landfill, upwind thermophilic actinomycetes significantly differed ($p < 0.05$) from onsite and 300 m downwind. A total of 40 and 69 airborne actinomycete isolates belonging to 8 genera were identified at the wastewater treatment plant and biosolid landfill. *Streptomyces* were the dominant actinomycete species. *Streptomyces diastaticus*, *Pseudonocardia compacta* and *Catellatospora ferruginea* were only detected at the biosolid landfill site. Meso- and thermophilic actinomycetes positively associated with relative humidity, and wind speed positively correlated with onsite thermophilic actinomycetes ($r = 0.65$) at the biosolid landfill. Temperature showed negative effect on survivability of mesophilic actinomycetes ($r = -0.8$) onsite of the wastewater treatment plant. Waste application facilities increase actinomycete concentrations onsite which may consequently deteriorate air quality in the nearby areas.

Keywords: Air, actinomycetes species, biosolid landfill, wastewater treatment plant, meteorological conditions



Corresponding Author:

Awad A. Abdel Hameed

☎ : +966-56-991-3502

☎ : +966-25-56-4955

✉ : abed196498@yahoo.com

Article History:

Received: 29 April 2013

Revised: 14 September 2013

Accepted: 15 September 2013

doi: 10.5094/APR.2014.001

1. Introduction

Sewage treatment plants and biosolid landfills have been considered as potential sources of bioaerosols (Karra and Katsivela, 2007; Heinonen–Tanski et al., 2009). Aeration tanks in wastewater treatment plants are considered the main sources for bioaerosol emissions (Pascual et al., 2003; Li et al., 2013).

Actinomycetes are important bio-pollutants in occupational environments (Nielsen et al., 1997), and the major component of bioaerosols emitted from composting facilities (Lacey, 1997; Swan et al., 2003). Thermophilic actinomycetes have been known in moldy hay, compost and self heated substrates (Unaogu et al., 1994), and have been used as indicator for bioaerosols discharged from composts (Dutkiewicz, 1997). Several studies have been published on airborne actinomycete concentrations in the urban and occupational settings (Breza–Boruta and Paluszak, 2007; Fang et al., 2008), and have indicated that actinomycetes are distributed worldwide and found in soil, decaying organic material and wastes. In Egypt, Abdel Hameed (1996) found airborne actinomycetes at mean concentrations of 10² CFU/m³ at city center, and 10³ CFU/m³ at Zenein wastewater treatment plant. Airborne actinomycetes averaged 2.25x10⁴ CFU/m³, 2.9x10⁴ CFU/m³, and 4.1x10³ CFU/m³ at 20 m, 40 m, and 60 m downwind distances, respectively, of an agricultural non-point source during wheat harvesting season (Abdel Hameed and Khodr, 2001), and varied within 46–222 CFU/p/h along the main stream of the Nile river (Abdel Hameed et al., 2008). *Saccharopolyspora rectivirgula*, *Saccharomonospora* spp., *Thermoactinomyces thalophilus*, *Thermoactinomyces vulgaris* and *Thermomonospora* are the

common thermophilic actinomycete species, and *Streptomyces* are the common mesophilic species (Swan et al., 2003).

The exposure of waste workers to airborne bacteria and fungi vary depending on site location, type of waste, treatment technology, and meteorological conditions (Nielsen et al., 1998; Thorn and Kerekes, 2001). Residents living ~200 m away of a composting plant are exposed to bioaerosols and suffered from irritations (Herr et al., 2003). Actinomycete species are considered human pathogens (Taha et al., 2007), and prolonged inhalation of actinomycetes is linked to allergic alveolitis (Herr et al., 2004), as their spores can deeply penetrate into the lungs. The present study aims to evaluate airborne meso- and thermophilic actinomycete concentrations and species upwind, onsite and 300 m downwind, and to investigate their distribution patterns in association with meteorological factors, at two waste application facilities varied in waste type and location, a wastewater treatment plant and a biosolid landfill, in Egypt.

2. Materials and Methods

2.1. Description of the waste application facilities

Zenein municipal wastewater treatment plant and Shoubraiment biosolid landfill were selected for sample collection. The wastewater treatment plant is located in a suburban area of Giza governorate, Egypt ~7 km west of the Nile River. Zenein is a densely populated suburban area, characterized by various human and small industrial activities. The wastewater treatment plant capacity is 450 000 m³/day, the system has preliminary and

secondary settlement tanks, and aeration tanks. The biosolid landfill is ~1 714 acres located in the Shoubramant desert of the Giza governorate. It was constructed in 1989, and it is capable of holding at least ~30 years production of biosolid (domestic, commercial and soil refuses). The amounts of waste handled are ~70 000 tons/year. The refuse is spread and compacted to its maximum concentration by bulldozers.

2.2. Air sampling and analysis

The samples were collected 200 m upwind, onsite (zero point), and 300 m downwind. No-fixed sampling points were selected, as sampling points were chosen according to the prevailing wind direction (Table 1). Liquid impinger sampler (AGI-30, SKC) containing 40 mL sterilized phosphate buffered solution, held ~1.5–2 m above the ground level and 1–2 m far from the main source, was used to collect airborne actinomycetes. The air was aspirated at a flow rate of 12.5 L/min (manufacturer's recommended rate) for 20 min. At each sampling point, two consecutive samples were collected, two times per month, from June 2006 to 2007, between 9 am and 2 pm.

Aliquots (0.5 mL) of the original sample, and its serial dilutions (up to 10^{-2}) were spread-plated, in duplicate, onto two series of starch casein agar medium (Hi-media laboratories, Mumbai, India). The plates were incubated at 28 °C and 45 °C for 7–14 days to determine mesophilic and thermophilic actinomycetes, respectively. The resultant colonies were counted and expressed as colony forming unit per cubic meter of the air (CFU/m³).

2.3. Biosolid and wastewater microbial analysis

One sample of biosolid and sewage water was collected once every three months, a total of 4 samples. A 100 mg of biosolid material was dissolved in 50 mL buffer phosphate solution, and serial dilutions were prepared (up to 10^{-7}). Aliquots (0.1 mL) of the original and its serial dilutions were spread plated, in duplicate, onto the surface of starch casein agar media. Aliquots (0.1 mL) of the original wastewater (collected from aeration tank), and its serial dilutions (up to 10^{-3}) were examined for the presence of meso- and thermophilic actinomycetes. The plates were incubated at the same previously mentioned conditions. The resultant colonies were expressed as CFU/mg for the biosolid material, and CFU/mL for the wastewater.

2.4. Identification of actinomycetes

Identification of actinomycetes was carried out on the basis of morphological and biochemical features. The morphology of both spore chain and surface were examined (International Streptomyces Project, ISP) using Olympus microscopy (model CX 31, RBSF, Tokyo, Japan), and Scanning Electron Microscopy (model JEOL, JEM, Japan), respectively. Actinomycete isolates were screened on the basis of the color of colony, reverse side, and production of diffusible pigment (ISP 2, 3, 4 and 5), (Pridham and Lyons, 1961; Shirling and Gottlieb, 1966). Formation of melanin (ISP-6), (Trenser and Danga, 1958), utilization of carbon and nitrogen sources (Shirling and Gottlieb, 1966; Williams et al., 1989), producing of protease, lecithinase, and lipase enzymes (Nitsch and

Kutzner, 1969), catalase and hydrogen sulfide (Kuster and Williams, 1964), hydrolysis of pectin (Hankin et al., 1971), reduction of nitrate (Kutzner et al., 1978), degradation of tyrosine, xanthine, casein, cellulose, gelatin and starch (Jones, 1949), and resistant to different antibiotics (Goodfellow and Orchard, 1974) were tested.

2.5. Meteorological conditions

Ambient temperature and relative humidity were measured with a portable psychrometer (SATO; PC-5 000 TRH-II Sampler, China) during every sampling event. Wind speed records were obtained from the Egyptian Meteorological Authority. During the study period, temperature ranged from 15.5–37.5 °C, relative humidity 25.5–69%, and wind speed 1.85–5.75 m/s. The prevailing wind direction was north-south.

2.6. Statistical analysis

Due to non normal distribution of the analyzed variables, Spearman's rank correlation coefficient test was used to examine the relationships between airborne actinomycetes, and meteorological factors. The differences between airborne actinomycete concentrations at the different sampling points were analyzed by using student t-test and Mann Whitney-U-test. A probability of less or equal to $p \leq 0.05$ was considered significant.

3. Results

3.1. Overall concentrations

Airborne actinomycete concentrations fluctuated throughout the period of study with higher concentrations onsite than both upwind and 300 m downwind. In some months, the downwind concentrations exceeded the onsite ones (Figure 1). Airborne mesophilic actinomycetes ranged from 0.0–200 CFU/m³ upwind, and 0.0–7 360 CFU/m³ downwind at both waste facilities (Table 2). The highest mesophilic actinomycete concentrations were found in February at the wastewater treatment plant (Figure 1a), and March at the biosolid landfill (Figure 1b). Mesophilic actinomycete concentrations did not significantly differ ($t \leq 2$, $p > 0.05$) between the all wastewater treatment plant sampling points. However mesophilic actinomycete concentrations were significantly higher at 300 m downwind than upwind ($t = 2.7$, $p \leq 0.05$) at the biosolid landfill. Weak correlations were found between onsite and 300 m mesophilic actinomycete concentrations at the wastewater treatment plant ($r = 0.26$) and the biosolid landfill ($r = -0.12$).

At both waste facilities, airborne thermophilic actinomycete concentrations ranged from 0.0–70 CFU/m³ upwind, and 0.0–5 973 CFU/m³ downwind sampling points (Table 2). The greatest onsite and 300 m thermophilic actinomycete concentrations were found in January and November at the wastewater treatment plant (Figure 2a) and the biosolid landfill (Figure 2b), respectively. Upwind concentrations significantly differed ($p \leq 0.05$) from onsite at the wastewater treatment plant, and with both onsite and 300 m downwind sampling points at the biosolid landfill. Positive correlation ($r = 0.54$) was found between thermophilic actinomycetes detected onsite and 300 m downwind at the biosolid landfill.

Table 1. Description of sampling points at both waste application facilities

Location	Description
Zenin wastewater treatment plant	Onsite (zero point); it is located ~1–2 m adjacent to the aeration tanks of the first stage of the wastewater treatment plant, and it is considered the source of microbial emission.
	300 m downwind distance of aeration tanks, it is located at the surrounding residential areas outside the borders of the facility. Background, 200 upwind of the wastewater treatment plant, and it is mainly located outside the borders of the plant.
Shoubramant biosolid landfill	Onsite (zero point), it is located ~1–2 m adjacent to the fresh biosolid pile.
	300 m downwind distance of the main fresh biosolid pile. Background, 200 m upwind of the biosolid landfill according to wind direction.

Download English Version:

<https://daneshyari.com/en/article/4434727>

Download Persian Version:

<https://daneshyari.com/article/4434727>

[Daneshyari.com](https://daneshyari.com)