



ELSEVIER

Contents lists available at ScienceDirect

Environmental Research

journal homepage: www.elsevier.com/locate/envres

Reports from the Field

Arsenic occurrence in drinking water supply systems in ten municipalities in Vojvodina Region, Serbia [☆]

Dragana Jovanovic ^{a,*}, Branko Jakovljević ^b, Zorica Rašić-Milutinović ^c, Katarina Paunović ^b, Gordana Peković ^c, Tanja Knezević ^a

^a Institute of Public Health of Serbia “Dr Milan Jovanovic Batut”, Dr Subotica 5, Belgrade, Serbia

^b Institute of Hygiene and Medical Ecology, School of Medicine, Belgrade, Serbia

^c Department of Endocrinology, University Hospital Zemun, Belgrade, Serbia

ARTICLE INFO

Article history:

Received 9 August 2010

Received in revised form

15 November 2010

Accepted 30 November 2010

Available online 24 December 2010

Keywords:

Arsenic

Drinking water

Public water supply

Pannonian Basin

Vojvodina

ABSTRACT

Vojvodina, a northern region of Serbia, belongs to the Pannonian Basin, whose aquifers contain high concentrations of arsenic. This study represents arsenic levels in drinking water in ten municipalities in Serbia. Around 63% of all water samples exceeded Serbian and European standards for arsenic in drinking water. Large variations in arsenic were observed among supply systems. Arsenic concentrations in public water supply systems in Vojvodina were much higher than in other countries in the Pannonian Basin.

© 2010 Elsevier Inc. All rights reserved.

1. Introduction

The Ministry of Health of the Republic of Serbia has implemented a national plan entitled “Monitoring Programme of Drinking Water Quality from Public Water Supply Systems” in 1978. It obliged public water supply systems and public health institutes to perform routine monitoring over the quality of drinking water in Serbia. One of its goals was to estimate inorganic arsenic levels in groundwater sources used for public water supply.

Vojvodina is a northern region of Serbia, geographically belonging to the southern part of the Pannonian Basin, together with Hungary, Romania, Slovakia and Slavonia in Croatia. The Pannonian Basin and the greater part of the Balkan Peninsula in south-east Europe belong to the Danube River Basin, which is geologically and geochemically complex (Dangic, 2007). Quaternary sedimentary aquifers within the Pannonian Basin contain high concentrations of naturally occurring arsenic (Varsanyi and Kovacs, 2006). In the water-sediment system, arsenic is specially enriched in clay fraction of the soil; tertiary magmatic rocks contain 3.1 mg As/kg, whereas soils contain 10 mg As/kg (Dangic, 2007). On the other

side, arsenic concentrations are lower in the Danube River water and shallow riparian groundwaters (0.1–8 µg/L) (Dangic, 2007).

Other sources of arsenic include foodstuffs (primarily fish and marine organisms), air and soil, and some human activities, such as the use of arsenical pesticides, burning of fossil fuels and disposal of industrial wastes (Mandal and Suzuki, 2002). Exposure to inorganic arsenic as arsenite (As(III)) or arsenate (As(V)) occurs from water and dust particles, whereas fish and seafood predominantly contain organic arsenic compounds (Francesconi and Kuehnelt, 2002). Arsenic concentrations in different types of soil in urban areas in Serbia range from 6 to 10 mg/kg (Ministry of Environment and Spatial Planning of Serbia, 2009). Average annual emission of arsenic in atmospheric total suspended particles in several urban zones in Serbia ranges from 2 to 18.6 ng/m³ (Institute of Public Health of Serbia, 2009).

The European Food Safety Authority Panel on Contaminants in the Food Chain (CONTAM) has in 2009 estimated national inorganic arsenic exposures from food and water across 19 European countries. Based on their report, this exposure ranged from 0.13 to 0.56 µg/kg of body weight per day for average consumers and from 0.37 to 1.22 µg/kg of body weight per day for 95th percentile consumers (EFSA Panel, 2009).

Although arsenic is a known water pollutant in Vojvodina for many decades, public water supply systems lack technical and financial resources for its removal. Since arsenic does not change water color, odor or taste, exposed populations may continue to use the contaminated water for drinking, cooking and soil/garden/plants watering all their lives, thus ingesting large amounts of

Abbreviations: As, arsenic; CONTAM, the European Food Safety Authority Panel on Contaminants in the Food Chain; SRPS ISO, Serbian Standardization Organization and International Standardization Organization

[☆] This work was performed at the Institute of Public Health of Serbia “Dr Milan Jovanovic Batut”, Belgrade, Serbia

* Corresponding author.

E-mail address: dragana_jovanovic@batut.org.rs (D. Jovanovic).

arsenic. Knowing that arsenic causes skin, bladder, and internal cancers (Abernathy et al., 1999), its presence in water represents a major public health topic. In the nearest future, Serbia will implement many European Union Directives regarding water quality and will have to find means to treat this problem. The aim of this study was to report current distribution of arsenic in drinking water from public water supply systems in ten municipalities in Vojvodina Region, Serbia.

2. Materials and methods

Vojvodina Region administratively consists of 45 municipalities. National Monitoring Programme of Drinking Water Quality in Serbia shows that public water supply systems in 20 municipalities (44.4% of all municipalities in Vojvodina) contain high amounts of arsenic. Regional authorities in these 20 municipalities were invited to participate in the study by collecting and analyzing arsenic in their water supply systems in 2008. Those who failed to respond were invited again in 2009. The response rate was 50%, i.e. 10 municipalities provided all data on time. Other municipalities were excluded from the study because of their refusal or lack of financial resources to analyze arsenic (7 municipalities), or because of inadequate sampling or analysis procedures (3 municipalities).

Total arsenic concentrations were measured in water samples from public water supply systems during 2008 and 2009. No statistical difference was observed between the samples taken in various years and therefore the most recent data were included in the analysis. Water samples from individual wells or bottled water were not taken into consideration.

Sampling and analyses of drinking water were performed at the following laboratories: at the Institute of Public Health of Vojvodina (in Novi Sad), the Institute of Public Health in Subotica, the Institute of Public Health in Zrenjanin and at the Institute of Public Health of Serbia in Belgrade. All laboratories were accredited and authorized according to SRPS ISO/IEC 17025 and SRPS ISO 9001 standards. Laboratory procedures for sample management, analytical methods and quality control measures (accuracy, precision and detection limits) were standardized by Serbian law (Book of Regulations on the Hygienic Correctness of Drinking Water 98/42, 1998). Current Serbian regulations limit arsenic levels at 10 µg/L (Book of Regulations on the Hygienic Correctness of Drinking Water 98/42, 1998).

Total arsenic water was determined using the HG-AAS technique (Perkin-Elmer 1100 atomic absorption spectrometer equipped with a MHS-20 hydride generation system, Perkin-Elmer Corp., Norwalk, CT, USA). Reagents of analytical grade or higher quality were used. Hydride generation was performed using a 3% (w/v) NaBH₄ in 1% NaOH (both Merck suprapur; E. Merck, Darmstadt, Germany) solution. The radiation source was a hollow cathode lamp of arsenic (Perkin-Elmer) used at a wavelength of 193.7 nm and a spectral slit width of 0.7 nm. Hydride generation (HG) was performed with 0.6% NaBH₄ dissolved in 0.5% NaOH and 5 mol/dm³ HCl. The hydrides were then transported to a heated quartz cell and the atomic absorption of the analyte was measured. With this method a limit of detection of 0.5 µ/L and limit of quantification of 2 µ/L were obtained by analyzing 4 series of 10 repeated analyses of blank samples and calculating the three standard deviations of these responses. Recovery of standards was 80–120%.

Descriptive statistic was presented as median values, 10th and 90th percentiles for numeric variables and percents for categorical variables. Differences between several samples were analyzed using Kruskal Wallis test for non-parametric data. STATISTICA software was used for all data analyses (Version 6, StatSoft Inc., Tulsa, OK, USA).

3. Results

In total, 577 water samples from public water supply systems from ten municipalities in Vojvodina were analyzed for arsenic. As shown in Table 1, almost 63% of all samples exceeded the current standard. Srbobran and Subotica municipalities (in the north-west of Vojvodina) had the lowest number of samples that exceeded the arsenic levels. Arsenic was detected in all water samples from Senta and Kanjiza (north-east of Vojvodina), and in 94% of samples in Zrenjanin (eastern part of Vojvodina) (Table 1).

Geographical distribution of arsenic in the study area and in the whole region of Vojvodina is shown in Fig. 1. High levels of arsenic are present in ten investigated municipalities (red circles) and are known to be present in ten more municipalities (yellow squares), but more precise data were unavailable at the present.

The distribution of total arsenic in drinking water by public water supply systems in the investigated municipalities is shown in Table 2.

Table 1

Number of analyzed samples, number and percent of samples with exceeded arsenic levels in public water supply systems in Vojvodina Region, according to Serbian maximum allowed value.^a

Municipality	No. of analyzed samples	No. of samples with exceeded ^a As levels	% of samples with exceeded ^a As levels
Zrenjanin	144	135	93.8
Bačka Palanka	174	126	72.4
Bački Petrovac	60	24	40.0
Bač	8	3	37.5
Srbobran	42	3	7.1
Titel	18	16	88.9
Temerin	55	27	49.1
Subotica	52	5	9.6
Kanjiza	12	12	100.0
Senta	12	12	100.0
Total	577	363	62.9

As—arsenic.

^a Over 10 µg/L.




Fig. 1. Geographical distribution of municipalities in Vojvodina according to arsenic levels. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

Large variations in arsenic concentration were observed among water supply systems (median ranged from 2 to 250 µg/L; maximum value ranged from 5 to 349 µg/L). The highest concentrations were reported in the city of Zrenjanin and most of its villages (eastern part of Vojvodina), in municipalities Temerin (central to north-east), Subotica (north) and village Čelarevo (west).

Fig. 2 represents the distribution of mean arsenic concentrations in water supply systems. In total, only 11% of all water supply

Download English Version:

<https://daneshyari.com/en/article/4470247>

Download Persian Version:

<https://daneshyari.com/article/4470247>

[Daneshyari.com](https://daneshyari.com)