



Activity of natural compounds from peanut skins on *Fusarium verticillioides* growth and fumonisin B₁ production

Romina P. Pizzolitto^{a,b,*}, José S. Dambolena^a, María P. Zunino^a, Mariana Larrauri^c, Nelson R. Grosso^c, Valeria Nepote^c, Ana M. Dalcero^b, Julio A. Zygodlo^a

^a Instituto Multidisciplinario de Biología Vegetal (IMBiV-CONICET), Cátedra de Química Orgánica, Facultad de Ciencias Exactas Físicas y Naturales, Universidad Nacional de Córdoba, Córdoba, Argentina

^b Departamento Microbiología e Inmunología, Cátedra de Micología, Facultad de Ciencias Exactas Físico-Químicas y Naturales, Universidad Nacional de Río Cuarto, Río Cuarto, Córdoba, Argentina

^c IMBiV (CONICET), Cátedra de Química Biológica, Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba, Córdoba, Argentina

ARTICLE INFO

Article history:

Received 25 October 2012

Received in revised form 8 March 2013

Accepted 19 March 2013

Keywords:

Peanut skin

Antifungal activity

Fusarium verticillioides

Fumonisin B₁

By-products

ABSTRACT

The objectives of this investigation were to evaluate the antifungal and antimycotoxigenic properties of peanut skin extracts (PSE) against *Fusarium verticillioides*. The PSE were prepared by a multisolvent extraction procedure, and the activity on growth parameters and fumonisin B₁ (FB₁) production of the three PSE was explored at different concentrations on potato dextrose agar and in artificially infected maize kernels, respectively. The results demonstrated that all PSE had a significant influence on growth rate or lag phase of *F. verticillioides*. The yellow and purple (250–500 µg ml⁻¹) extracts decreased growth rate, whereas brown extract extended the lag phase. Only the yellow extract at 62.5 µg ml⁻¹ was able to affect both growth rate and lag phase. With respect to mycotoxin production a significant stimulation on FB₁ production was observed with purple (62.5 µg ml⁻¹) and brown (250 µg ml⁻¹) extracts. In contrast, a significant decreased in FB₁ was observed at 62.5 µg ml⁻¹ of yellow extract. These findings showed that natural compounds from PSE possess inhibitory effects on *F. verticillioides* growth and mycotoxin production. Thus, PSE could be use as an alternative to minimize fungal contamination.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

Fusarium is a large complex genus with species adapted to a wide range of habitats. They are worldwide in distribution and many are important plant pathogens. However, many species are soil borne and exist as saprophytes important in breaking down plant residues. Fumonisin is a fungal secondary metabolite produced by species of *Fusarium*, mainly *F. verticillioides* and *F. proliferatum* (Krska et al., 2007). There are several identified fumonisins, but fumonisin B₁ (FB₁) and fumonisin B₂ (FB₂) are the most important and constitute up to 70% of the fumonisins found in naturally contaminated foods (Niderkorn et al., 2009). Fumonisin B₁ and FB₂ are phytotoxic to corn (Lamprecht et al., 1994), cytotoxic to various mammalian cell lines (Abbas et al., 1993) and FB₁ is carcinogenic in rat liver and kidney (IARC, 2002). The occurrence of these analogs in home-grown corn has been associated with an

increased risk of esophageal cancer in humans (Shephard et al., 2000). Fumonisin B₁ is the most toxic fumonisin analog and it is considered possible carcinogens to human and classified as class 2B (IARC, 2002). Considering the high incidence of *Fusarium* species on crops and the impact of fumonisin on human and animal health, the application of strategies to prevent their formation in foods, as well as, to eliminate, inactivate or reduce their presence in food products, becomes necessary. For many years, synthetic fungicides have been used for control plant pathogenic fungi. However, the extensive use of these chemicals led to the development of resistance in many areas around the world and also increases the risk of toxic residues in the products (Marei et al., 2012). Thus, the exploitation of natural substances with bioactivity against fungi has been the target of interest in the search for ecologically safe products (Dambolena et al., 2012). Agricultural wastes represent a largely ignored source of high-value phytochemicals and value-added industrial products that could contribute to sustainability objectives (Das and Singh, 2004). Peanut skin is a by-product of the peanut blanching operation that has low economic value despite its high content of active components including flavonoids, phenolic acids, phytosterols, alkaloids, and stilbenes. Some therapeutic effects have been reported for peanut seed extracts, such as antioxidative, antibacterial, antifungal, and anti-inflammatory activities

* Corresponding author at: Departamento Microbiología e Inmunología, Facultad de Ciencias Exactas, Físico-Químicas y Naturales, Universidad Nacional de Río Cuarto, Ruta 36 Km. 601, 5800 Río Cuarto, Córdoba, Argentina.
Tel.: +54 358 4676421; fax: +54 358 4676232.

E-mail address: rpizzolitto@exa.unrc.edu.ar (R.P. Pizzolitto).

(Lopes et al., 2011; Yu et al., 2005). In previous studies peanut skin compounds have been reported as antimicrobial agents (Sarnoski et al., 2012); however, to our best knowledge there is no information of the effects of peanut skin extracts (PSE) on FB_1 production and fungal growth. Thus, the objective of this investigation was to evaluate the antifungal and antifumonisin property of PSE against *Fusarium verticillioides*.

2. Materials and methods

2.1. Materials

Peanut skins from Argentine peanuts (*Arachis hypogaea*) (cv Runner, 2010 crop year) were obtained by a blanching process and were provided for the Company “Lorenzati, Ruesch y Cia”, Ticino, Córdoba, Argentina in March, 2010. The peanut skins were kept in a sealed plastic bag and stored at 4 °C until used.

2.2. Peanut skin extraction

The ethanolic extract (EE) from peanut skin was obtained according to the method published by Nepote et al. (2004) with slight modifications. Briefly, to obtain the EE, the peanut skins were previously defatted by two extractions with n-hexane (50 ml each 10 g peanut skins) during 12 h each one at room temperature. The dry defatted peanut skins (20 g) were extracted by solid–liquid extraction using ethanol 70%, during 24 h by maceration in darkness at room temperature. The extract was filtered and the residue was extracted again under the same conditions. The combined filtrate was evaporated to dryness in a rotary evaporator at 35 °C.

2.3. Separation of the ethanolic extract

The crude EE was purified by partition with 90 ml hexane, 300 ml ethyl acetate (EtOAc) and 55 ml water. The EtOAc fraction was evaporated in a rotary evaporator and separated with ethanol in a column packed with Sephadex LH-20 (internal diameter: 10 mm, length: 33.5 cm, elution flow 0.5 ± 0.05 ml min⁻¹). Fractions with different colors in visible light were separated from the column and identified as: yellow, purple and brown.

2.4. Fungal culture

The culture of *F. verticillioides* M3125, isolated from maize in California is a fumonisin-producing strain (Leslie et al., 1992). The strain obtained from carnation leaves-agar by monosporic isolation, was used in all experiments.

2.5. Antifungal activity of peanut skin extracts

Inoculum was prepared by growing on PDA agar for 7 days at 25 °C to obtained heavily sporulating cultures. A conidial suspension was placed in aqueous solution, after homogenizing, the suspension was counted using a Neubauer chamber and adjusted to 10^6 conidia ml⁻¹.

The PSE were tested at different concentrations of 500, 250, 125 and 62.5 µg ml⁻¹ of PDA agar to control growth of *F. verticillioides* M 3125. Peanut skin extracts were dissolved in 70% ethyl alcohol and added to the autoclaved based medium. PDA plates were inoculated centrally with 10 µl of the conidia suspension and were incubated for 8 days at 25 °C.

The radial mycelial growth was determined by periodical measurement of two right-angled diameters of the colonies. Colony diameters versus time were plotted and radial growth rates (mm day⁻¹) were evaluated from the slope by linear regression.

Lag phase was determined as the abscissa from growth rate curves. All the experiments were performed in triplicate.

2.6. Effect of peanut skin extracts on FB_1 biosynthesis on maize

Fumonisin B₁ biosynthesis was determined by using healthy maize as substratum. Corn grain free from FB_1 (25 g), was placed in 250 ml dark Erlenmeyer flasks and sterilized for 2 consecutive days in an autoclave for 15 min at 121 °C. The PSE first dissolved in 70% ethyl alcohol and then mixed with water. Water solution (8 ml) was added on autoclaved maize in order to reach 35% humidity. The used concentrations were 62.5 and 250 µg ml⁻¹. Maize was then inoculated with 50 µl of a conidial suspension (10^6 conidia ml⁻¹) of *F. verticillioides* prepared as described above. Immediately, in order to obtain good homogenization, flasks with inoculated maize were shaken vigorously. Treatments were incubated 28 days in dark at 25 °C, with manual stirring the first 5 days. Control flasks were prepared following the same procedure; however, no PSE were added on water. Four replications of each treatment were done.

2.7. FB_1 quantification

Briefly, after incubation, fermented maize was sterilized in an autoclave for 15 min at 121 °C and dried in a vacuum oven at 60 °C until constant weight was achieved. Later, 10 g of dried maize was finely ground. The FB_1 was extracted with ultrapure water by shaking the powder and water for 2 h in an orbital shaker. The aqueous extracts were centrifuged at $9000 \times g$, and filtered through filter paper (Whatman no. 4, Whatman International, Maidstone, UK). Samples (500 µl) from the aqueous extracts were diluted with acetonitrile (500 µl). The quantification of the diluted extracts was performed following the methodology proposed by Shephard et al. (1990). Briefly, an aliquot (50 µl) of this solution was derivatized with 200 µl of o-phthalaldehyde. This solution was obtained by adding 5 ml of 0.1 M sodium tetraborate and 50 ml of 2-mercaptoethanol to 1 ml of methanol containing 40 mg of o-phthalaldehyde. The derivatized samples were analyzed by Hewlett Packard HPLC equipped with a fluorescence detector. The wavelengths used were 335 and 440 nm for excitation and emission, respectively. An analytical reversed-phase column C18 (150 mm × 4.6 mm internal diameter and 5 mm particle size) was connected to a precolumn C18 (20 mm × 4.6 mm and 5 mm particle size). The mobile phase was methanol, NaH₂PO₄ 0.1 M (75:25); the pH was set at 3.35 ± 0.2 with orthophosphoric acid, and a flow rate of 1.5 ml min⁻¹. The quantification of FB_1 was carried out by comparing the peak areas obtained from aqueous extracts with those corresponding to the standards of 10.544, 5.135 and 2.567 µg ml⁻¹ of FB_1 (PROMEC, Programme on Mycotoxins and Experimental Carcinogenesis, Tygerberg, Republic of South Africa).

2.8. Statistical evaluation

Statistical analyses were conducted using INFOSTAT/Professional 2005p.1 (F.C.A.-Universidad Nacional de Córdoba, Argentina) at $p=0.05$. Data from these studies were analyzed through two-way analysis of variance (ANOVA). Normality of data was tested using the Shapiro–Wilk test. Comparisons between treatments were performed by the DGC (Di Rienzo, Guzmán and Casanoves) test (Di Rienzo et al., 2002). Results giving P values <0.05 were considered significantly different.

3. Results and discussion

The effect of PSE on *F. verticillioides* M3125 growth was evaluated in a range of concentrations between 0 and 500 µg ml⁻¹. The

Download English Version:

<https://daneshyari.com/en/article/4513592>

Download Persian Version:

<https://daneshyari.com/article/4513592>

[Daneshyari.com](https://daneshyari.com)